




**Review of Environmental Factors-  
Addendum  
Hunter-Central Coast REZ  
Network Infrastructure**

Appendix D- Ecology

**Wired for good.™**





**Updated Ecological Assessment Report  
with modifications  
for  
Ausgrids  
HUNTER REZ PROJECT  
NSW**



Prepared by: **WILDTHING** Environmental Consultants

38c Stapleton Street  
WALLSEND NSW 2287  
ABN: 41 033 509 215

For: **Ausgrid**  
145 Newcastle Rd,  
Wallsend NSW 2287

Job No: 12875

September 2026



|                |                                                                                               |                                                                                      |
|----------------|-----------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------|
| Project Name   | Updated Ecological Assessment Report with modifications for Ausgrids Hunter REZ Project, NSW. |                                                                                      |
| Project Number | 12875                                                                                         |                                                                                      |
| Prepared By    | Daryl Harman<br>Senior Ecologist BEnvSc<br>BAAS17074                                          |   |
|                | Nicola Mohr<br>Ecologist BScMSc<br>BAAS23007                                                  |  |
| Reviewed By    | Dr Kylie Bridges<br>Ecologist BEnvSc Hons PhD<br>BAAS20005                                    |  |
| Date           | Updated 16 September 2026                                                                     |                                                                                      |
| Version Number | 6                                                                                             |                                                                                      |

## Disclaimer

This report has been prepared in accordance with the proposal provided by the Client and outlined within this report. All findings, conclusions or recommendations contained within this report are based upon the data and results collected under the times and conditions specified in the report and are only applicable for the proposal considered within this report. This report has been prepared for use exclusively by the Client. No responsibility for its use by any other party is accepted by WILDTHING Environmental Consultants.

## Summary

Wildthing Environmental Consultants have been commissioned by Ausgrid to undertake an updated ecological assessment to reflect the proposed eleven (11) modifications to the original network upgrade of the existing electricity network between Kurri Kurri and Muswellbrook as part of Hunter-Central Coast Renewable Energy Project (REZ).

### The REZ proposal will be completed in Stages:

**Stage 1:** A proposed dual circuit high-capacity 132kV subtransmission lines ~60km between the existing Singleton and Kurri Sub-Transmission Substations (STS's). This will include pole replacements, access track works and vegetation clearing. This stage also includes upgrades at Singleton and Kurri STSs. The 132kV subtransmission line will be almost entirely located within Ausgrids existing maintained easement. Depending on the width of the required easement (either 20 or 45m) some areas of native vegetation along the edge of the easement will be impacted. In most cases only minor branch trimming will be required. Sections of new easement are also proposed which will require the removal of native vegetation in some areas.

### **Stage 2:**

A proposed dual circuit high-capacity 132kV subtransmission line (~25km) between the proposed Eastern Hub 132kV Switching Station located on the northern side of Lake Liddell to the existing Singleton STS.

Rearrangement of Ausgrid's existing Upper Hunter subtransmission network to provide connection between an existing subtransmission line known as 95U, which ultimately will connect to Eastern Hub SS and Ausgrid's existing Muswellbrook STS. The subtransmission line (~5km) will be almost entirely located within Ausgrids existing maintained easement. Some branch trimming and the removal of a small number of trees will likely be required.

### **Modifications to the original network upgrade**

A total of eleven modifications to the original network upgrade have been proposed. These modifications are shown below in Table 1.

Table 1: Proposed modifications to the Renewable Energy Project (REZ).

| No. | Modification Name                                             | Description/ Justification                                                                                                                                                                                                                                                                                                                      |
|-----|---------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1   | Hunter Expressway crossing – Branxton                         | The line deviates further to the south to cross the Hunter Expressway before turning back to meet the original alignment (~500m). This modification addresses some significant constructability issues with the original route.                                                                                                                 |
| 2   | ARTC crossing off Standen Drive, Belford                      | There is an existing building within the easement of the original route. the proposed route modification relocates the ARTC crossing approximately 200m west of the existing crossing thereby avoiding this interaction. In addition, the revised crossing is also more perpendicular to the railway line which is a preferred outcome for ARTC |
| 3   | Pothana Lane, Belford                                         | The proposed modification diverts north through the TSR for approximately 500m to the New England Highway and then heads west for approximately 1.9km before rejoining the original route.                                                                                                                                                      |
| 4   | Golden Highway, Whittingham                                   | The line diverts to the south of the original route for approximately 1km before rejoining in the original route due to detailed design.                                                                                                                                                                                                        |
| 5   | Mitchell Line of Road - Singleton Military Area 1 (Dochra)    | Relocation of existing 66kV line underground and revision of the 132KV OH to achieve height compliance with Defence requirements re Dochra Air Field.                                                                                                                                                                                           |
| 6   | Mitchell Line of Road - Singleton Military Area 2             | The proposed modification of approximately 1.3km diverts immediately to the south due to detailed design.                                                                                                                                                                                                                                       |
| 7   | Mitchell Line of Road - Singleton Military Area 3             | Approximately 600m diverts to the south to stay on the southern side of Mitchell line of road and avoid two road crossings.                                                                                                                                                                                                                     |
| 8   | Mitchell Line of Road - Singleton Military Area 4 (Redgroves) | The line diverts to the north along Mitchell Line of Road for approximately 2km to move out of the existing easement in Singleton Military Area.                                                                                                                                                                                                |



| No. | Modification Name                | Description/ Justification                                                                                                                                                                                                                                                                                                           |
|-----|----------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 9   | Jerry's Plains Road - Mt Thorley | Relocation from existing Ausgrid easement to west to the Jerry's Plains Road for approximately 3.5km before rejoining the original route where it crosses the highway at Gouldsville. This option removes two railway line crossings and reduces impact on Yancoal buffer lands. Also included is a minor realignment on Putty Road. |
| 10  | Glendell Mine, Ravensworth       | Minor realignment to immediately to east of existing easement for approximately 1 km due to inaccessibility of existing easement for construction and future maintenance works.                                                                                                                                                      |
| 11  | Antiene STSS – Hebden            | Minor realignment of southern connection to Antiene STSS due to detailed design and constructability issues. Addition of new northern connection approximately 1km.                                                                                                                                                                  |

### Vegetation

Vegetation within the REZ Project area ranged from highly disturbed land that had been historically cleared and subject to prolonged grazing and coal mining impacts to areas of woodland and open forest that were relatively intact. Most of the vegetation within the required 132kV subtransmission lines easement was contained within the existing Ausgrid easements, and had been kept low to the ground by routine maintenance. Other areas impacted by the Muswellbrook STS upgrade and new sections of easement such as an area between the Kurri STS and Hunter Expressway will require the removal of native vegetation in the form of woodland and open forest.

The vegetation assemblages along the REZ were stratified where possible by assigning the vegetation to Plant Community Types (PCTs) detailed in the NSW Vegetation Information System (VIS) classification database. A total of 11 Plant Community Types were identified: these were:

- PCT 3631 Kurri Sand Clay Woodland
- PCT 3433 Hunter Coast Foothills Spotted Gum-Ironbark Forest
- PCT 3634 Quorrobolong Sand Flats Forest
- PCT 3444 Lower Hunter Spotted Gum-Ironbark Forest
- PCT 3446 Lower North Foothills Spotted Gum-Ironbark Grassy Forest
- PCT 4023 Coastal Valleys Riparian Forest
- PCT 3315 Central Hunter Ironbark-Spotted Gum Forest
- PCT 4015 Central Hunter Swamp Oak Riparian Forest
- PCT 4073 Lower North Hinterland River Oak Forest
- PCT 3431 Central Hunter Ironbark Grassy Woodland
- PCT 3525 Upper Hunter Box-Blakely's Red Gum Grassy Forest

### Threatened Ecological Communities listed under the state BC Act 2016

Seven TEC's listed under the BC Act were identified within the REZ or in close proximity. These TEC's were:

- Kurri Sand Swamp Woodland in the Sydney Basin Bioregion
- Hunter Lowland Redgum Forest in the Sydney Basin and New South Wales North Coast Bioregions
- Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW North Coast Bioregions
- Central Hunter Ironbark – Spotted Gum – Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions
- Central Hunter Grey Box – Ironbark Woodland in the NSW North Coast and Sydney Basin Bioregions
- Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions
- White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grasslands

Table 2 shows impacted areas of each impacted PCT and aligned TEC's.



**Table 2: Areas of impacted vegetation within REZ including PCT's and aligned TEC's.**

| PCT                                                                | Condition                                                               | BC Act 2016                                                                                                              | EPBC Act 1999                                                                            | Total Area (ha) |
|--------------------------------------------------------------------|-------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------|-----------------|
| PCT 3631 Kurri Sand Clay Woodland                                  | Forested                                                                | EEC - Kurri Sand Swamp Woodland in the Sydney Basin Bioregion                                                            | EEC - Kurri sand swamp woodland of the Sydney Basin bioregion                            | 0.86            |
| PCT 3433 Hunter Coast Foothills Spotted Gum-Ironbark Grassy Forest | Forested                                                                | EEC - Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW North Coast Bioregions                        |                                                                                          | 0.31            |
| PCT 3444 Lower Hunter Spotted Gum-Ironbark Forest                  | Forested                                                                | EEC - Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW North Coast Bioregions                        |                                                                                          | 0.22            |
| PCT 4023 Coastal Valley Riparian Forest                            | Forested                                                                | EEC - Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions      | EEC- Swamp Oak Floodplain Forest                                                         | 0.08            |
| PCT 3315 Central Hunter Ironbark-Spotted Gum Forest                | Forested                                                                | EEC - Central Hunter Ironbark-Spotted Gum-Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions | Central Hunter Valley Eucalypt Forest and Woodland                                       | 13.24           |
| PCT 4015 Central Hunter Swamp Oak Riparian Forest                  | Forested                                                                | Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions            | Swamp Oak Floodplain Forest                                                              | 0.1             |
| PCT 4073 Lower North Hinterland River Oak Forest                   | Forested                                                                | EEC - Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions      |                                                                                          | 0.73            |
| PCT 3431 Central Hunter Ironbark Grassy Woodland                   | Forested                                                                | EEC - Central Hunter Grey Box-Ironbark Woodland in the New South Wales North Coast and Sydney Basin Bioregions           | Central Hunter Valley Eucalypt Forest and Woodland                                       | 4.61            |
| PCT 3525 Upper Hunter Box-Blakelys Red Gum Grassy Forest           | Forested                                                                | White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grasslands                                 | White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grasslands | 0.59            |
| PCT 3634 Quorrobolong Sand Flats Forest                            | Forested                                                                |                                                                                                                          |                                                                                          | 0.17            |
| Introduced <i>Juncus acutus</i>                                    |                                                                         |                                                                                                                          |                                                                                          | 0.46            |
| Native Regeneration                                                |                                                                         |                                                                                                                          |                                                                                          | 6.68            |
| Planted Native Trees                                               |                                                                         |                                                                                                                          |                                                                                          | 0.55            |
| Planted native vegetation                                          |                                                                         |                                                                                                                          |                                                                                          | 0.49            |
|                                                                    | Total native vegetation impacted by proposal outside managed easements. |                                                                                                                          |                                                                                          | 31.98           |
|                                                                    | Total native vegetation within existing easements (under management).   |                                                                                                                          |                                                                                          | 147.88          |

The proposal will result in an incremental reduction in narrow sections of these threatened ecological communities, particularly the EEC Central Hunter Ironbark – Spotted Gum – Grey Box Forest; however, it is considered unlikely to have a significant impact which might lead to extinction locally.

#### Threatened flora species listed under the state BC Act 2016

Four threatened flora species were recorded during fieldwork, these were

- *Eucalyptus glaucina* (Slaty Red Gum)
- *Eucalyptus parramattensis* subsp. *decadens* (Drooping Red Gum)
- *Grevillea parviflora* subsp. *parviflora* (Small-flowered Grevillea)
- *Diuris tricolor* (Pine Donkey Orchid)

The proposal will directly impact a small number of specimens of Slaty Red Gum and Drooping Red Gum. Specimens of Small-flowered Grevillea and Pine Donkey Orchid were recorded within the existing easement and are not required to be impacted by the proposal. However, these flora species may potentially be inadvertently impacted during the construction of the subtransmission line. Considerations under Section 7.3 of the BC Act (2016) found the REZ is unlikely to significantly affect the life cycle of any of these flora species or place any viable local populations of these species at risk of extinction.

### Threatened fauna species listed under the state BC Act 2016

Three threatened fauna species were recorded during fieldwork. These species were:

- *Pomatostomus temporalis* subsp. *temporalis* (Grey-crowned Babbler);
- *Climacteris picumnus victoriae* (Brown Treecreeper);
- *Haliaeetus leucogaster* (White-bellied Sea-Eagle).

The proposal will impact suitable habitat for the Grey-crowned Babbler and Brown Treecreeper. A small number of Grey-crowned Babbler nests/roosts may also be impacted. No suitable habitat is likely to be impacted for *Haliaeetus leucogaster* (White-bellied Sea-Eagle). *Delma vescolineata* (Hunter Valley Delma) has been recorded in 2024 within the neighbouring property to the west of the Muswellbrook proposed STS within similar habitat (Wildthing Environmental Consultants, 2024). According to the BioNet Atlas records (DCCEEW, 2026a), Additional records of *D. vescolineata* have also been found within close proximity to the Muswellbrook STS. Taking this into consideration it is highly likely that *D. vescolineata* would utilise habitat within the impact area of the Muswellbrook STS. Considerations under Section 7.3 of the BC Act (2016) found the REZ is unlikely to significantly affect the life cycle of any of the addressed threatened fauna species or place any viable local populations of these species at risk of extinction.

### Endangered Populations listed under the state BC Act 2016

Specimens of *Acacia pendula* (Weeping Myall) listed as an endangered population in the Hunter Catchment Weeping Myall were recorded long the Golden Highway, Warkworth and Liddell STS Hub at Antiene. The specimens along the Golden Highway, Warkworth were found in one location and are likely naturally occurring specimens. A fire had relatively recently burnt through all the specimens of *A. pendula* at Warkworth (first half of 2026) leaving burnt stems. As the non-pendulous form of *A. pendula* readily suckers from the roots (Umwelt 2006), these plants are likely to reshoot. It is thought that any suckering specimens of *A. pendula* along within the new easement at Warkworth could be protected in-situ and managed to not impact the overhead wires. Specimens of Weeping Myall at Liddell STS had been planted within a treed belt running east-west along Hebdon Road on the northern boundary of the Lake Liddell Recreation Area. The two specimens of *A. pendula* are required to be removed for access to the Liddell STS works. As these specimens have been planted as part of a treed belt the removal of these specimens would not likely be considered significant.

### Commonwealth Environment Protection and Biodiversity Conservation Act 1999 & Matters of National Environmental Significance

Four Nationally Listed Threatened Ecological Communities were identified within the REZ:

- Kurri sand swamp woodland of the Sydney Basin bioregion;
- Central Hunter Valley eucalypt forest and woodland;
- Swamp Oak Floodplain Forest;
- White Box – Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland

It was determined the REZ will have an incremental impact on these TEC's particularly that of Central Hunter Valley eucalypt forest and woodland within new alternate easements along the Mitchell Line of road, however it is unlikely to have a significant impact on any of these nationally listed threatened communities. A referral is unlikely to be required.

Three Nationally threatened flora species were recorded during fieldwork:

- *Eucalyptus glaucina* (Slaty Red Gum)
- *Eucalyptus parramattensis* subsp. *decadens* (Drooping Red Gum)
- *Grevillea parviflora* subsp. *parviflora* (Small-flowered Grevillea)

One Nationally threatened fauna species *Climacteris picumnus victoriae* (Brown Treecreeper) was recorded within the REZ land. *Delma vescolineata* (Hunter Valley Delma) was also considered likely to occur within parts of the REZ particularly in the far west.

It was determined the REZ is unlikely to have a significant impact on any of these nationally listed threatened species.



## Significant/Hollow-bearing Trees

A total of 56 significant (larger trees and/or those containing hollows) trees were recorded along the proposed REZ. Of the 56, 28 trees are likely to require removal. Seven trees will also require branch trimming. A summary of the significant tree impacts is shown in Table 3.

**Table 3.: Summary of the significant tree impacts**

| TREE                    | REQUIRING REMOVAL | REQUIRING BRANCH TRIMMING |
|-------------------------|-------------------|---------------------------|
| Habitat Tree            | 16                | 7                         |
| Non-habitat Tree        | 12                | 0                         |
| Total of impacted trees | 28                | 7                         |

A number of mitigation measures have been specified to minimise the impact of the proposal and to protect biodiversity values.

In conclusion, the proposal will result in an incremental impact to native vegetation and habitat for a number of threatened communities and species. Given the mitigation measures the proposal is unlikely to disrupt the life cycle of any addressed threatened species, endangered population or endangered ecological community such that local extinction would occur.

**CONTENTS**

|            |                                                                            |           |
|------------|----------------------------------------------------------------------------|-----------|
| <b>1.0</b> | <b>INTRODUCTION</b>                                                        | <b>1</b>  |
| 1.1        | THE PROPOSAL                                                               | 1         |
| 1.2        | PROPOSED MODIFICATIONS                                                     | 2         |
| <b>2.0</b> | <b>SUBJECT LAND CONTEXT</b>                                                | <b>3</b>  |
| 2.1        | HYDROGEOGRAPHY                                                             | 3         |
| <b>3.0</b> | <b>LEGISLATIVE CONTEXT</b>                                                 | <b>5</b>  |
| 3.1        | NSW ENVIRONMENTAL PLANNING AND ASSESSMENT AMENDMENT ACT 2017               | 5         |
| 3.2        | NSW BIODIVERSITY CONSERVATION ACT 2016                                     | 5         |
| 3.3        | STATE ENVIRONMENTAL PLANNING POLICY (BIODIVERSITY AND CONSERVATION) 2021   | 7         |
| 3.4        | BIOSECURITY ACT 2015                                                       | 7         |
| 3.5        | COMMONWEALTH ENVIRONMENT PROTECTION AND BIODIVERSITY CONSERVATION ACT 1999 | 7         |
| 3.6        | LICENSING                                                                  | 8         |
| <b>4.0</b> | <b>METHODOLOGY</b>                                                         | <b>9</b>  |
| 4.1        | DESKTOP ASSESSMENT                                                         | 9         |
| 4.2        | FIELD ASSESSMENT                                                           | 10        |
| 4.2.1      | VEGETATION ASSESSMENT                                                      | 10        |
| 4.2.2      | DIURNAL FAUNA SURVEY                                                       | 10        |
| 4.2.3      | GENERAL HABITAT FOR NATIVE SPECIES                                         | 10        |
| 4.2.4      | HABITAT FOR SIGNIFICANT SPECIES                                            | 10        |
| 4.2.5      | TREE SURVEY                                                                | 12        |
| 4.3        | SIGNIFICANT SPECIES                                                        | 12        |
| <b>5.0</b> | <b>RESULTS</b>                                                             | <b>17</b> |
| 5.1        | FLORA ASSEMBLAGES                                                          | 17        |
| 5.1.1      | THREATENED ECOLOGICAL COMMUNITIES                                          | 40        |
| 5.1.2      | THREATENED AND RARE FLORA SPECIES                                          | 41        |
| 5.1.3      | ENDANGERED POPULATIONS                                                     | 44        |
| 5.1.4      | PRIORITY WEEDS AND WEEDS OF STATE AND NATIONAL SIGNIFICANCE                | 46        |
| 5.2        | HABITAT APPRASIAL                                                          | 48        |
| 5.2.1      | HABITAT DESCRIPTION AND DISTRIBUTION IN THE VICINITY                       | 48        |
| 5.2.2      | HABITAT CORRIDORS                                                          | 48        |
| 5.2.3      | TREE SURVEY                                                                | 48        |
| 5.3        | HABITAT FOR SIGNIFICANT SPECIES                                            | 50        |
| 5.4        | FAUNA APPRASIAL RESULTS                                                    | 69        |



|             |                                                                                                                   |            |
|-------------|-------------------------------------------------------------------------------------------------------------------|------------|
| 5.4.1       | DIURNAL SURVEYS                                                                                                   | 69         |
| 5.5         | SURVEY LIMITATIONS                                                                                                | 70         |
| <b>6.0</b>  | <b>IMPACT ASSESSMENT</b>                                                                                          | <b>71</b>  |
| 6.1         | AVOIDANCE AND MINIMISATION OF IMPACTS                                                                             | 71         |
| 6.2         | DIRECT IMPACT                                                                                                     | 71         |
| 6.3         | INDIRECT IMPACTS                                                                                                  | 72         |
| 6.4         | MITIGATION MEASURES                                                                                               | 75         |
| <b>7.0</b>  | <b>CONSIDERATIONS UNDER SECTION 7.3 OF THE BC ACT 2016</b>                                                        | <b>77</b>  |
| <b>8.0</b>  | <b>ASSESSMENT OF SERIOUS AND IRREVERSIBLE IMPACTS</b>                                                             | <b>82</b>  |
| 8.1         | POTENTIAL SAI ENTITIES                                                                                            | 82         |
| <b>9.0</b>  | <b>CONSIDERATIONS UNDER THE COMMONWEALTH ENVIRONMENT PROTECTION AND BIODIVERSITY CONSERVATION ACT 1999</b>        | <b>84</b>  |
| 9.1         | CENTRAL HUNTER VALLEY EUCALYPT FOREST AND WOODLAND                                                                | 87         |
| 9.2         | WHITE BOX -YELLOW BOX- BLAKELYS RED GUM GRASSY WOODLAND AND DERIVED NATIVE GRASSLAND                              | 90         |
| 9.3         | KURRI SAND SWAMP WOODLAND OF THE SYDNEY BASIN BIOREGION                                                           | 93         |
| 9.4         | COASTAL SWAMP OAK (CASUARINA GLAUCA) FOREST OF THE NEW SOUTH WALES AND SOUTH-EAST QUEENSLAND ECOLOGICAL COMMUNITY | 95         |
| 9.5         | BROWN TREECREEPER                                                                                                 | 97         |
| <b>10.0</b> | <b>CONCLUSION</b>                                                                                                 | <b>99</b>  |
| <b>11.0</b> | <b>BIBLIOGRAPHY</b>                                                                                               | <b>100</b> |

## LIST OF APPENDICES

APPENDIX A – SURVEY POINTS ALONG REZ ROUTE

APPENDIX B – DETAILED MAPS OF REZ

APPENDIX C – SECTION 7.3 OF THE BC ACT 2016 TEST OF SIGNIFICANCE

APPENDIX D – FLORA SPECIES LIST

APPENDIX E – SURVEYED TREE DATA

APPENDIX F – SIGNIFICANT FLORA SPECIES INFORMATION

APPENDIX G – FAUNA SPECIES LIST

**Acronyms and Abbreviations used in this report**

|          |                                                                 |
|----------|-----------------------------------------------------------------|
| AOBV     | Area of outstanding Biodiversity Value                          |
| BAAS     | Biodiversity Assessors Accreditation System                     |
| BAM      | Biodiversity Assessment Method                                  |
| BAMC     | Biodiversity Assessment Method Calculator                       |
| BAR      | Biodiversity Assessment Report                                  |
| BC Act   | Biodiversity Conservation Act 2016                              |
| BDAR     | Biodiversity Development Assessment Report                      |
| BOAMS    | Biodiversity Offsets and Agreement Management System            |
| BOS      | Biodiversity Offset Scheme                                      |
| BOSET    | Biodiversity Offsets Scheme Entry Tool                          |
| DCCEEW   | Department of Climate Change, Energy, the Environment and Water |
| DPE      | Department of Planning and Environment (NSW)                    |
| EEC      | Endangered Ecological Community                                 |
| EPBC Act | Environmental Protection & Biodiversity Conservation Act 1999   |
| EP&A Act | Environmental Planning & Assessment Act 1979                    |
| IBRA     | Interim Biogeographic Regionalisation for Australia             |
| LGA      | Local Government Area                                           |
| LLS Act  | Local Land Services Act 2013                                    |
| NES      | Matters of National Significance under the EPBC Act             |
| NPW Act  | National Parks & Wildlife Act 1974                              |
| OEH      | Office of Environment & Heritage (now DPE)                      |
| PCT      | Plant Community Type                                            |
| PMST     | Protected Matters Search Tool                                   |
| SAII     | Serious and Irreversible Impacts                                |
| SEPP     | State Environmental Planning Policy                             |
| REZ      | Renewable Energy Zone                                           |
| SP       | Single Pole                                                     |
| TEC      | Threatened Ecological Community                                 |

## 1.0 INTRODUCTION

Flora, fauna and habitat studies have been undertaken to update the ecological assessment to reflect the proposed eleven (11) modifications to the original network upgrade of the existing electricity network between Kurri Kurri and Muswellbrook as part of Ausgrids Hunter-Central Coast Renewable Energy Project (REZ). The investigations were in accordance with the requirements of the *Environmental Planning and Assessment Amendment Act 2017* (EP&A Act 2017), the *Biodiversity Conservation Act 2016* (BC Act 2016) and the *Commonwealth Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act 1999). The results are presented here in the form of an Ecological Assessment.

### 1.1 THE PROPOSAL

The NSW energy system is undergoing a major transformation to transition its reliance on coal-fired power to renewable energy. By 2040, the remaining four of NSW's coal-fired generators will have been retired, while increasingly higher demand for electricity will be required.

The construction of new and upgraded network infrastructure is required to ensure energy can be transferred from Renewable Energy Zones (REZs) to major load centres in NSW. The Hunter and Central Coast REZ (HCC REZ) was formally declared on 9 December 2022 and comprises a specified geographical area which lies within the Ausgrid distribution network. EnergyCo as the infrastructure planner has nominated that planned, new and existing network infrastructure with the geographical area is specified as REZ infrastructure.

The HCC REZ is expected to accommodate renewable energy generation and storage projects, and become a hub for low-emission industrial development in the region. EnergyCo conducted an initial engagement process with industry regarding the potential investment in renewable generation and storage projects in the HCC REZ. From this, an intended network capacity of 1GW has been identified for HCC REZ.

#### **The REZ proposal will be completed in Stages:**

**Stage 1:** A proposed dual circuit high-capacity 132kV subtransmission lines ~60km between the existing Singleton and Kurri Sub-Transmission Substations (STS's). This will include pole replacements, access track works and vegetation clearing. This stage also includes upgrades at Singleton and Kurri STSs. The 132kV subtransmission line will be almost entirely located within Ausgrids existing maintained easement. Depending on the width of the required easement (either 20 or 45m) some areas of native vegetation along the edge of the easement will be impacted. In most cases only minor branch trimming will be required. Some sections of new easement are also proposed which will require the removal of native vegetation in some areas.



## Stage 2:

A proposed dual circuit high-capacity 132kV subtransmission line (~25km) between the proposed Eastern Hub 132kV Switching Station located on the northern side of Lake Liddell to the existing Singleton STS.

Rearrangement of Ausgrid's existing Upper Hunter subtransmission network to provide connection between an existing subtransmission line known as 95U, which ultimately will connect to Eastern Hub SS and Ausgrid's existing Muswellbrook STS. The subtransmission line (~5km) will be almost entirely located within Ausgrid's existing maintained easement. Some branch trimming and the removal of a small number of trees will likely be required.

The location of the Hunter REZ Project is shown in Figure 1.1.

## 1.2 PROPOSED MODIFICATIONS

Eleven modifications have been proposed for the REZ project. These modifications are described below in Table 2.1 and shown in maps in Appendix B. With the exception of modification 3 Pothana Lane Belford, Ausgrid proposes to remove the existing poles and wires along the original route at these locations.

**Table 2.1. Proposed modifications to the proposal**

| No. | Modification Name                                             | Description/ Justification                                                                                                                                                                                                                                                                                                                      | Figure No.      |
|-----|---------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------|
| 1   | Hunter Expressway crossing – Branxton                         | The line deviates further to the south to cross the Hunter Expressway before turning back to meet the original alignment (~500m). This modification addresses some significant constructability issues with the original route.                                                                                                                 | B10             |
| 2   | ARTC crossing off Standen Drive, Belford                      | There is an existing building within the easement of the original route. The proposed route modification relocates the ARTC crossing approximately 200m west of the existing crossing thereby avoiding this interaction. In addition, the revised crossing is also more perpendicular to the railway line which is a preferred outcome for ARTC | B10             |
| 3   | Pothana Lane, Belford                                         | The proposed modification diverts north through the TSR for approximately 500m to the New England Highway and then heads west for approximately 1.9km before rejoining the original route.                                                                                                                                                      | B11, B11a, B11b |
| 4   | Golden Highway, Whittingham                                   | The line diverts to the south of the original route for approximately 1km before rejoining in the original route due to detailed design.                                                                                                                                                                                                        | B13             |
| 5   | Mitchell Line of Road - Singleton Military Area 1 (Dochra)    | Relocation of existing 66kV line underground and revision of the 132KV OH to achieve height compliance with Defence requirements re Dochra Air Field.                                                                                                                                                                                           | B13             |
| 6   | Mitchell Line of Road - Singleton Military Area 2             | The proposed modification of approximately 1.3km diverts immediately to the south due to detailed design.                                                                                                                                                                                                                                       | B14             |
| 7   | Mitchell Line of Road - Singleton Military Area 3             | Approximately 600m diverts to the south to stay on the southern side of Mitchell line of road and avoid two road crossings.                                                                                                                                                                                                                     | B15             |
| 8   | Mitchell Line of Road - Singleton Military Area 4 (Redgroves) | The line diverts to the north along Mitchell Line of Road for approximately 2km to move out of the existing easement in Singleton Military Area.                                                                                                                                                                                                | B15, B16        |
| 9   | Jerry's Plains Road - Mt Thorley                              | Relocation from existing Ausgrid easement to west to the Jerry's Plains Road for approximately 3.5km before rejoining the original route where it crosses the highway at Gouldsville.                                                                                                                                                           | B16, B17, B18   |

| No. | Modification Name          | Description/ Justification                                                                                                                                                      | Figure No. |
|-----|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|
|     |                            | This option removes two railway line crossings and reduces impact on Yancoal buffer lands. Also included is a minor realignment on Putty Road.                                  |            |
| 10  | Glendell Mine, Ravensworth | Minor realignment to immediately to east of existing easement for approximately 1 km due to inaccessibility of existing easement for construction and future maintenance works. | B30        |
| 11  | Antiene STSS – Hebden      | Minor realignment of southern connection to Antiene STSS due to detailed design and constructability issues. Addition of new northern connection approximately 1km.             | B31, B32   |


## 2.0 SUBJECT LAND CONTEXT

The REZ project is located entirely within the Sydney Basin Bioregion and Hunter Sub-bioregion (regions gazetted by the Minister, or an Interim Biogeographical Regionalisation of Australia (IBRA Bioregion). Mitchell Landscapes located within the REZ project are Newcastle Coastal Ramp, Central Hunter Foothills, Central Hunter Alluvial Plains. The REZ project is located within the Cessnock, Singleton and Muswellbrook Local Government Areas (LGA's).

### 2.1 HYDROGEOGRAPHY

The entire REZ project is located entirely within the greater Hunter River Catchment. The easement crosses a number of minor and major streams including the Hunter River, Glennies and Bowmans Creeks.

Figure 1.1 Location Map of REZ Project



|                                                                                                                                                                                                                                                                                                                                                                                                                                         |           |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------|
| Job No.                                                                                                                                                                                                                                                                                                                                                                                                                                 | 12875     |
| A4 Scale                                                                                                                                                                                                                                                                                                                                                                                                                                | 1:300,000 |
| Disclaimer: While all reasonable care has been taken to ensure the information contained on this map is up to date and accurate, no guarantee is given that the information portrayed is free from error or omission. In addition the spatial accuracy of the map is wholly dependent on source data. Please verify the accuracy of all information prior to use. Development footprint areas should be used for indicative areas only. |           |




Figure 1.1

Location of proposed  
Transmission Lines, Eastern  
Hub and existing STSs

8 November 2024

**WILDTHING**  
Environmental Consultants  
(a Division of Tattersall Lander Pty Ltd)  
ABN 41 003 509 215



### 3.0 LEGISLATIVE CONTEXT

The following sections detail the legislative frameworks relevant to this report.

#### 3.1 NSW ENVIRONMENTAL PLANNING AND ASSESSMENT AMENDMENT ACT 2017

The assessment of development applications in NSW is regulated under Part 4 or Part 5 of the EP&A Act. Under Section 110B(1)(a) of the EP&A Act, Ausgrid is both the proponent and determining authority and, as such, acts under Part 5 of the EP&A Act. Section 1.7 of the EP&A Act links proponents to Part 7 of the BC Act for the operation of the EP&A Act in connection with the terrestrial environment. The EP&A Act is also supported by other statutory environmental planning instruments, including State Environmental Planning Policies (SEPPs). The following SEPPs and acts are relevant to this report:

- State Environmental Planning Policy (Biodiversity and Conservation) 2021 (Biodiversity and Conservation SEPP)
- State Environmental Planning Policy (Resilience and Hazards)

#### 3.2 NSW BIODIVERSITY CONSERVATION ACT 2016

The Biodiversity Offsets Scheme (BOS) Entry Test (BOSET) does not apply to Part 5 developments as it does for Part 4. The BOS only applies to Part 5 in the circumstances outlined below:

1. Is the proposed activity to be carried out in an Area of Outstanding Biodiversity Value (AOBV)?
2. Is the proposed activity likely to significantly affect threatened species?

The Review of Environmental Factors (REF), is prescribed by Part 5 of the Environmental Planning and Assessment Act 1979 and Part 8 of the Environmental Planning and Assessment Regulation 2021. As part of the REF, a Test of Significance is undertaken to determine if the activity is likely to harm a threatened species. If the answer is yes to either there are two options as follows:

- Option 1 - Prepare a Species Impact Statement (SIS) in consultation with Department of Climate Change, Energy, the Environment and Water (NSW DCCEEW);  
or
- Option 2 - Opt in to the Biodiversity Offset Scheme (BOS)

The BOS criteria as it applies to Part 5 for the proposal is shown in Table 3.1. A Biodiversity Values Map and Threshold Tool (BMAT) map for the site is shown in Figure 3.1.

##### 3.2.1 SERIOUS AND IRREVERSIBLE IMPACTS

The BC Act imposes various obligations on determining authorities in relation to impacts on biodiversity values that are at risk of a serious and irreversible impact. Under the BC Act, a determination of whether an impact is serious and irreversible must be made in accordance with the principles prescribed in section 6.7 of the BC Regulation 2017. Species and ecological communities (SAIL entities) that are likely to be the subject of serious and irreversible impacts are listed on the NSW DCCEEW website (NSW DCCEEW 2025f). No candidate SAIL entities were found to be present within the site.

**Table 3.1: Criteria for entry into the Biodiversity Offsets Scheme in relation to the proposed development.**

| CRITERIA FOR ENTRY INTO THE BIODIVERSITY OFFSETS SCHEME (BOS)                                                                                                                                                                                                                                          | SECTION CRITERIA ADDRESSED | ASSESSMENT OF CRITERIA                                                                                                                                    |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------|
| Development activities that have the potential to impact Areas of Outstanding Biodiversity Value (AOBV) as listed under Part 3 of the BC Act.                                                                                                                                                          | Section 7.0                | No declared areas of outstanding biodiversity value were located within or in proximity to the site.                                                      |
| Development activities that have the potential to cause a significant impact on a threatened species, population or ecological community, listed under Schedules 1 and 2 of the BC Act, as determined by application of a five-part-test of significance in accordance with Section 7.3 of the BC Act; | Section 7.0                | The five-part test found no significant impact on threatened species, populations or ecological communities listed under Schedules 1 and 2 of the BC Act. |

### 3.3 STATE ENVIRONMENTAL PLANNING POLICY (BIODIVERSITY AND CONSERVATION) 2021

The State Environmental Planning Policy (Biodiversity and Conservation) 2021 (Biodiversity and Conservation SEPP) consolidates, transfers and repeals provisions of the following 11 SEPPs (or deemed SEPPs):

1. SEPP (Vegetation in Non-Rural Areas) 2017 (Vegetation SEPP)
2. SEPP (Koala Habitat Protection) 2020 (Koala SEPP 2020)
3. SEPP (Koala Habitat Protection) 2021 (Koala SEPP 2021)
4. Murray Regional Environmental Plan No 2—Riverine Land (Murray REP)
5. SEPP No 19—Bushland in Urban Areas (SEPP 19)
6. SEPP No 50—Canal Estate Development (SEPP 50)
7. SEPP (Sydney Drinking Water Catchment) 2011 (Sydney Drinking Water SEPP)
8. Sydney Regional Environmental Plan No 20 – Hawkesbury – Nepean River (No 2 – 1997) (Hawkesbury–Nepean River SREP)
9. Sydney Regional Environmental Plan (Sydney Harbour Catchment) 2005 (Sydney Harbour Catchment SREP)
10. Greater Metropolitan Regional Environmental Plan No 2 – Georges River Catchment (Georges River REP)
11. Willandra Lakes Regional Environmental Plan No 1 – World Heritage Property (Willandra Lakes REP).

Each consolidated SEPP now makes up a chapter in the SEPP (Biodiversity and Conservation) 2021. The Koala Habitat Protection SEPPs only apply to 'development' under Part 4 of the NSW Environmental Planning and Assessment Act 1979 (EP&A Act), specifically excluding Part 5 'activities'. Additionally, no other listed SEPPs apply to the proposal.

### 3.4 BIOSECURITY ACT 2015

The NSW Biosecurity Act 2015 provides regulatory controls and powers to manage priority weeds in NSW. For weed management this Act divides NSW into regions based on combined LGAs and priority weeds for a region are listed. Some weeds are managed at a state level as they form part of a broader containment strategy. The legislation compliments listed Weeds of National Significance (WoNS).

### 3.5 COMMONWEALTH ENVIRONMENT PROTECTION AND BIODIVERSITY CONSERVATION ACT 1999

The purpose of the EPBC Act is to ensure that actions likely to cause a significant impact on Matters of National Environmental Significance (MNES) undergo a process of assessment. Under the EPBC Act, an action includes a project, undertaking, development or activity that may impact MNES. An action that 'has, will have or is likely to have a significant impact on a MNES' is deemed to be a 'controlled action' and may not be undertaken without prior approval from the commonwealth minister for the Department of Climate Change, Energy, the Environment and Water (DCCEEW).

MNES categories listed under the EPBC Act are:

- World heritage properties;



- National heritage places;
- Wetlands of international importance (Ramsar wetlands);
- Threatened species and ecological communities (Section 18 and 18A);
- Migratory species;
- Commonwealth marine areas;
- Nuclear actions (including uranium mining); and
- A water resource, in relation to coal seam gas development and large coal mining development.

Initially MNES protected under the EPBC Act are assessed in accordance with the Significant Impact Guidelines 1.1 - Matters of National Environmental Significance (DoE 2013). This is performed to determine if there is likelihood for an action to have a significant impact on MNES. An action will require referral to, and may require the approval of, the commonwealth minister for the Environment (in addition to any local or state government consent or approval) if that action will have, or is likely to have, a significant impact on the environment or on a MNES.

### **3.6 LICENSING**

Fieldwork undertaken by Wildthing Environmental Consultants was carried out under NPWS Scientific Investigation Licence SL100345 and under Animal Care and Ethics Approval: Animal Research Authority Issue by the Department of Primary Industries (Trim File No. 13/251) for Fauna Survey for Biodiversity and Impact Assessment.

## 4.0 METHODOLOGY

### 4.1 DESKTOP ASSESSMENT

A site-specific literature and database review was undertaken prior to conducting the field survey and the preparation of this report. A list of the resources reviewed, the date they were accessed and the spatial extent of the search conducted, where relevant, is provided in Table 4.1.

**Table 4.1: Desktop Resources**

| RESOURCE                                                                                                                                                 | LAST ACCESS DATE           | SPATIAL EXTENT                 |
|----------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------|--------------------------------|
| <b>Biodiversity Values and Landscape Maps</b>                                                                                                            |                            |                                |
| BioNet Atlas of NSW Wildlife (BioNet) (NSW DCCEE, 2026a)                                                                                                 | 28 May 2024<br>6 June 2026 | 10x10km radius of subject land |
| Commonwealth Protected Matters Search Tool (PMST) (DCCEE 2024a)                                                                                          | 5 June 2024<br>June 2026   | 10x10km radius of subject land |
| NSW Biodiversity Values Map (DPE 2024b)                                                                                                                  | 28 May 2024                | Entire subject land            |
| SIX Maps<br>-Base Map - LPI 1:25,000 digital topographic databases (DTDB) (LPI 2024)<br>-Cadastral data LPI digital cadastral database (DCDB) (LPI 2024) | Various dates              | Entire subject land            |
| NSW Government SEED Mapping (NSW Government 2024)                                                                                                        | Various dates              | Entire subject land            |
| BioNet NSW (Mitchell) Landscapes – Version 3.1 (OE 2016a)                                                                                                | 5 November 2024            |                                |
| NSW Interim Biogeographic Regions of Australia (IBRA region and sub-regions) – Version 7 (DAWE 2016).                                                    | 5 November 2024            | Entire subject land            |
| <b>Threatened Species and Vegetation Databases</b>                                                                                                       |                            |                                |
| Commonwealth species profiles and threats database (SPRAT) (DCCEE 2024a)                                                                                 | Various dates              | -                              |
| DPE Profiles of threatened species, population, and ecological communities (DPE 2024d)                                                                   | Various dates              | -                              |
| DPE BioNet vegetation classification database (DPE 2024c)                                                                                                | Various dates              |                                |

## **4.2 FIELD ASSESSMENT**

A summary of the time spent on site during fieldwork and the prevailing weather conditions at the time is contained in Table 4.2.

### **4.2.1 VEGETATION ASSESSMENT**

The initial determination of the basic vegetation community boundaries was undertaken through the review of an orthophoto covering the site. Following this, a detailed ground survey was conducted in accordance with the Department of Environment and Conservation's (NSW) Threatened Biodiversity Survey and Assessment Guidelines – Working Draft (Department of Environment and Conservation, 2004). Flora searches were undertaken in the manner described by Cropper (1993) as the 'Random Meander Technique'. This involved walking in a random manner throughout the easement route, particularly within areas of impacted native vegetation.

A list of all flora species identified on site has been provided in Appendix A.

### **4.2.2 DIURNAL FAUNA SURVEY**

Opportunistic sightings of species and secondary indications (scats, scratches, diggings, tracks etc.) of resident fauna were noted and included:

- Surveys for avifauna;
- Surveys for herpetofauna;
- checks for obvious nests of raptors;
- checking trees (particularly smooth-barked species) for scratches consistent with arboreal mammals.

### **4.2.3 GENERAL HABITAT FOR NATIVE SPECIES**

From the vegetation appraisal, diurnal fauna survey and a general inspection of the site and surrounding areas, a subjective assessment of the general habitat value of this site was made. Considered in this assessment were:

- occurrence of that habitat type in the general vicinity;
- degree of disturbance and degradation;
- area occupied by that habitat on site;
- continuity with similar habitat adjacent to the site, or connection with similar habitat off site by way of corridors; and
- structural and floral diversity.

### **4.2.4 HABITAT FOR SIGNIFICANT SPECIES**

The subject land area was evaluated as potential habitat for each of the threatened species reported on the BioNet (NSW DCCEEW, 2026a) and PMST (DCCEEW, 2026a) databases from within 10km of the site. This evaluation was based on home range, feeding, roosting, breeding, movement patterns and corridor requirements for fauna and hydrology, soil types, aspect and structural formation for flora species. The list of threatened species recorded within these databases is provided within Table 4.3 and an assessment of the likelihood of occurrence of these threatened species within the subject land is provided in Table 5.4.



**Table 4.2: Survey Dates, Times and Weather Conditions**

| DATE                          | TIME        | LOCATIONS                                                                                                                       | WEATHER                                                    |
|-------------------------------|-------------|---------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------|
| Wednesday<br>29 May 2024      | 0800 - 1530 | Kurri Kurri STS, Scales Road, Hart Road, Bishops Bridge, Majors Lane                                                            | 0/8 Cloud, 6.7°C, 700% relative humidity, Wind 6km/hr WNW  |
| Thursday<br>30 May 2024       | 8:00 - 1500 | Lovedale - Greta,                                                                                                               | 0/8 Cloud, 9.7°C, 100% relative humidity, Wind 2km/hr ESE  |
| Monday<br>3 June 2024         | 0830 - 1630 | Greta - Branxton                                                                                                                | 0/8 Cloud, 10.6°C, 82% relative humidity, Wind 19km/hr WNW |
| Tuesday<br>4 June 2024        | 0830 - 1630 | Lower Belford                                                                                                                   | 7/8 Cloud, 9.7°C, 71% relative humidity, Wind 17km/hr NW   |
| Thursday<br>6 June 2024       | 0800 - 1630 | From intersection of Golden Highway & New England Highway. Mining buffer land, Long Point                                       | 1/8 Cloud, 10.5°C, 93% relative humidity, Wind 4km/hr WSW  |
| Tuesday<br>11 June 2024       | 0830 - 1400 | Singleton STS, Maison Dieu, Glennies Creek                                                                                      | 2/8 Cloud, 6.7°C, 100% relative humidity, Wind 7km/hr W    |
| Thursday<br>13 June 2024      | 0930 - 1500 | Singleton Military Base, Bulga Coal Offset                                                                                      | 0/8 Cloud, 8.8°C, 60% relative humidity, Wind 19km/hr SSW  |
| Monday<br>1 July 2024         | 0830 - 1500 | Camberwell, Glennies Creek, Eastern Hub (Lake Liddell, Muswellbrook)                                                            | 3/8 Cloud, 10.6°C, 63% relative humidity, Wind 11km/hr SSW |
| Thursday<br>22 August 2024    | 1030 - 1200 | Kurri Kurri STS, Hart Road<br>Mapping new easement alignment, Bushfire Assessor also on site.                                   | 6/8 Cloud, 17.6°C, 75% relative humidity, Wind 13km/hr NW  |
| Thursday<br>29 August 2024    | 0900 - 1430 | Muswellbrook STS area & existing easement to the NE.                                                                            | 8/8 Cloud, 15°C, 80% relative humidity, Wind 13km/hr SW    |
| Friday<br>4 April 2025        | 1000 - 1130 | Western side of Hunter Expressway over from Kurri STS.                                                                          | 0/8 Cloud, 20.5°C, 69% relative humidity, Wind 15km/hr WNW |
| Wednesday<br>12 November 2025 | 0800 - 1200 | Assessment of new route along Mitchell Line of Road bordering Singleton Army Base, Putty Road and Golden Highway Mount Thorley. | 0/8 Cloud, 14.4°C, 39% relative humidity, Wind 4km/hr WSW  |
| Wednesday<br>20 November 2025 | 1130 - 1230 | Inspection of alternate option on Golden Highway near Muddies Creek                                                             | 3/8 Cloud, 32.5°C, 25% relative humidity, Wind 15km/hr NW  |
| Thursday<br>22 January 2026   | 1045 - 1200 | Inspecting revised impact areas NW of Muswellbrook STS                                                                          | 0/8 Cloud, 32.5°C, 44% relative humidity, Wind 15km/hr WNW |
| Thursday<br>22 January 2026   | 1300- 1400  | Inspecting revised impact areas Eastern Hub (Lake Liddell, Muswellbrook).                                                       | 0/8 Cloud, 32°C, 41% relative humidity, Wind 15km/hr WNW   |
| Wednesday<br>20 May 2026      | 1100 - 1430 | Assessment of new route along New England Highway at Lower Belford.                                                             | 3/8 Cloud, 18°C, 47% relative humidity, Wind 11km/hr SW    |

A detailed methodology for the surveys listed within Table 4.2 above have been described in the following Sections 4.2.1 – 4.2.5:

---

## 4.2.5 TREE SURVEY

During the fieldwork, a survey was undertaken to identify significant trees (larger trees and/or those containing hollows or other habitat attributes) which may be impacted by the proposal. Hollow-bearing trees are a habitat resource utilised by a variety of native avifaunal and mammalian species. This resource is usually a limiting factor in the occurrence of hollow-dependent species on a site, due to the time taken for hollows to form in trees. It must be noted that observations made from ground level may fail to record a small number of hollows that are obscured. Some entrances may also not lead to a cavity. The internal dimensions of the hollows are also impossible in many cases to determine from the ground.

## 4.3 SIGNIFICANT SPECIES

The following threatened species listed in Table 4.3 have been recorded on the BioNet (DPE, 2026a) and PMST (DCCEEW, 2026a) Databases as occurring within 10km of the subject land. Species marked with an asterisk (\*) are listed on the DCCEEW Database as having habitat likely to occur within 10km of the subject land.

**Table 4.3: Threatened species, endangered populations and ecological communities considered**

| Scientific Name                                          | Common Name                | BC Act 2016 | EPBC Act 1999 |
|----------------------------------------------------------|----------------------------|-------------|---------------|
| <b>Flora Species</b>                                     |                            |             |               |
| <i>*Wollemia nobilis</i>                                 | Wollemi Pine               | E4A         | CE            |
| <i>*Caladenia tessellata</i>                             | Thick-lipped Spider-orchid | V           | V             |
| <i>Corybas dowlingii</i>                                 | Red Helmet Orchid          | E1          |               |
| <i>*Cryptostylis hunteriana</i>                          | Leafless Tongue Orchid     | V           | V             |
| <i>Diuris pedunculata</i>                                | Small Snake Orchid         | E1          | E             |
| <i>Diuris tricolor</i>                                   | Pine Donkey Orchid         | V           |               |
| <i>Prasophyllum petilum</i>                              | Tarengo Leek Orchid        | E1          | E             |
| <i>*Prasophyllum</i> sp. <i>Wybong</i>                   | A Leek Orchid              |             | CE            |
| <i>Pterostylis chaetophora</i>                           | Tall Rustyhood             | V           |               |
| <i>*Pterostylis gibbosa</i>                              | Illawarra Greenhood        | E1          | E             |
| <i>*Rhizanthella slateri</i>                             | Eastern Underground Orchid | V           | E             |
| <i>*Thelymitra adorata</i>                               | Wyong Sun-orchid           | E4A         | CE            |
| <i>*Dichanthium setosum</i>                              | Bluegrass                  | V           | V             |
| <i>*Cynanchum elegans</i>                                | White-flowered Wax Plant   | E1          | E             |
| <i>*Brachyscome brownii</i>                              | Brachyscome brownii        |             | CE            |
| <i>*Ozothamnus tessellatus</i>                           | Ozothamnus tessellatus     | V           | V             |
| <i>*Picris evae</i>                                      | Hawkweed                   | V           | V             |
| <i>*Rutidosia heterogama</i>                             | Heath Wrinklewort          | V           | V             |
| <i>*Lepidium aschersonii</i>                             | Spiny Peppergrass          | V           | V             |
| <i>*Tetratheca juncea</i>                                | Black-eyed Susan           | V           | V             |
| <i>*Tylophora linearis (Vincetoxicum forsteri)</i>       |                            | V           | E             |
| <i>*Acacia bynoeana</i>                                  | Bynoe's Wattle             | E1          | V             |
| <i>Dillwynia tenuifolia</i>                              |                            | V           |               |
| <i>*Swainsona murrayana</i>                              | Slender Darling-pea        | V           | V             |
| <i>*Prostanthera cineolifera</i>                         | Singleton Mint Bush        | V           | V             |
| <i>*Angophora inopina</i>                                | Charmhaven Apple           | V           | V             |
| <i>Callistemon linearifolius</i>                         | Netted Bottle Brush        | V           |               |
| <i>Eucalyptus castrensis</i>                             | Singleton Mallee           | E1          |               |
| <i>Eucalyptus fracta</i>                                 | Broken Back Ironbark       | E1          |               |
| <i>*Eucalyptus glaucina</i>                              | Slaty Red Gum              | V           | V             |
| <i>Eucalyptus largeana</i>                               | Craven Grey Box            | E1          | E             |
| <i>*Eucalyptus parramattensis</i> subsp. <i>decadens</i> | Drooping Red Gum           | V           | V             |
| <i>*Eucalyptus pumila</i>                                | Pokolbin Mallee            | V           | V             |
| <i>*Melaleuca biconvexa</i>                              | Biconvex Paperbark         | V           | V             |
| <i>Melaleuca groveana</i>                                | Grove's Paperbark          | V           |               |
| <i>*Rhodamnia rubescens</i>                              | Scrub Turpentine           | E4A         | CE            |
| <i>*Rhodomyrtus psidioides</i>                           | Native Guava               | E4A         | CE            |
| <i>*Syzygium paniculatum</i>                             | Magenta Lilly Pilly        | E1          | V             |
| <i>*Euphrasia arguta</i>                                 |                            | E4A         | CE            |
| <i>*Persicaria elatior</i>                               | Tall Knotweed              | V           | V             |
| <i>*Banksia penicillata</i>                              |                            |             | E             |
| <i>*Grevillea parviflora</i> subsp. <i>parviflora</i>    | Small-flowered Grevillea   | V           | V             |
| <i>*Olearia cordata</i>                                  |                            | V           | V             |
| <i>*Persoonia hirsuta</i>                                | Hairy Geebung              | E1          | E             |
| <i>*Persoonia pauciflora</i>                             | North Rothbury Persoonia   | E4A         | CE            |
| <i>*Androcalva procumbens</i>                            |                            |             | V             |
| <i>*Pomaderris brunnea</i>                               | Brown Pomaderris           | E1          | V             |
| <i>Pomaderris queenslandica</i>                          | Scant Pomaderris           | E1          |               |
| <i>*Asperula asthenes</i>                                | Trailing Woodruff          | V           | V             |
| <i>*Homoranthus darwinoides</i>                          |                            | V           | V             |
| <i>*Leonema lamprophyllum</i> subsp. <i>fractum</i>      |                            | E4A         | CE            |
| <i>*Thesium australe</i>                                 | Austral Toadflax           | V           | V             |

| Scientific Name                         | Common Name                | BC Act 2016 | EPBC Act 1999 |
|-----------------------------------------|----------------------------|-------------|---------------|
| <i>Bertya mollissima</i>                | Bertya mollissima          | E1          | E             |
| <i>Maundia triglochoides</i>            |                            | V           |               |
| <i>Zannichellia palustris</i>           |                            | E1          |               |
| <b>Amphibians</b>                       |                            |             |               |
| <i>*Heleioporus australiacus</i>        | Giant Burrowing Frog       | V           | V             |
| <i>*Litoria aurea</i>                   | Green and Golden Bell Frog | E1          | V             |
| <i>*Litoria booroolongensis</i>         | Booroolong Frog            | E1          | E             |
| <i>Litoria brevipalmata</i>             | Green-thighed Frog         | V           |               |
| <i>Litoria littlejohni</i>              | Littlejohn's Tree Frog     | E1          | E             |
| <i>*Mixophyes balbus</i>                | Stuttering Frog            | E1          | V             |
| <i>*Mixophyes iteratus</i>              | Giant Barred Frog          | V           | V             |
| <i>Pseudophryne australis</i>           | Red-crowned Toadlet        | V           |               |
| <b>Reptiles</b>                         |                            |             |               |
| <i>*Aprasia parapulchella</i>           | Pink-tailed Worm-lizard    | V           | V             |
| <i>*Delma vescolineata</i>              | Hunter Valley Delma        | E1          | E             |
| <i>Varanus rosenbergi</i>               | Rosenberg's Goanna         | V           |               |
| <i>*Hoplocephalus bungaroides</i>       | Broad-headed Snake         | E1          | E             |
| <b>Birds</b>                            |                            |             |               |
| <i>*Calidris acuminata</i>              | Sharp-tailed Sandpiper     |             | V & M         |
| <i>*Calidris ferruginea</i>             | Curlew Sandpiper           | E4A         | CE & M        |
| <i>*Tringa nebularia</i>                | Common Greenshank          |             | E & M         |
| <i>*Numenius madagascariensis</i>       | Eastern Curlew             |             | CE & M        |
| <i>*Charadrius leschenaultii</i>        | Greater Sand-plover        | V           | V & M         |
| <i>*Gallinago hardwickii</i>            | Latham's Snipe             | V           | V             |
| <i>*Rostratula australis</i>            | Australian Painted Snipe   | E1          | E             |
| <i>*Botaurus poiciloptilus</i>          | Australian Bittern         | E1          | E             |
| <i>Ixobrychus flavicollis</i>           | Black Bittern              | V           |               |
| <i>Ephippiorhynchus asiaticus</i>       | Black-necked Stork         | E1          |               |
| <i>Anseranas semipalmata</i>            | Magpie Goose               | V           |               |
| <i>Oxyura australis</i>                 | Blue-billed Duck           | V           |               |
| <i>Stictonetta naevosa</i>              | Freckled Duck              | V           |               |
| <i>Irediparra gallinacea</i>            | Comb-crested Jacana        | V           |               |
| <i>Burhinus grallarius</i>              | Bush Stone-curlew          | E1          |               |
| <i>Turnix maculosus</i>                 | Red-backed Button-quail    | V           |               |
| <i>Sternula albifrons</i>               | Little Tern                | E1          | M             |
| <i>Ptilinopus magnificus</i>            | Wompoo Fruit-Dove          | V           |               |
| <i>Ptilinopus regina</i>                | Rose-crowned Fruit-Dove    | V           |               |
| <i>Ptilinopus superbus</i>              | Superb Fruit-Dove          | V           |               |
| <i>*Calyptorhynchus lathamii</i>        | Glossy Black-Cockatoo      | V           | V             |
| <i>*Callocephalon fimbriatum</i>        | Gang Gang Cockatoo         | E1          | E             |
| <i>*Lathamus discolor</i>               | Swift Parrot               | E1          | CE            |
| <i>*Neophema chrysostoma</i>            | Blue-winged Parrot         | V           | V             |
| <i>Neophema pulchella</i>               | Turquoise Parrot           | V           |               |
| <i>*Polytelis swainsonii</i>            | Superb Parrot              | V           | V             |
| <i>Glossopsitta pusilla</i>             | Little Lorikeet            | V           |               |
| <i>*Aphelocephala leucopsis</i>         | Southern Whiteface         | V           | V             |
| <i>*Hirundapus caudacutus</i>           | White-throated Needletail  | V           | V & M         |
| <i>Artamus cyanopterus cyanopterus</i>  | Dusky Woodswallow          | V           |               |
| <i>*Pycnophilus floccosus</i>           | Pilotbird                  | V           | V             |
| <i>Epthianura albifrons</i>             | White-fronted Chat         | V           |               |
| <i>*Melanodryas cucullata cucullata</i> | Hooded Robin               | E1          | E             |
| <i>Petroica boodang</i>                 | Scarlet Robin              | V           |               |
| <i>Petroica phoenicea</i>               | Flame Robin                | V           |               |
| <i>*Climacteris picumnus victoriae</i>  | Brown Treecreeper          | V           | V             |
| <i>*Stagonopleura guttata</i>           | Diamond Firetail           | V           | V             |



| Scientific Name                                                                                  | Common Name                                   | BC Act 2016 | EPBC Act 1999 |
|--------------------------------------------------------------------------------------------------|-----------------------------------------------|-------------|---------------|
| <i>Pomatostomus temporalis</i> subsp. <i>temporalis</i>                                          | Grey-crowned Babbler                          | V           |               |
| <i>Chthonicola sagittata</i>                                                                     | Speckled Warbler                              | V           |               |
| * <i>Anthochaera phrygia</i>                                                                     | Regent Honeyeater                             | E4A         | CE            |
| <i>Certhionyx variegatus</i>                                                                     | Pied Honeyeater                               | V           |               |
| * <i>Grantiella picta</i>                                                                        | Painted Honeyeater                            | V           | V             |
| <i>Melithreptus gularis gularis</i>                                                              | Black-chinned Honeyeater (eastern subspecies) | V           |               |
| <i>Daphoenositta chrysoptera</i>                                                                 | Varied Sittella                               | V           |               |
| <i>Circus assimilis</i>                                                                          | Spotted Harrier                               | V           |               |
| <i>Pandion cristatus</i>                                                                         | Eastern Osprey                                | V           |               |
| <i>Lophoictinia isura</i>                                                                        | Square-tailed Kite                            | V           |               |
| * <i>Erythrorhynchus radiatus</i>                                                                | Red Goshawk                                   | E1          | E             |
| <i>Hieraaetus morphnoides</i>                                                                    | Little Eagle                                  | V           |               |
| <i>Haliaeetus leucogaster</i>                                                                    | White-bellied Sea Eagle                       | V           | M             |
| <i>Hamirostra melanosternon</i>                                                                  | Black-breasted Buzzard                        | V           |               |
| * <i>Falco hypoleucos</i>                                                                        | Grey Falcon                                   | V           | V             |
| <i>Falco subniger</i>                                                                            | Black Falcon                                  | V           |               |
| <i>Ninox connivens</i>                                                                           | Barking Owl                                   | V           |               |
| <i>Ninox strenua</i>                                                                             | Powerful Owl                                  | V           |               |
| <i>Tyto longimembris</i>                                                                         | Eastern Grass Owl                             | V           |               |
| <i>Tyto novaehollandiae</i>                                                                      | Masked Owl                                    | V           |               |
| <i>Tyto tenebricosa</i>                                                                          | Sooty Owl                                     | V           |               |
| <b>Mammals</b>                                                                                   |                                               |             |               |
| * <i>Dasyurus maculatus maculatus</i>                                                            | Spotted-tailed Quoll                          | V           | E             |
| <i>Phascogale tapoatafa</i>                                                                      | Brush-tailed Phascogale                       | V           |               |
| * <i>Phascogale cinereus</i>                                                                     | Koala                                         | E1          | E             |
| * <i>Petrogale penicillata</i>                                                                   | Brush-tailed Rock-wallaby                     | E1          | V             |
| * <i>Notamacropus parma</i>                                                                      | Parma Wallaby                                 | V           | V             |
| * <i>Potorous tridactylus tridactylus</i>                                                        | Long-nosed Potoroo                            |             | V             |
| <i>Cercartetus nanus</i>                                                                         | Eastern Pygmy-possum                          | V           |               |
| * <i>Petaurus australis</i>                                                                      | Yellow-bellied Glider                         | V           | V             |
| <i>Petaurus norfolcensis</i>                                                                     | Squirrel Glider                               | V           |               |
| * <i>Petauroides volans</i>                                                                      | Greater Glider                                | E1          | E             |
| * <i>Pseudomys novaehollandiae</i>                                                               | New Holland Mouse                             |             | V             |
| <i>Pseudomys oralis</i>                                                                          | Hastings River Mouse                          | E1          | E             |
| * <i>Pteropus poliocephalus</i>                                                                  | Grey-headed Flying-fox                        | V           | V             |
| <i>Saccolaimus flaviventris</i>                                                                  | Yellow-bellied Sheath-tail-bat                | V           |               |
| <i>Micronomus norfolkensis</i>                                                                   | Eastern Freetail-bat                          | V           |               |
| <i>Falsistrellus tasmaniensis</i>                                                                | Eastern False Pipistrelle                     | V           |               |
| <i>Miniopterus australis</i>                                                                     | Little Bentwing-bat                           | V           |               |
| <i>Miniopterus orianae oceanensis</i>                                                            | Large Bentwing-bat                            | V           |               |
| <i>Myotis macropus</i>                                                                           | Southern Myotis                               | V           |               |
| * <i>Nyctophilus corbeni</i>                                                                     | Corben's Long-eared Bat                       | V           | V             |
| <i>Scoteanax rueppellii</i>                                                                      | Greater Broad-nosed Bat                       | V           |               |
| * <i>Chalinolobus dwyeri</i>                                                                     | Large Pied Bat                                | E1          | E             |
| <i>Vespadelus troughtoni</i>                                                                     | Eastern Cave Bat                              | V           |               |
| <b>Endangered Populations</b>                                                                    |                                               |             |               |
| <i>Acacia pendula</i> (Weeping Myall) – population in the Hunter Catchment                       |                                               | E2          |               |
| <i>Cymbidium canaliculatum</i> population in the Hunter Catchment                                |                                               | E2          |               |
| <i>Diuris tricolor</i> (Pine Donkey Orchid) population in the Muswellbrook local government area |                                               | E2          |               |
| <i>Eucalyptus camaldulensis</i> (River Red Gum) – population in the Hunter Catchment             |                                               | E2          |               |
| <i>Leionema lamprophyllum</i> subsp. <i>obovatum</i> in the Hunter Catchment                     |                                               | E2          |               |
| <i>Spyridium burragorang</i> – in the Cessnock local government area                             |                                               | E2          |               |

| Scientific Name                                                                                                                                                                                                                                              | Common Name | BC Act 2016 | EPBC Act 1999 |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------|-------------|---------------|
| <b>Endangered Ecological Communities</b>                                                                                                                                                                                                                     |             |             |               |
| Central Hunter Grey Box-Ironbark Woodland in the New South Wales North Coast and Sydney Basin Bioregions                                                                                                                                                     |             | E3          |               |
| Central Hunter Ironbark-Spotted Gum-Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions                                                                                                                                           |             | E3          |               |
| Central Hunter Valley eucalypt forest and woodland                                                                                                                                                                                                           |             |             | CE            |
| Coolibah - Black Box Woodlands of the Darling Riverine Plains and the Brigalow Belt South Bioregions                                                                                                                                                         |             |             | E             |
| Freshwater Wetlands on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions                                                                                                                                 |             | E3          |               |
| Hunter Floodplain Red Gum Woodland in the NSW North Coast and Sydney Basin Bioregions                                                                                                                                                                        |             | E3          |               |
| Hunter Lowland Redgum Forest in the Sydney Basin and New South Wales North Coast Bioregions                                                                                                                                                                  |             | E3          |               |
| Hunter Valley Foothills Slaty Gum Woodland in the Sydney Basin Bioregion                                                                                                                                                                                     |             | V2          |               |
| Hunter Valley Vine Thicket in the NSW North Coast and Sydney Basin Bioregions                                                                                                                                                                                |             | E3          |               |
| Hunter Valley Weeping Myall (Acacia pendula) Woodland                                                                                                                                                                                                        |             |             | CE            |
| Hunter Valley Weeping Myall Woodland in the Sydney Basin Bioregion                                                                                                                                                                                           |             | E4B         |               |
| Kurri Sand Swamp Woodland in the Sydney Basin Bioregion                                                                                                                                                                                                      |             | E3          |               |
| Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW North Coast Bioregions                                                                                                                                                                  |             | E3          |               |
| Lower Hunter Valley Dry Rainforest in the Sydney Basin and NSW North Coast Bioregions                                                                                                                                                                        |             | V2          |               |
| Lowland Rainforest in the NSW North Coast and Sydney Basin Bioregions                                                                                                                                                                                        |             | E3          |               |
| Lowland Rainforest of Subtropical Australia                                                                                                                                                                                                                  |             |             | CE            |
| Lowland Rainforest on Floodplain in the New South Wales North Coast Bioregion                                                                                                                                                                                |             | E3          |               |
| Quorrobolong Scribbly Gum Woodland in the Sydney Basin Bioregion                                                                                                                                                                                             |             | E3          |               |
| River-flat eucalypt forest on coastal floodplains of southern New South Wales and eastern Victoria                                                                                                                                                           |             |             | CE            |
| River-Flat Eucalypt Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions                                                                                                                          |             | E3          |               |
| Subtropical Coastal Floodplain Forest of the New South Wales North Coast Bioregion                                                                                                                                                                           |             | E3          |               |
| Subtropical eucalypt floodplain forest and woodland of the New South Wales North Coast and South East Queensland bioregions                                                                                                                                  |             |             | E             |
| Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions                                                                                                                                                |             | E3          |               |
| Sydney Freshwater Wetlands in the Sydney Basin Bioregion                                                                                                                                                                                                     |             | E3          |               |
| Warkworth Sands Woodland in the Sydney Basin Bioregion                                                                                                                                                                                                       |             | E3          |               |
| Warkworth Sands Woodland of the Hunter Valley                                                                                                                                                                                                                |             |             | CE            |
| Weeping Myall Woodlands                                                                                                                                                                                                                                      |             | E           |               |
| White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and |             | E4B         |               |
| White Box-Yellow Box-Blakely's Red Gum Grassy Woodland and Derived Native Grassland                                                                                                                                                                          |             |             | CE            |

**E1/E=Endangered Species   E2=Endangered Population   E3=Endangered Ecological Community**  
**V=Vulnerable Species   V2= Vulnerable Ecological Community   E4A/E4B/CE=Critically Endangered**  
**M=Migratory Species**

## 5.0 RESULTS

### 5.1 FLORA ASSEMBLAGES

Vegetation within the REZ Project area ranged from highly disturbed land that had been historically cleared and subject to prolonged grazing and coal mining impacts to areas of woodland and open forest that were relatively intact. Most of the vegetation within the required 132kV subtransmission line easement was contained within the existing Ausgrid easement and had been kept low to the ground by routine maintenance. The required easement for the 132kV subtransmission line will require some additional impacts in certain areas. These impacts are mostly in the form of tree branch trimming to widen the easement; however, some tree removal will be required. Areas impacted by the Muswellbrook STS upgrade and new sections of easement such as an area between the Kurri STS and Hunter Expressway, along the New England Highway at Lower Belford and along the Mitchell Line of Road will require removal of native vegetation.

The vegetation assemblages along the HEZ were stratified where possible by assigning the vegetation to Plant Community Types (PCTs) detailed in the NSW Vegetation Information System (VIS) classification database. A total of 11 Plant Community Types were identified: these were:

- 01. PCT 3631 Kurri Sand Clay Woodland
- 02. PCT 3433 Hunter Coast Foothills Spotted Gum-Ironbark Forest
- 03. PCT 3634 Quorrobolong Sand Flats Forest
- 04. PCT 3444 Lower Hunter Spotted Gum-Ironbark Forest
- 05. PCT 3446 Lower North Foothills Spotted Gum-Ironbark Grassy Forest
- 06. PCT 4023 Coastal Valleys Riparian Forest
- 07. PCT 3315 Central Hunter Ironbark-Spotted Gum Forest
- 08. PCT 4015 Central Hunter Swamp Oak Riparian Forest
- 09. PCT 4073 Lower North Hinterland River Oak Forest
- 10. PCT 3431 Central Hunter Ironbark Grassy Woodland
- 11. PCT 3525 Upper Hunter Box-Blakely's Red Gum Grassy Forest

These PCT's within the REZ varying conditions from derived grassland to open forest. A description of each PCT has been given in Tables 5.1 – 5.11. The vegetation assemblages have been mapped and are shown in Appendix B. A full list of the flora species recorded during the fieldwork is listed in Appendix D.

**Table 5.1 PCT 3631 Kurri Sand Clay Woodland**

| PCT 3631 Kurri Sand Clay Woodland               |                                                                                                                                                                                                                                                                                                                                                                                                                              |
|-------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| PCT ID                                          | PCT 3631                                                                                                                                                                                                                                                                                                                                                                                                                     |
| PCT name                                        | Kurri Sand Heathy Woodland                                                                                                                                                                                                                                                                                                                                                                                                   |
| Vegetation Formation                            | Dry Sclerophyll Forests (Shrubby sub-formation)                                                                                                                                                                                                                                                                                                                                                                              |
| Vegetation Class                                | Sydney Sand Flats Dry Sclerophyll Forests                                                                                                                                                                                                                                                                                                                                                                                    |
| Area Impacted beyond existing easement          | 0.86ha                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Per cent cleared value (%)                      | 58.04                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Extent of PCT within REZ                        | This community was present between the Kurri Kurri STS and the Hunter Expressway and North-west from Hart Road.                                                                                                                                                                                                                                                                                                              |
| Description of PCT 3631 within the subject land | Common canopy species were <i>Eucalyptus parramattensis</i> subsp. <i>decadens</i> (Drooping Red Gum) and <i>Angophora bakeri</i> (Narrow-leaved Apple). Common shrub species were <i>Melaleuca nodosa</i> (Ball-Honeymyrtle), <i>Daviesia ulicifolia</i> , <i>Acacia longifolia</i> (Sydney Golden Wattle), <i>Lissanthe strigosa</i> , <i>Grevillea montana</i> and <i>Grevillea parviflora</i> subsp. <i>parviflora</i> . |
| Condition States                                | Between the Kurri STS and the Hunter Expressway there is no existing easement. This area of Kurri Sand Heathy Woodland has been subject to previous disturbance with areas of this EEC highly modified to relatively intact. Between                                                                                                                                                                                         |



## PCT 3631 Kurri Sand Clay Woodland

Hart Road and the Hunter Expressway this EEC occurs within an existing easement and is composed of low maintained grown covers with a moderate diversity. Better quality areas of Kurri Sand Clay Woodland occur either side of the easement.

BC Act Status

Kurri Sand Swamp Woodland in the Sydney Basin Bioregion

EPBC Act Status

Kurri sand swamp woodland of the Sydney Basin bioregion



Plate 01: PCT 3631 Kurri Sand Clay Woodland between Kurri STS and the Hunter Expressway.



PCT 3631 Kurri Sand Clay Woodland



Plate 02: PCT 3631 Kurri Sand Clay Woodland (West of Hart Road)



Plate 03: PCT 3631 Kurri Sand Clay Woodland (West of Hart Road)



**Table 5.2 PCT 3433 Hunter Coast Foothills Spotted Gum-Ironbark Forest**

| PCT 3433 Hunter Coast Foothills Spotted Gum-Ironbark Forest |                                                                                                                                                                                                                                                                                                                                                                           |
|-------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| PCT ID                                                      | PCT 3433                                                                                                                                                                                                                                                                                                                                                                  |
| PCT name                                                    | Hunter Coast Foothills Spotted Gum-Ironbark Forest                                                                                                                                                                                                                                                                                                                        |
| Vegetation Formation                                        | Dry Sclerophyll Forests (Shrubby sub-formation)                                                                                                                                                                                                                                                                                                                           |
| Vegetation Class                                            | Hunter-Macleay Dry Sclerophyll Forest                                                                                                                                                                                                                                                                                                                                     |
| Per cent cleared value (%)                                  | 68.60                                                                                                                                                                                                                                                                                                                                                                     |
| Area Impacted beyond existing easement                      | 0.31ha                                                                                                                                                                                                                                                                                                                                                                    |
| Extent of PCT within REZ                                    | Sections of PCT 3433 were present along the Hunter Expressway near Kurri Kurri, NW of Harts Road.                                                                                                                                                                                                                                                                         |
| Description of PCT 3433 within the REZ                      | The dominant canopy species were <i>Corymbia maculata</i> (Spotted Gum) and <i>Eucalyptus fibrosa</i> (Broad-leaved Ironbark). Common shrub species were <i>Melaleuca nodosa</i> (Ball-Honeymyrtle), <i>Daviesia ulicifolia</i> , <i>Acacia longifolia</i> (Sydney Golden Wattle), <i>Bursaria spinosa</i> (Blackthorn) and <i>Dillwynia retorta</i> (Heathy Parrot Pea). |
| Condition States                                            | PCT 3433 was found in a variety of condition states. Areas of PCT are largely located along the existing easement with canopy species present along the edge of the easement and derived native low groundcovers within the easement.                                                                                                                                     |
| BC Act Status                                               | Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW North Coast Bioregions                                                                                                                                                                                                                                                                               |
| EPBC Act Status                                             | Not listed                                                                                                                                                                                                                                                                                                                                                                |



**Plate 04: PCT 3433 Hunter Coast Foothills Spotted Gum-Ironbark Forest just NW of Hart Road.**

PCT 3433 Hunter Coast Foothills Spotted Gum-Ironbark Forest



Plate 05: PCT 3433 Hunter Coast Foothills Spotted Gum-Ironbark Forest just NW of Hart Road.

Table 5.3 PCT 3634 Quorrobolong Sand Flats Forest

| PCT 3634 Quorrobolong Sand Flats Forest |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|-----------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| PCT ID                                  | PCT 3634                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| PCT name                                | Quorrobolong Sand Flats Forest                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Vegetation Formation                    | Dry Sclerophyll Forests (Shrubby sub-formation)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Vegetation Class                        | Sydney Sand Flats Dry Sclerophyll Forests                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Per cent cleared value (%)              | 78.12                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| Area Impacted beyond existing easement  | 0.17ha                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Extent of PCT within REZ                | PCT 3634 Quorrobolong Sand Flats Forest was present along Scales Road at Loxford. <b>Points 6-13.</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| Description of PCT 3634 within the REZ  | <p>Common canopy species were <i>Angophora floribunda</i> (Rough-barked Apple), <i>Eucalyptus punctata</i> (Grey Gum) and <i>Eucalyptus tereticornis</i> (Forest Red Gum). Native shrub species present were <i>Melaleuca nodosa</i>, <i>Acacia parvipinnula</i>, <i>Acacia longifolia</i>, <i>Acacia ulicifolia</i>, <i>Breynia oblongifolia</i> <i>Persoonia linearis</i> and <i>Bursaria spinosa</i>.</p> <p>Common native ground covers were <i>Imperata cylindrica</i> (Blady Grass), <i>Microlaena stipoides</i> (Weeping Meadow Grass), <i>Lomandra longifolia</i>, <i>Pteridium esculentum</i> (Bracken) and <i>Cheilanthes sieberi</i> (Mulga Fern).</p> <p>Common introduced species were <i>Lantana camara</i> (Lantana), <i>Anthoxanthum odoratum</i> (Sweet Vernal Grass), <i>Hyparrhenia hirta</i> (Coolatai Grass), <i>Coreopsis lanceolata</i> (Coreopsis) and <i>Ambrosia artemisiifolia</i> (Annual Ragweed).</p> |
| Condition States                        | PCT within the REZ varying conditions from derived grassland to open forest.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| BC Act Status                           | Hunter Lowland Redgum Forest in the Sydney Basin and New South Wales North Coast Bioregions                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| EPBC Act Status                         | NA                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |



PCT 3634 Quorrobolong Sand Flats Forest



**Plate 06: PCT 3634 Quorrobolong Sand Flats Forest along Scales Road, Loxford.**



**Plate 07: PCT 3634 Quorrobolong Sand Flats Forest along Scales Road, Loxford.**



**Table 5.4 PCT 3444 Lower Hunter Spotted Gum-Ironbark Forest**

| PCT 3444 Lower Hunter Spotted Gum-Ironbark Forest |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|---------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| PCT ID                                            | PCT 3444                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| PCT name                                          | Hunter Coast Foothills Spotted Gum-Ironbark Grassy Forest                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| Vegetation Formation                              | Dry Sclerophyll Forests (Shrubby sub-formation)                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Vegetation Class                                  | Hunter-Macleay Dry Sclerophyll Forest                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Per cent cleared value (%)                        | 62.33                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Area Impacted beyond existing easement            | 0.22ha                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| Extent of PCT within REZ                          | East of Old Maitland Road.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| Description of PCT 3631 within the subject land   | <p>The dominant canopy species were <i>Corymbia maculata</i> (Spotted Gum) and <i>Eucalyptus fibrosa</i> (Broad-leaved Ironbark).</p> <p>Common shrub species were <i>Melaleuca nodosa</i> (Ball-Honeymyrtle), <i>Melaleuca nodosa</i>, <i>Melaleuca stypheloides</i>, <i>Acacia parvipinnula</i>, <i>Acacia longifolia</i>, <i>Acacia ulicifolia</i>, <i>Bursaria spinosa</i>, <i>Breynia oblongifolia</i> and <i>Persoonia linearis</i>.</p> <p>Groundcovers <i>Aristida vagans</i>, <i>Entolasia stricta</i>,.</p> |
| Condition States                                  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| BC Act Status                                     | Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW North Coast Bioregions                                                                                                                                                                                                                                                                                                                                                                                                                           |
| EPBC Act Status                                   | NA                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |



**Plate 08: PCT 3444 Lower Hunter Spotted Gum-Ironbark Forest (east of Old Maitland Road) (Point 16).**



**Table 5.5 PCT 3446 Lower North Foothills Ironbark-Box-Gum Grassy Forest**

| PCT 3446 Lower North Foothills Ironbark-Box-Gum Grassy Forest |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
|---------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| PCT ID                                                        | PCT 3446                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| PCT name                                                      | Lower North Foothills Ironbark-Box-Gum Grassy Forest                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Vegetation Formation                                          | Dry Sclerophyll Forests (Shrubby sub-formation)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| Vegetation Class                                              | Hunter-Macleay Dry Sclerophyll Forest                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Per cent cleared value (%)                                    | 74.93                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Area Impacted beyond existing easement                        | 0ha                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Extent of PCT within REZ                                      | Keinbah – Data Point 26                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Description of PCT 3631 within the subject land               | <p>The dominant canopy species were <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Eucalyptus moluccana</i> (Grey Box) and <i>Eucalyptus tereticornis</i> (Forest Red Gum).</p> <p>Shrub species were <i>Acacia falcata</i>, <i>Acacia implexa</i>, <i>Breynia oblongifolia</i>, <i>Leucopogon juniperinus</i> and <i>Persoonia linearis</i>.</p> <p>Common native ground covers were <i>Cymbopogon refractus</i>, <i>Lobelia purpurascens</i>, <i>Cheilanthes sieberi</i> subsp. <i>sieberi</i>, <i>Themeda triandra</i>, <i>Microlaena stipoides</i>, <i>Dichondra repens</i> and <i>Lomandra multiflora</i> subsp. <i>multiflora</i></p> |
| Condition States                                              | Highly disturbed within easement.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| BC Act Status                                                 | Hunter Lowland Redgum Forest in the Sydney Basin and New South Wales North Coast Bioregions                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| EPBC Act Status                                               | Central Hunter Valley eucalypt forest and woodland                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |



**Plate 09: PCT 3446 Lower North Foothills Ironbark-Box-Gum Grassy Forest (Point 26).**

**Table 5.6 PCT 4023 Coastal Valley Riparian Forest**

| PCT 4023 Coastal Valley Riparian Forest         |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|-------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| PCT ID                                          | PCT 4023                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| PCT name                                        | Coastal Valley Riparian Forest                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Vegetation Formation                            | Forested Wetlands                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Vegetation Class                                | Coastal Floodplain Wetlands                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Per cent cleared value (%)                      | 78.22                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Area Impacted beyond existing easement          | 0.08ha                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| Extent of PCT within REZ                        | Found adjacent to drainage lines such as Black Creek & Jump up Creek at Belford.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Description of PCT 4023 within the subject land | <p><i>Casuarina glauca</i> (Swamp Oak) was the dominant canopy species. Included other canopy species <i>Eucalyptus tereticornis</i> (Forest Red Gum) and <i>Eucalyptus moluccana</i> (Grey Box).</p> <p>Common native ground covers were <i>Microlaena stipoides</i> (Weeping Meadow Grass), <i>Cynodon dactylon</i> (Couch), <i>Austrostipa verticillata</i>, <i>Dichondra repens</i> (Kidney Weed) <i>Lomandra longifolia</i> and <i>Lobelia purpurescens</i>.</p> <p>Common introduced species were <i>Cardiospermum grandiflorum</i> (Balloon Vine), <i>Tradescantia fluminensis</i> (Trad).</p> |
| Condition States                                | Low-Moderate High Weed Infestation.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| BC Act Status                                   | Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| EPBC Act Status                                 | Swamp Oak Floodplain Forest                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |



**Plate 10: PCT 4023 Coastal Valley Riparian Forest (Black Creek crossing) (Point 90).**



PCT 4023 Coastal Valley Riparian Forest



Plate 11: PCT 4023 Coastal Valley Riparian Forest (Jump up Creek crossing) (Point 118).

Table 5.7 PCT 3315 Central Hunter Ironbark-Spotted Gum Forest

| PCT 3315 Central Hunter Ironbark-Spotted Gum Forest |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|-----------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| PCT ID                                              | PCT 3315                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| PCT name                                            | Central Hunter Ironbark-Spotted Gum Forest                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Vegetation Formation                                | Grassy Woodlands                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Vegetation Class                                    | Coastal Valley Grassy Woodlands                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| Per cent cleared value (%)                          | 77.70                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| Area Impacted beyond existing easement              | 13.27ha                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Extent of PCT within REZ                            | Branxton, Huntlee, Lower Belford, Singleton Army Base & Mount Thorley                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| Description of PCT 3315 within the subject land     | <p>Canopy species were <i>Corymbia maculata</i> (Spotted Gum) <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Eucalyptus moluccana</i> (Grey Box) and <i>Eucalyptus fibrosa</i> (Broad-leaved Ironbark), <i>Eucalyptus tereticornis</i> (Forest Red Gum) <i>Allocasuarina luehmannii</i> (Bulloak) was a common mid storey species.</p> <p>Common shrub species were <i>Acacia parvipinnula</i>, <i>Acacia falcata</i>, <i>Melaleuca decora</i>, <i>Melaleuca nodosa</i>, <i>Pultenaea spinosa</i>, <i>Persoonia linearis</i>, <i>Grevillea montana</i>, <i>Lissanthe strigosa</i>, <i>Daviesia ulicifolia</i>, <i>Breynia oblongifolia</i>, <i>Melichrus urceolatus</i>, <i>Hibbertia pedunculata</i>, <i>Phyllanthus hirtellus</i></p> <p>Common groundcovers were <i>Cymbopogon refractus</i>, <i>Aristida racemosa</i>, <i>Aristida vagans</i>, <i>Lomandra multiflora</i> subsp. <i>multiflora</i>, <i>Cheilanthes sieberi</i> subsp. <i>sieberi</i> and <i>Dianella revoluta</i>.</p> |
| Condition States                                    | <p>Moderate-Good within new proposed easements at Lower Belford &amp; Mount Thorley. Derived Grassland within existing easements – Low-Good Condition on edges of the easement.</p> <p>A large portion of the trees that will be impacted for the new proposed easement along the southern side of the New England Highway at Lower Belford were planted as part of rehabilitation for the road duplication undertaken in the early</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |



**PCT 3315 Central Hunter Ironbark-Spotted Gum Forest**

|                 |                                                                                                                    |
|-----------------|--------------------------------------------------------------------------------------------------------------------|
|                 | 1990's. This was evidenced by the remains of plastic tree guard around the bases of a number of trees.             |
| BC Act Status   | Central Hunter Ironbark-Spotted Gum-Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions |
| EPBC Act Status | Central Hunter Valley Eucalypt Forest and Woodland                                                                 |



**Plate 12: PCT 3315 Central Hunter Ironbark-Spotted Gum Forest (Point 69)**



**Plate 13: PCT 3315 Central Hunter Ironbark-Spotted Gum Forest (Point 44)**



PCT 3315 Central Hunter Ironbark-Spotted Gum Forest



Plate 14: PCT 3315 Central Hunter Ironbark-Spotted Gum Forest (Point 85)



Plate 15: PCT 3315 Central Hunter Ironbark-Spotted Gum Forest (Point 342) Lower Belford along southern side of the New England Highway.



PCT 3315 Central Hunter Ironbark-Spotted Gum Forest



Plate 16: PCT 3315 Central Hunter Ironbark-Spotted Gum Forest (Point 347) Lower Belford along southern side of the New England Highway.



Plate 17: Tree with remains of plastic guard along New England Highway, Lower Belford.



PCT 3315 Central Hunter Ironbark-Spotted Gum Forest



Plate 18: PCT 3315 Central Hunter Ironbark-Spotted Gum Forest (Point 265) Mitchell Line of Road, Mount Thorley, NSW.



Plate 19: PCT 3315 Central Hunter Ironbark-Spotted Gum Forest (Point 266) Mitchell Line of Road, Mount Thorley, NSW.



PCT 3315 Central Hunter Ironbark-Spotted Gum Forest



Plate 20: PCT 3315 Central Hunter Ironbark-Spotted Gum Forest (Point 266) Mitchell Line of Road, Mount Thorley, NSW.

Table 5.8 PCT 4015 Central Hunter Riparian Forest

| PCT 4015 Central Hunter Riparian Forest         |                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|-------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| PCT ID                                          | PCT 4015                                                                                                                                                                                                                                                                                                                                                                                                                               |
| PCT name                                        | Central Hunter Riparian Forest                                                                                                                                                                                                                                                                                                                                                                                                         |
| Vegetation Formation                            | Forested Wetlands                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Vegetation Class                                | Coastal Floodplain Wetlands                                                                                                                                                                                                                                                                                                                                                                                                            |
| Per cent cleared value (%)                      | 87.87                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Area Impacted beyond existing easement          | 0.1ha                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Extent of PCT within REZ                        | Found adjacent to drainage lines near the Singleton Army Base. Points 145, 146 Muds Creek Crossing.                                                                                                                                                                                                                                                                                                                                    |
| Description of PCT 4015 within the subject land | <p><i>Casuarina glauca</i> (Swamp Oak) was the dominant canopy species. Very little mid stratum was present.</p> <p>Native ground covers were <i>Austrostipa verticillata</i>, <i>Microlaena stipoides</i> (Weeping Meadow Grass), <i>Cynodon dactylon</i> (Couch) and <i>Lobelia purpurascens</i> (White Root).</p> <p>The invasive climber <i>Anredera cordifolia</i> (Madeira Vine) was very common at the Muds Creek crossing.</p> |
| Condition States                                | Low-Moderate – High weed infestation.                                                                                                                                                                                                                                                                                                                                                                                                  |
| BC Act Status                                   | Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions                                                                                                                                                                                                                                                                                                                          |
| EPBC Act Status                                 | Swamp Oak Floodplain Forest                                                                                                                                                                                                                                                                                                                                                                                                            |



PCT 4015 Central Hunter Riparian Forest



Plate 21: PCT 4015 Central Hunter Riparian Forest (Muds Creek Golden Highway) (Point 145)

Table 5.9 PCT 4073 Lower North Hinterland River Oak Forest


| PCT 4015 Central Hunter Riparian Forest         |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|-------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| PCT ID                                          | PCT 4073                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| PCT name                                        | Coastal Valley Riparian Forest                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Vegetation Formation                            | Forested Wetlands                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Vegetation Class                                | Eastern Riverine Forests                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Per cent cleared value (%)                      | 80.40                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| Area Impacted beyond existing easement          | 0.73ha                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| Extent of PCT within REZ                        | Found along Bowmans and Glennies Creek crossings.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Description of PCT 4073 within the subject land | <p><i>Casuarina cunninghamiana</i> subsp. <i>cunninghamiana</i> (River Oak) was the dominant canopy species. Other native tree species included <i>Casuarina glauca</i> (Swamp Oak), <i>Melia azedarach</i> (White Cedar), <i>Angophora floribunda</i> (Rough-barked Apple).</p> <p>Native ground covers were <i>Austrostipa verticillata</i>, <i>Microlaena stipoides</i> (Weeping Meadow Grass) and <i>Cynodon dactylon</i> (Couch), <i>Lomandra longifolia</i> and <i>Lobelia purpurascens</i> (White Root).</p> <p>Common introduced flora species were <i>Cestrum parqui</i> (Green Cestrum), <i>Ricinus communis</i> (Castor Oil Plant), <i>Lycium ferocissimum</i> (African Boxthorn), <i>Galium aparine</i> (Clevers), <i>Bidens pilosa</i> (Bidens).</p> |
| Condition States                                | Moderate                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| BC Act Status                                   | Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| EPBC Act Status                                 | Swamp Oak Floodplain Forest                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |



PCT 4015 Central Hunter Riparian Forest



**Plate 22: PCT 4073 Lower North Hinterland River Oak Forest Glennies Creek crossing (Point 190)**



**Plate 23: PCT 4073 Lower North Hinterland River Oak Forest (Bowmans Creek)(Point 194)**



PCT 4015 Central Hunter Riparian Forest



Plate 24: PCT 4073 Lower North Hinterland River Oak Forest (Bowmans Creek) (Point 223)



Plate 25: PCT 4073 Lower North Hinterland River Oak Forest (Bowmans Creek) (Point 225)



**Table 5.10 PCT 3431 Central Hunter Ironbark Grassy Woodland**

| PCT 3431 Central Hunter Ironbark Grassy Woodland |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|--------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| PCT ID                                           | PCT 3431                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| PCT name                                         | Central Hunter Ironbark Grassy Woodland                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Vegetation Formation                             | Dry Sclerophyll Forests (Shrub/grass sub-formation)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Vegetation Class                                 | Hunter-Macleay Dry Sclerophyll Forests                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Per cent cleared value (%)                       | 86.47                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| Area Impacted beyond existing easement           | 4.61ha                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Extent of PCT within REZ                         | In and around the Muswellbrook STS and existing easement to the NE, Ravensworth.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Description of PCT 3431 within the subject land  | <p>Areas of native vegetation within and around the proposed Muswellbrook STS and along the existing easement east to the north-east were consistent with PCT 3431 Central Hunter Ironbark Grassy Woodland. Areas of this PCT have been subject to historic native vegetation clearance and past agricultural practices particularly cattle grazing and the construction of the existing easement. A large portion of PCT 3431 within the existing easement and parts of the proposed STS was composed of a derived grassland variation of this PCT. Other areas were composed of more concentrated remnant trees and regrowth trees.</p> <p><i>Eucalyptus crebra</i> (Narrow-leaved Ironbark) was the dominant canopy species. A small number of specimens of <i>Eucalyptus moluccana</i> (Grey Box) were also noted outside the impact areas. Mid-storey species were largely absent. A small number of specimens of <i>Allocasuarina luehmannii</i> (Bulloak) were present.</p> <p>Native shrub species were uncommon within the impact areas. Species recorded were <i>Notelaea microcarpa</i> var. <i>microcarpa</i> (Native Olive), <i>Acacia paradoxa</i> (Kangaroo Wattle), <i>Acacia decora</i> (Western Silver Wattle), <i>Solanum cinereum</i> (Narrawa Burr) and <i>Maireana microphylla</i> (Eastern Cottonbush).</p> <p>Native grasses were common groundcovers over the subject land within derived and treed areas. Common native grasses were <i>Aristida ramosa</i> (Three-awn Grass), <i>Sporobolus creber</i> (Rats Tail Grass), <i>Bothriochloa decipiens</i> var. <i>decipiens</i> (Red Leg Grass), <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Chloris ventricosa</i> (Tall Windmill Grass), <i>Austrostipa verticillata</i> (Slender Bamboo Grass), <i>Digitaria divaricatissima</i> (Umbrella Grass), <i>Eriochloa pseudoacrotricha</i> (Early Spring Grass), <i>Rytidosperma fulvum</i> (Wallaby Grass), <i>Cynodon dactylon</i> (Couch), <i>Eragrostis leptostachya</i> (Paddock Lovegrass) and <i>Microlaena stipoides</i> (Weeping Meadow Grass). Other common native ground included <i>Chrysocephalum apiculatum</i> (Common Everlasting), <i>Sida corrugata</i> (Corrugated Sida), <i>Sida hackettiana</i> (Golden Rod), <i>Glycine tabacina</i>, <i>Dichondra repens</i> (Kidneyweed), <i>Einadia hastata</i> (Berry Saltbush), <i>Eremophila debilis</i> (Amulla), <i>Vittadinia cuneata</i> (Fuzzweed), <i>Commelina cyanea</i> (Scurvy Weed), <i>Stackhousia viminea</i> (Slender Stackhousia), <i>Rumex brownii</i> (Slender Dock), <i>Dianella revoluta</i> (Blueberry Lily), <i>Erodium crinitum</i> (Blue Storksbill) and <i>Cheilanthes sieberi</i> subsp. <i>sieberi</i> (Mulga Fern).</p> <p>Common introduced species included <i>Paspalum dilatatum</i> (Paspalum), <i>Lolium perenne</i> (Perennial Ryegrass), <i>Senecio madagascariensis</i> (Fireweed), <i>Sida rhombifolia</i> (Paddy's Lucerne), <i>Plantago lanceolata</i> (Plantago), <i>Gomphocarpus fruticosus</i> (Narrow-leaved Cottonbush), <i>Galenia pubescens</i> (Galenia), <i>Bidens pilosa</i> (Cobblers Pegs), <i>Verbena bonariensis</i> (Purple-top Verbena) and <i>Lycium ferocissimum</i> (African Boxthorn). <i>Galenia pubescens</i> was particularly invasive in some areas.</p> |
| Condition States                                 | Low-Moderate                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| BC Act Status                                    | Central Hunter Grey Box-Ironbark Woodland in the New South Wales North Coast and Sydney Basin Bioregions                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| EPBC Act Status                                  | Central Hunter Grey Box-Ironbark Woodland where a patch satisfies condition thresholds as per Section 1.5.3 of the Conservation Advice.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |



PCT 3431 Central Hunter Ironbark Grassy Woodland



**Plate 26: PCT 3431 Central Hunter Ironbark Grassy Woodland within impact area of the proposed new Muswellbrook STS.**



**Plate 27: PCT 3431 Central Hunter Ironbark Grassy Woodland within impact area of the proposed new Muswellbrook STS.**



PCT 3431 Central Hunter Ironbark Grassy Woodland



**Plate 28: PCT 3431 Central Hunter Ironbark Grassy Woodland within impact area of the proposed new Muswellbrook STS.**



**Plate 29: PCT 3431 Central Hunter Ironbark Grassy Woodland within impact area of the proposed new Muswellbrook STS.**



PCT 3431 Central Hunter Ironbark Grassy Woodland



Plate 30: PCT 3431 Central Hunter Ironbark Grassy Woodland within impact area of the proposed new Muswellbrook STS.



Plate 31: PCT 3431 Central Hunter Ironbark Grassy Woodland within NW impact area of the proposed new Muswellbrook STS (Note: Tree is a *Lophostemon confertus* (Brush box) which is not endemic to the area.



**Table 5.11 PCT 3525 Upper Hunter Box-Blakely's Red Gum Grassy Forest**

| PCT 3525 Upper Hunter Box-Blakely's Red Gum Grassy Forest |                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|-----------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| PCT ID                                                    | PCT 3525                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| PCT name                                                  | Upper Hunter Box-Blakely's Red Gum Grassy Forest                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Vegetation Formation                                      | Dry Sclerophyll Forests (Shrub/grass sub-formation)                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Vegetation Class                                          | North-west Slopes Dry Sclerophyll Woodlands                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| Per cent cleared value (%)                                | 58.34                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Area Impacted beyond existing easement                    | 0.59ha                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Extent of PCT within REZ                                  | Areas of PCT 3525 were only recorded north-east of the Muswellbrook STS. Points 250,251,252.                                                                                                                                                                                                                                                                                                                                                                                  |
| Description of PCT 3525 within the subject land           | <p>Common canopy species were <i>Eucalyptus blakelyi</i> (Blakely's Red Gum), <i>Eucalyptus moluccana</i> (Grey Box). Shrub species included <i>Breynia oblongifolia</i> (Breynia), <i>Notelaea microcarpa</i>, and <i>Pittosporum undulatum</i> (Sweet Pittosporum).</p> <p>Common grasses were <i>Microlaena stipoides</i> (Weeping Meadow Grass), <i>Aristida racemosa</i>, <i>Cymbopogon refractus</i> (Barbed Wire Grass) and <i>Dichondra repens</i> (Kidney Weed).</p> |
| Condition States                                          | Low-Moderate                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| BC Act Status                                             | White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grasslands (Critically Endangered under BC Act)                                                                                                                                                                                                                                                                                                                                                 |
| EPBC Act Status                                           | White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grasslands                                                                                                                                                                                                                                                                                                                                                                                      |



**Plate 32: PCT 3525 Upper Hunter Box-Blakely's Red Gum Grassy Forest north-west of Muswellbrook STS. Note tree on right No. 23 will require removal.**



### 5.1.1 THREATENED ECOLOGICAL COMMUNITIES

Twenty-one threatened ecological communities (TECs) have been recorded within the locally of the REZ the region according to both the BioNet (DPE, 2024a) and PMST databases. Results of the database search conducted for TECs are shown within Table 4.3.

Seven TEC's listed under the BC Act were identified within the REZ or in close proximity. These TEC's were:

- Kurri Sand Swamp Woodland in the Sydney Basin Bioregion
- Hunter Lowland Redgum Forest in the Sydney Basin and New South Wales North Coast Bioregions
- Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW North Coast Bioregions
- Central Hunter Ironbark – Spotted Gum – Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions
- Central Hunter Grey Box – Ironbark Woodland in the NSW North Coast and Sydney Basin Bioregions
- Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions
- White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grasslands

Table 5.12 below shows the PCT's which are consistent within the threatened Ecological Communities listed under the BC Act 2016 and National EPBC Act 1999.

The impact of the proposed REZ on threatened Ecological Communities listed under the BC Act 2016 and National EPBC Act 1999 has been undertaken in Sections 7 & 10.

**Table 5.12 PCT's WITHIN REZ ALIGNED WITH STATE AND NATIONALLY LISTED THREATENED ECOLOGICAL COMMUNITIES.**

| Plant Community Type (PCT)                                        | Threatened Ecological Community<br>BC Act 2016                                                                     | Threatened Ecological Community<br>EPBC Act 1999<br>(provided criteria are met)         |
|-------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------|
| PCT 3631 Kurri Sand Clay Woodland                                 | Kurri Sand Swamp Woodland in the Sydney Basin Bioregion                                                            | Kurri sand swamp woodland of the Sydney Basin bioregion                                 |
| PCT 3433 Hunter Coast Foothills Spotted Gum-Ironbark Forest       | Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW North Coast Bioregions                        |                                                                                         |
| PCT 3444 Lower Hunter Spotted Gum-Ironbark Forest                 |                                                                                                                    |                                                                                         |
| PCT 3634 Quorrobolong Sand Flats Forest                           | Hunter Lowland Redgum Forest in the Sydney Basin and New South Wales North Coast Bioregions                        |                                                                                         |
| PCT 3446 Lower North Foothills Spotted Gum-Ironbark Grassy Forest |                                                                                                                    |                                                                                         |
| PCT 3315 Central Hunter Ironbark-Spotted Gum Forest               | Central Hunter Ironbark-Spotted Gum-Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions | Central Hunter Valley eucalypt forest and woodland                                      |
| PCT 3431 Central Hunter Ironbark Grassy Woodland                  | Central Hunter Grey Box-Ironbark Woodland in the New South Wales North Coast and Sydney Basin Bioregions           |                                                                                         |
| PCT 4023 Coastal Valley Riparian Forest                           |                                                                                                                    |                                                                                         |
| PCT 4015 Central Hunter Riparian Forest                           | Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions      | Swamp Oak Floodplain Forest                                                             |
| PCT 3525 Upper Hunter Box-Blakely's Red Gum Grassy Forest         | White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grassland                            | White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland |

### 5.1.2 THREATENED AND RARE FLORA SPECIES

Fifty-five (55) threatened flora species have been recorded within 10km of the subject land according to the BioNet database (DPE, 2024) or are considered to have suitable habitat on the PMST database. The results of the database search conducted for threatened flora species is shown within Table 4.3.

Threatened flora species recorded during fieldwork were:

- *Eucalyptus glaucina* (Slaty Gum)
- *Eucalyptus parramattensis* subsp. *decadens* (Drooping Red Gum)
- *Grevillea parviflora* subsp. *parviflora* (Small-flowered Grevillea)
- *Diuris tricolor* (Pine Donkey Orchid)

A number of specimens of *Eucalyptus glaucina* (Slaty Gum) were present within the area around Belford and the Singleton Military Base (Plate 34).

Specimens of *Eucalyptus parramattensis* subsp. *decadens* (Drooping Red Gum) were common within areas mapped as Kurri Sand Swamp Forest. Several specimens occurring between the Kurri STS and the Hunter Expressway and across the Hunter Expressway from the Kurri STS (35) will require removal for the creation of a new easement. The locations of *E. parramattensis* subsp. *decadens* are shown in the mapping in Appendix B. Details of each tree are contained in Appendix F.

*Grevillea parviflora* subsp. *parviflora* (Small-flowered Grevillea) were found to be common within areas of existing easement containing low maintained derived Kurri Sand Swamp Forest (Plate 36). Details of each tree are contained in Appendix F.

Approximately 4 specimens of *Diuris tricolor* (Pine Donkey Orchid) were present in the REZ within the existing easement north-east of the Muswellbrook STS (Plate 37). Details of each tree are contained in Appendix F.





Plate 34: Photo of *Eucalyptus glaucina* (Slaty Red Gum) occurring along the New England Highway at Belford below Point 118.



Plate 35: Photo of *Eucalyptus parramattensis* subsp. *decadens* (Drooping Red Gum) occurring west of Hunter Expressway across from the Kurri STS.





Plate 36: Photo of *Grevillea parviflora* subsp. *parviflora* (Small-flowered Grevillea) west of Hart Road.



Plate 37: Photo of *Diuris tricolor* (Pine Donkey Orchid) North-east of the Muswellbrook STS (SW of Point 255).



### 5.1.3 ENDANGERED POPULATIONS

Four Endangered Populations are listed as potentially occurring along the proposed REZ Zone:

- *Acacia pendula* (Weeping Myall) – population in the Hunter Catchment;
- *Diuris tricolor*, the Pine Donkey Orchid population in the Muswellbrook local government area;
- *Cymbidium canaliculatum* population in the Hunter Catchment;
- *Eucalyptus camaldulensis* (River Red Gum) – population in the Hunter Catchment.

Two Endangered Populations were recorded within the proposed REZ Zone: *Diuris tricolor* (Pine Donkey Orchid) and *Acacia pendula* (Weeping Myall). As previously stated approximately 4 specimens of *Diuris tricolor* (Pine Donkey Orchid) were present in the REZ within the existing easement north-east of the Muswellbrook STS. Details of each specimen are contained in Appendix F.

Specimens of Weeping Myall were recorded at Lower Belford, Golden Highway, Warkworth and Liddell STS Hub at Antiene (Point 285). Specimens of Weeping Myall at Liddell STS had been planted within a treed belt running east-west along Hebdon Road on the northern boundary of the Lake Liddell Recreation Area (Plate 38). The specimens at Lower Belford and those particularly along the Golden Highway, Warkworth are likely naturally occurring specimens (Plate 39). All specimens identified were the non-pendulous form of *A. pendula*. A fire had relatively recently burnt through all the specimens of *A. pendula* at Warkworth (first half of 2026) leaving burnt stems (Plate 40). As the non-pendulous form of *A. pendula* readily suckers from the roots (Umwelt 2006), these plants are likely to reshoot. Two specimens of *A. pendula* are required to be removed for access to the Liddell STS works (Point 293). As these specimens have been planted as part of a treed belt the removal of these specimens would not likely be considered significant. It is thought that any suckering specimens of *A. pendula* along within the new easement at Warkworth could be protected and managed to not impact the overhead wires. Specimens recorded at Lower Belford were situated well outside the impact area. Details of each *A. pendula* specimen are contained in Appendix F.

Whilst suitable habitat for *Cymbidium canaliculatum* was present, no specimens were recorded during fieldwork. Specimens of *Eucalyptus camaldulensis* (River Red Gum) are known to occur within the vicinity of the transmission crossing of the Hunter River at the transmission crossing at Maison Dieu. As the transmission lines will be well above the canopy no specimens of *E. camaldulensis* will likely be impacted.



Plate 38: Planted line of *Acacia pendula* (Weeping Myall) below area of proposed Eastern Hub.



Plate 39: Specimens of *Acacia pendula* (Weeping Myall) along Golden Highway, Warkworth (Point 285)(12/11/2025).





Plate 40: Specimens of Weeping Myall along Golden Highway, Warkworth (Point 285) (17/06/2026) after fire.

#### 5.1.4 PRIORITY WEEDS AND WEEDS OF STATE AND NATIONAL SIGNIFICANCE

Twenty-eight (28) priority weed species listed under the Biosecurity Act 2015 were identified on site and are listed below in Table 5.14. The site lies within the Hunter Regional Weed Committee (HRWC).

Table 5.14: Priority Weeds recorded within the REZ.

| WEED SPECIES                                     | LEGAL REQUIREMENTS                                                                              | ADDITIONAL SIGNIFICANCE |
|--------------------------------------------------|-------------------------------------------------------------------------------------------------|-------------------------|
| <i>Echium plantagineum</i><br>Paterson's Curse   | General Biosecurity Duty<br>Regional Recommended Measure<br>(Hunter)                            |                         |
| <i>Asparagus aethiopicus</i><br>Ground Asparagus | General Biosecurity Duty<br>Prohibition on dealings<br>Regional Recommended Measure             |                         |
| <i>Asparagus asparagoides</i><br>Bridal Creeper  | General Biosecurity Duty<br>Prohibition on dealings<br>Regional Recommended Measure<br>(Hunter) | T                       |
| <i>Juncus acutus</i><br>Spiny Rush               | General Biosecurity Duty<br>Regional Recommended Measure<br>(Hunter)                            |                         |
| <i>Arundo donax</i><br>Giant Reed                | General Biosecurity Duty<br>Regional Recommended Measure<br>(Hunter)                            |                         |
| <i>Cortaderia selloana</i><br>Pampas Grass       | General Biosecurity Duty<br>Prohibition on dealings<br>Regional Recommended Measure<br>(Hunter) | T                       |
| <i>Eragrostis curvula</i><br>African Lovegrass   | General Biosecurity Duty<br>Regional Recommended Measure<br>(Hunter)                            | T                       |

| WEED SPECIES                                                  | LEGAL REQUIREMENTS                                                                              | ADDITIONAL SIGNIFICANCE |
|---------------------------------------------------------------|-------------------------------------------------------------------------------------------------|-------------------------|
| <i>Hyparrhenia hirta</i><br>Coolatai Grass                    | General Biosecurity Duty<br>Regional Recommended Measure                                        | T                       |
| <i>Ageratina adenophora</i><br>Crofton Weed                   | General Biosecurity Duty<br>Regional Recommended Measure<br>(Hunter)                            |                         |
| <i>Carthamus lanatus</i><br>Saffron Thistle                   | General Biosecurity Duty                                                                        |                         |
| <i>Senecio madagascariensis</i><br>Fireweed                   | General Biosecurity Duty<br>Regional Recommended Measure                                        | N                       |
| <i>Xanthium spinosum</i><br>Bathurst Burr                     | General Biosecurity Duty                                                                        |                         |
| <i>Ailanthus altissima</i><br>Tree-of-heaven                  | General Biosecurity Duty<br>Regional Recommended Measure<br>(Hunter)                            |                         |
| <i>Dolichandra unguis-cati</i><br>Cat's claw creeper          | General Biosecurity Duty<br>Prohibition on dealings<br>Regional Recommended Measure<br>(Hunter) | T                       |
| <i>Erythrina crista-galli</i><br>Cockspur Coral Tree          | General Biosecurity Duty<br>Prohibition on dealings<br>Regional Recommended Measure<br>(Hunter) |                         |
| <i>Gleditsia triacanthos</i><br>Honey Locust                  | General Biosecurity Duty<br>Regional Recommended Measure<br>(Hunter)                            |                         |
| <i>Hypericum perforatum</i><br>St. John's Wort                | General Biosecurity Duty<br>Regional Recommended Measure<br>(Hunter)                            |                         |
| <i>Galenia pubescens</i><br>Galenia                           | General Biosecurity Duty<br>Regional Recommended Measure<br>(Hunter)                            |                         |
| <i>Lycium ferocissimum</i><br>African Boxthorn                | General Biosecurity Duty<br>Regional Recommended Measure                                        | N                       |
| <i>Opuntia stricta</i><br>Prickly Pear                        | General Biosecurity Duty<br>Prohibition on dealings<br>Regional Recommended Measure             |                         |
| <i>Lantana camara</i><br>Lantana                              | General Biosecurity Duty<br>Prohibition on dealings<br>Regional Recommended Measure<br>(Hunter) | N, T                    |
| <i>Bryophyllum</i> species<br>Mother-of-millions              | General Biosecurity Duty<br>Regional Recommended Measure                                        |                         |
| <i>Rubus fruticosus</i> species aggregate<br>Blackberry       | General Biosecurity Duty<br>Prohibition on dealings<br>Regional Recommended Measure             |                         |
| <i>Heliotropium amplexicaule</i><br>Blue Heliotrope           | General Biosecurity Duty<br>Regional Recommended Measure<br>(Hunter)                            |                         |
| <i>Cinnamomum camphora</i><br>Camphor Laurel                  | General Biosecurity Duty<br>Regional Recommended Measure                                        |                         |
| <i>Olea europaea</i> subsp. <i>cuspidata</i><br>African Olive | General Biosecurity Duty<br>Regional Recommended Measure                                        | T                       |
| <i>Cotoneaster glaucophyllus</i><br>Cotoneaster               | General Biosecurity Duty<br>Regional Recommended Measure*                                       |                         |
| <i>Cestrum parqui</i><br>Green Cestrum                        | General Biosecurity Duty<br>Regional Recommended Measure<br>(Hunter)                            |                         |

T – Listed as a Threatening Process under the NSW BC Act 2016.

N – Weed of National Significance.

**\*Priorities under the Biosecurity Act 2015**

General Biosecurity Duty - any person dealing with plant matter must take measures to prevent, minimise or eliminate the biosecurity risk (as far as is reasonably practicable).

Prohibition on dealings - Must not be imported into the State or sold.

Regional Recommended Measure - Land managers mitigate the risk of the plant being introduced to their land. Land managers reduce impacts from the plant on priority assets. Land managers prevent spread from their land where feasible. The plant or parts of the plant are not traded, carried, grown or released into the environment.



## 5.2 HABITAT APPRASIAL

### 5.2.1 HABITAT DESCRIPTION AND DISTRIBUTION IN THE VICINITY

The vegetation and landforms present along the REZ offer a variety of habitats for native fauna species. The broad habitat types included Open Forest, Woodland, Grassland and Aquatic.

#### Sclerophyll Forest/Woodland

Sclerophyll Forest/Woodland occurring along the REZ route would provide suitable habitat opportunities for a variety of species. Frugivorous, nectivorous, granivorous and insectivorous birds and microchiropteran bat species would all find potential foraging resources within this complex. Hollow-bearing trees would provide nesting and roosting sites for a variety of fauna species such as arboreal mammals, microchiropteran bats, reptile and avifauna species. Hunting opportunities exist for birds of prey, given that the variable tree coverage and understorey vegetation has created a myriad of ecotones and habitat densities. Such habitat is suitable for terrestrial species including small and medium sized mammals, macropods, reptiles and potentially for some frog species adapted to drier areas.

#### Grassland

Grassland was in the form of highly disturbed grazing land to areas of derived native grassland. Grassland provides habitat for a number of avifauna species, including predominantly terrestrial species preferring open spaces, seed eating birds and several birds of prey, which may hunt over this area in search of potential prey species. Macropods may also frequent such areas whilst grazing. Some species of bats may also forage over this cleared area for insects. The scarcity of trees and shrubs along with the often limits the value of such areas for many vulnerable species, particularly some reptiles, small mammals and birds. Areas of better quality derived native grassland may favour threatened species such as the threatened *Delma vescolineata* (Hunter Valley Delma).

#### Aquatic Habitat

Rivers, streams, dams and wetlands would provide suitable habitat for a range of fish, frog, reptile, waterbird and mammal species. This area would also act as a water source for other native animals such as macropods and offer potential hunting habitat for microchiropteran bats and birds such as *Myotis macropus* (Southern Myotis) and *Haliaeetus leucogaster* (White-bellied Sea Eagle) that prefer to hunt above or around larger water bodies such as Lake Liddel.

### 5.2.2 HABITAT CORRIDORS

The REZ crossing a number of habitat corridors of various sizes and functionality. As a result of the relatively small size and narrow nature of the REZ the proposal is unlikely to have any significant impact on the movement of any threatened fauna species.

### 5.2.3 TREE SURVEY

A total of 56 significant (larger trees and/or those containing hollows) trees were recorded along the proposed REZ. Of the 56, 28 trees are likely to require removal. Seven trees will also require branch trimming. A summary of the significant tree impacts is shown in Table 5.15

**Table 5.15: Summary of the significant tree impacts**

| TREE                    | REQUIRING REMOVAL | REQUIRING BRANCH TRIMMING |
|-------------------------|-------------------|---------------------------|
| Habitat Tree            | 16                | 7                         |
| Non-habitat Tree        | 12                | 0                         |
| Total of impacted trees | 28                | 7                         |

It is recommended that tree removal be avoided wherever possible. Details of each of the 56 trees including height, diameter at breast height (DBH), coordinates and fauna habitat attributes such as hollows and are contained Appendix E.



### 5.3 HABITAT FOR SIGNIFICANT SPECIES

An assessment of habitat attributes on site has been undertaken for the significant species listed in Table 4.3. The results of the assessment using definitions shown in Table 5.16 are displayed in Table 5.17. Threatened species identified in this assessment as having potential habitat available on site have been considered further in Section 7.0 of this report.

**Table 5.16: Definitions of likelihood of occurrence criteria.**

| Likelihood of Occurrence | Threatened Fauna                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               | Threatened Flora                                                                                                                                                                                                                                                                                                                                                                            |
|--------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Unlikely                 | Suitable habitat is absent from the subject land and/or the subject land is outside of the species known distribution                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |                                                                                                                                                                                                                                                                                                                                                                                             |
| Low                      | <ul style="list-style-type: none"> <li>The species has not been recorded in the locality (10km) within the last five years; and/or</li> <li>Although suitable habitat is present in the subject land the suitable habitat is in a highly modified, limited or degraded state; and/or</li> <li>This species may be an occasional visitor, but habitat similar or of higher quality is widely distributed in the local area.</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                          | <ul style="list-style-type: none"> <li>The species has not been recorded in the locality (10km) within the last five years, and/or</li> <li>Although suitable habitat is present in the subject land the suitable habitat is in a highly modified or degraded state</li> </ul>                                                                                                              |
| Moderate                 | <ul style="list-style-type: none"> <li>The species has been recorded in the locality (10km) within the last five years; and/or</li> <li>It is unlikely to be dependent on habitat within the subject land (i.e., for breeding or important life cycle periods) or to maintain a permanent resident population. However, the species may seasonally, opportunistically or occasionally use resources within the subject land; and/or</li> <li>Although suitable habitat is present in the subject land the suitable habitat is in a moderately modified, limited or degraded state</li> </ul> <p>This category includes fauna species that were targeted by seasonal surveys and were not recorded, wide ranging species which may fly-over the site, regardless of the habitat types present and generalist species with non-specific habitat requirements</p> | <ul style="list-style-type: none"> <li>The species has been recorded in the locality (10km) within the last five years; and/or.</li> <li>Although potential habitat is present in the subject land the suitable habitat is in a moderately modified or degraded state.</li> </ul> <p>This category includes flora species that were targeted by seasonal surveys and were not recorded.</p> |
| High                     | <ul style="list-style-type: none"> <li>The species has been recorded in the locality (10km) within the last five years; and/or</li> <li>It is highly likely that the species inhabits the subject land and is dependent on identified suitable habitat (i.e., for breeding or important life cycle periods) and is likely to maintain a resident population. This includes species that are known to visit the subject land during regular seasonal movements or migration.</li> </ul>                                                                                                                                                                                                                                                                                                                                                                         | <ul style="list-style-type: none"> <li>The species has been recorded in the locality (10km) within the last five years; and/or</li> <li>It is highly likely to inhabit the subject land and is dependent on identified suitable habitat.</li> </ul>                                                                                                                                         |
| Known                    | The species was observed in the subject land during the current survey and/or was recorded during a survey conducted on the site during the last 5 years.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |                                                                                                                                                                                                                                                                                                                                                                                             |

**Table 5.17: Habitat Assessment for Significant Species (Oceanic Birds have been removed from assessment)**

| SPECIES                                                   | STATUS            |                      |      | HABITAT DESCRIPTION AND LOCALLY KNOWN POPULATIONS                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | LIKELIHOOD OF OCCURRENCE<br>WITHIN THE REZ SITE                                                                                                                              |
|-----------------------------------------------------------|-------------------|----------------------|------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                           | BC<br>Act<br>2016 | EPB<br>C Act<br>1999 | SAIL |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |                                                                                                                                                                              |
| FLORA                                                     |                   |                      |      |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |                                                                                                                                                                              |
| <i>Wollemia nobilis</i><br>Wollemi Pine                   | E4A               | CE                   | Yes  | Restricted to remote canyons in the Wollemi National Park, north-west of Sydney. Occurs in warm temperate rainforest and rainforest margins in remote sandstone canyons.                                                                                                                                                                                                                                                                                                                                                                                     | <b>Unlikely</b><br>No suitable habitat was present. Outside known distribution.                                                                                              |
| <i>Caladenia tessellata</i><br>Thick-lipped Spider-orchid | V                 | V                    | Yes  | Generally found in grassy sclerophyll woodland on clay loam or sandy soils, though the population near Braidwood is in low woodland with stony soil. Is known from the Sydney area (old records), Wyong, Ulladulla and Braidwood in NSW.                                                                                                                                                                                                                                                                                                                     | <b>Unlikely</b><br>Only marginal habitat present. Outside known distribution.                                                                                                |
| <i>Corybas dowlingii</i><br>Red Helmet Orchid             | E1                |                      | No   | Forms colonies and typically grows in gullies in tall open forest on well-drained gravelly soil at elevations of 10-200m. Is restricted to New South Wales where it is currently known from 4 localities including Port Stephens (2 localities), Bulahdelah and Freemans Waterhole south of Newcastle. A population is known from Stoney Ridge Reserve on Soldiers Point on the southern shore of Port Stephens.                                                                                                                                             | <b>Low</b><br>Marginal habitat confined to the far east of the REZ.                                                                                                          |
| <i>Cryptostylis hunteriana</i><br>Leafless Tongue Orchid  | V                 | V                    | No   | Grows in swamp-heath on sandy soils, chiefly in coastal districts, south from the Gibraltar Range. It is known historically from several localities on the NSW south coast and has been observed in recent years at many sites between Batemans Bay and Nowra (although it is uncommon at all sites). Also recorded at Munmorah State Conservation Area, Nelson Bay, Wyee, Washpool National Park, Nowendoc State Forest, Ku-Ring-Gai Chase National Park and Ben Boyd National Park.                                                                        | <b>Unlikely</b><br>Only marginal habitat present. Outside known distribution.                                                                                                |
| <i>Diuris pedunculata</i><br>Small Snake Orchid           | E1                | E                    | No   | Confined to northeast NSW. It was originally found scattered from Tenterfield south to the Hawkesbury River. Mainly found on the New England Tablelands, around Armidale, Uralla, Guyra and Ebor. Grows on grassy slopes or flats, often on peaty soils in moist areas. Also occurs on shale and trap soils, on fine granite, and among boulders. It flowers during August-October.                                                                                                                                                                          | <b>Unlikely</b><br>Only marginal habitat present. Outside known distribution.                                                                                                |
| <i>Diuris tricolor</i><br>Pine Donkey Orchid              | V                 |                      | No   | Sporadically distributed on the western slopes of NSW, extending from south of Narrandera all the way to the north of NSW. Localities in the south include several sites west of Wagga Wagga, Dubbo in the Central West and Pilliga National in the north. The population in the Muswellbrook LGA is at the eastern limit of the geographic range of the species. Grows in sclerophyll forest among grass with native Cypress Pine ( <i>Callitris</i> spp.). It is found in sandy soils, either on flats or small rises. Flowers from September to November. | <b>Known</b><br>Recorded within easement to the NE of the Muswellbrook STS. Suitable habitat confined to western half of REZ. More likely to be present around Muswellbrook. |



| SPECIES                                                      | STATUS            |                      |     | HABITAT DESCRIPTION AND LOCALLY KNOWN POPULATIONS                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  | LIKELIHOOD OF OCCURRENCE<br>WITHIN THE REZ SITE                                                                      |
|--------------------------------------------------------------|-------------------|----------------------|-----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------|
|                                                              | BC<br>Act<br>2016 | EPB<br>C Act<br>1999 | SAI |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |                                                                                                                      |
| <i>Prasophyllum petilum</i><br>Tarengo Leek Orchid           | E1                | E                    | No  | In NSW natural populations are known in a total of five sites. These are near Boorowa, Queanbeyan area, Ilford, Delegate and west of Muswellbrook. Grows in open sites within Natural Temperate Grassland.                                                                                                                                                                                                                                                                                                                                                                                         | <b>Low</b><br>Suitable habitat confined to western half of REZ. More likely to be present around Muswellbrook.       |
| <i>Prasophyllum</i> sp.<br>Wybong<br>A Leek Orchid           |                   | CE                   | Yes | Leek orchids are generally found in shrubby and grassy habitats in dry to wet soil (Jones 2006). Known to occur in open eucalypt woodland and grassland.                                                                                                                                                                                                                                                                                                                                                                                                                                           | <b>Low</b><br>Suitable habitat confined to western half of REZ. More likely to be present around Muswellbrook.       |
| <i>Pterostylis chaetophora</i><br>Tall Rustyhood             | V                 |                      | No  | In NSW it is currently known from 18 scattered locations in a relatively small area between Taree and Kurri Kurri, extending to the south-east towards Tea Gardens and west into the Upper Hunter, with additional records near Denman and Wingen. The preferred habitat is seasonally moist, dry sclerophyll forest with a grass and shrub understorey.                                                                                                                                                                                                                                           | <b>Low</b><br>Suitable habitat confined to the far east of the REZ.                                                  |
| <i>Pterostylis gibbosa</i><br>Illawarra Greenhood            | E1                | E                    | No  | All known sub-populations occur in open forest and woodland on flat or gently sloping land with poorly drained soils. Within the Hunter Valley this orchid species is confined to the Milbrodale area.                                                                                                                                                                                                                                                                                                                                                                                             | <b>Unlikely</b><br>Only marginal habitat present. Outside known distribution.                                        |
| <i>Rhizanthella slateri</i><br>Eastern<br>Underground Orchid | V                 | E1                   | Yes | Occurs from south-east Queensland to south-east NSW. In NSW, currently known from fewer than 10 locations, including near Bulahdelah, the Watagan Mountains, the Blue Mountains, Wiseman's Ferry area, Agnes Banks and near Nowra. Grows in sclerophyll forest in shallow to deep loams.                                                                                                                                                                                                                                                                                                           | <b>Low</b><br>Marginal habitat was present within better quality areas of native vegetation. No known local records. |
| <i>Thelymitra adorata</i><br>Wyong Sun Orchid                | E4A               | CE                   | Yes | Currently known from several local government areas (LGA) within the Central Coast Council region of New South Wales. These include but are not limited to the LGAs of Wyong, Warnervale and Wyongah. Also recorded in the southern portion of Lake Macquarie City Council area. Occurs in grassy woodland or occasionally derived grassland in well-drained clay loam or shale derived soils. The vegetation type in which the majority of populations occur (including the largest colony) is a Spotted Gum - Ironbark Forest with a diverse grassy understorey and occasional scattered shrubs. | <b>Unlikely</b><br>No suitable habitat. Outside known distribution.                                                  |
| <i>Dichanthium setosum</i><br>Blue Grass                     | V                 | V                    | No  | Occurs on the New England Tablelands, Northwest Slopes and Plains and the Central Western Slopes of NSW, extending to northern Queensland. Associated with heavy basaltic black soils and red-brown loams with clay subsoil.                                                                                                                                                                                                                                                                                                                                                                       | <b>Low</b><br>Suitable habitat confined to the far west of the REZ.                                                  |
| <i>Cynanchum elegans</i><br>White-flowered Wax<br>Plant      | E1                | E                    | No  | This species occurs in scattered coastal localities from the QLD-NSW border south to Wollongong. Found in dry, littoral or subtropical rainforest, and occasionally in scrub and woodland from sea level to about 600m ASL.                                                                                                                                                                                                                                                                                                                                                                        | <b>Unlikely</b><br>No likely suitable habitat.                                                                       |
| <i>Brachyscome brownii</i>                                   |                   | CE                   | No  | Endemic to New South Wales (NSW). Associated with riparian zones and sandy soils in dry sclerophyll forest.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        | <b>Low</b><br>Suitable habitat was present.                                                                          |

| SPECIES                                                       | STATUS            |                      |     | HABITAT DESCRIPTION AND LOCALLY KNOWN POPULATIONS                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      | LIKELIHOOD OF OCCURRENCE<br>WITHIN THE REZ SITE                                                      |
|---------------------------------------------------------------|-------------------|----------------------|-----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------|
|                                                               | BC<br>Act<br>2016 | EPB<br>C Act<br>1999 | SAI |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |                                                                                                      |
| <i>Ozothamnus tessellatus</i>                                 | V                 | V                    | No  | East-west zone south of Bunnan and between west Bylong and east Ravensworth. Grows in eucalypt woodland.                                                                                                                                                                                                                                                                                                                                                                                                                               | <b>Low-Moderate</b><br>Suitable habitat present west of Singleton. Known records around Ravensworth. |
| <i>Picris evae</i><br>Hawkweed                                | V                 | V                    | No  | Found in NSW north from the Inverell area, in the north-western slopes and plains regions. Main habitat is open Eucalypt Forest.                                                                                                                                                                                                                                                                                                                                                                                                       | <b>Unlikely</b><br>No likely suitable habitat.                                                       |
| <i>Rutidosia heterogama</i><br>Heath Wrinklewort              | V                 | V                    | No  | Grows in heath on sandy soils and moist areas in open forest and has been recorded along disturbed roadsides. Recorded from near Cessnock to Kurri Kurri with an outlying occurrence at Howes Valley. On the Central Coast it is located north from Wyong to Newcastle.                                                                                                                                                                                                                                                                | <b>Low-Moderate</b><br>Suitable habitat confined to the east of the REZ.                             |
| <i>Lepidium aschersonii</i><br>Spiny Peppercress              | V                 | V                    | No  | Endemic to mainland southern Australia, north-eastern New South Wales to Western Australia. Occurs in the marginal central-western slopes and north-western plains regions of NSW. Associated with periodically wet sites such as gilgai depressions and the margins of freshwater and saline marshes and shallow lakes, usually on heavy clay soil (Harris & Smith 2000).                                                                                                                                                             | <b>Unlikely</b><br>No likely suitable habitat.                                                       |
| <i>Tetradlea juncea</i><br>Black-eyed Susan                   | V                 | V                    | No  | Confined to the northern portion of the Sydney Basin bioregion and the southern portion of the North Coast bioregion in the local government areas of Wyong, Lake Macquarie, Newcastle, Port Stephens, Great Lakes and Cessnock. Found in low open forest/woodland with a mixed shrub understorey and grassy groundcover. However, it has also been recorded in heathland and moist forest.                                                                                                                                            | <b>Unlikely</b><br>No likely suitable habitat. REZ outside known distribution.                       |
| <i>Tylophora linearis</i><br>( <i>Vincetoxicum forsteri</i> ) | V                 | E                    | No  | Occurs from southern Queensland into central NSW. Grows in dry scrub and open forest.                                                                                                                                                                                                                                                                                                                                                                                                                                                  | <b>Unlikely</b><br>No likely suitable habitat.                                                       |
| <i>Acacia bynoeana</i><br>Bynoe's Wattle                      | E1                | V                    | No  | Found in heath, woodland and dry sclerophyll forests on sandy soils derived from Hawkesbury Sandstone. Associated overstorey species include Red Bloodwood, Scribbly Gum, Parramatta Red Gum, Saw Banksia and Narrow-leaved Apple but has also been recorded within Spotted Gum – Ironbark Forest at its most northerly extent in North Rothbury in the Hunter Valley. Found in central eastern NSW, from the Hunter District (Morisset, Kurri Kurri & North Rothbury) south to the Southern Highlands and west to the Blue Mountains. | <b>Low-Moderate</b><br>Suitable habitat confined to the east of the REZ.                             |
| <i>Dillwynia tenuifolia</i>                                   | V                 |                      | No  | Core distribution is the Cumberland Plain from Windsor and Penrith east to Dean Park near Colebee. Other populations in western Sydney recorded from Voyager Point and Kemps Creek in the Liverpool LGA, Luddenham in the Penrith LGA and South Maroota in the Baulkham Hills Shire. Disjunct localities outside the Cumberland Plain include the Bulga Mountains at Yengo in the north, Kurrajong Heights, and Woodford in the Lower Blue Mountains. Occurs in dry sclerophyll woodland on sandstone, shale or laterite               | <b>Unlikely</b><br>No likely suitable habitat.                                                       |



| SPECIES                                                 | STATUS            |                      |      | HABITAT DESCRIPTION AND LOCALLY KNOWN POPULATIONS                                                                                                                                                                                                                                                                                                                                                                                                                                                                          | LIKELIHOOD OF OCCURRENCE<br>WITHIN THE REZ SITE                                                             |
|---------------------------------------------------------|-------------------|----------------------|------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------|
|                                                         | BC<br>Act<br>2016 | EPB<br>C Act<br>1999 | SAIL |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |                                                                                                             |
| <i>Swainsona murrayana</i><br>Slender Darling-pea       | V                 | V                    | No   | Found throughout NSW , recorded in the Jerilderie and Deniliquin areas of the southern riverine plain, the Hay plain as far north as Willandra National Park, near Broken Hill and in various localities between Dubbo and Moree. Grows in vegetation types including bladder saltbush, black box and grassland communities. Plants have been found in remnant native grasslands or grassy woodlands that have been intermittently grazed or cultivated                                                                    | <b>Low</b><br>Marginal habitat confined to the far west of the REZ.                                         |
| <i>Prostanthera cineolifera</i><br>Singleton Mint Bush  | V                 | V                    | No   | Grows in open woodlands on exposed sandstone ridges. Usually found in association with shallow or skeletal soils. Localities include Apsley Falls, east of Walcha; St Albans and the western side of Mangrove Creek Dam, near Bucketty; a site west of North Rothbury; Pokolbin State Forest (NSW Government, 2005) and Bellbird in the Hunter Valley (DECC NSW, 2008).                                                                                                                                                    | <b>Unlikely</b><br>No likely suitable habitat.                                                              |
| <i>Angophora inopina</i><br>Charmhaven Apple            | V                 | V                    | No   | Endemic to the Central Coast region of NSW. The known northern limit is near Karuah where a disjunct population occurs; to the south populations extend from Toronto to Charmhaven with the main population occurring between Charmhaven and Morisset. There is an unconfirmed record of the species near Bulahdelah. Approximately 1250 ha of occupied habitat has been mapped in the Wyong-southern Lake Macquarie area. Grows in open woodland with a dense shrub understorey on deep white sandy soils over sandstone. | <b>Unlikely</b><br>No likely suitable habitat.                                                              |
| <i>Callistemon linearifolius</i><br>Netted Bottle Brush | V                 |                      | No   | Grows in dry sclerophyll forest on the coast and adjacent ranges. From the Georges River to the Hawkesbury River in the Sydney area, and north to Nelson Bay. Known to occur within Stony Ridge Reserve on Soldiers Point.                                                                                                                                                                                                                                                                                                 | <b>Moderate-High</b><br>Suitable habitat within the eastern section of the REZ. Known records close to REZ. |
| <i>Eucalyptus castrensis</i><br>Singleton Mallee        | E1                |                      | Yes  | Occurs on a low broad ridgetop on loam over sandstone. The understorey consists of grasses and scattered shrubs, with bare ground and litter. Known only from a single dense stand within the Singleton Military Area. Here it is locally dominant stand over about ten hectares with a number of smaller outlying stands over a 2.5 km range.                                                                                                                                                                             | <b>Unlikely</b><br>No suitable habitat was present.                                                         |
| <i>Eucalyptus fracta</i><br>Broken Back Ironbark        | E1                |                      | Yes  | Occurs in dry eucalypt woodland in shallow soils. Associated species in slightly deeper soils include <i>Eucalyptus sparsifolia</i> , <i>E. punctata</i> , <i>Corymbia maculata</i> and <i>Angophora euryphylla</i> . Confined largely to State Forest. Locally common but restricted to the northern Broken Back Range near Cessnock, NSW. The dominant tree in a narrow band along the upper edge of a sandstone escarpment.                                                                                             | <b>Unlikely</b><br>No suitable habitat was present.                                                         |
| <i>Eucalyptus glaucina</i><br>Slaty Red Gum             | V                 | V                    | No   | Grows in grassy woodland and dry eucalypt forest, usually on deep, moderately fertile and well-watered soils. This species has only been recorded on the north coast of NSW and in small populations from Taree to Broke and west of Maitland.                                                                                                                                                                                                                                                                             | <b>Known</b><br>Tree recorded at several locations within or in close proximity to the REZ.                 |
| <i>Eucalyptus largeana</i><br>Craven Grey Box           | E1                | E                    | No   | Confined to Gloucester-Craven district and near Pokolbin, although a number of unsubstantiated records exist from outside the currently accepted range. Populations                                                                                                                                                                                                                                                                                                                                                        | <b>Unlikely</b>                                                                                             |

| SPECIES                                                                     | STATUS            |                      |     | HABITAT DESCRIPTION AND LOCALLY KNOWN POPULATIONS                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 | LIKELIHOOD OF OCCURRENCE<br>WITHIN THE REZ SITE                                                                                                       |
|-----------------------------------------------------------------------------|-------------------|----------------------|-----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                             | BC<br>Act<br>2016 | EPB<br>C Act<br>1999 | SAI |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |                                                                                                                                                       |
|                                                                             |                   |                      |     | are known from Copeland Tops State Conservation Area and Berrico Nature Reserve, with unconfirmed records from Talawahl and Glen Nature Reserves and Willi Willi National Park. The majority of remaining populations occur on private lands and roadsides, often as single trees or small clumps interspersed with other tree species. Occurs in wet forest on sloping sites in subcoastal ranges.                                                                                                                                                               | No suitable habitat was present. Outside known distribution.                                                                                          |
| <i>Eucalyptus parramattensis</i> subsp. <i>decadens</i><br>Drooping Red Gum | V                 | V                    | No  | Generally, occupies deep, low-nutrient sands, often those subject to periodic inundation or where water tables are relatively high. It occurs in dry sclerophyll woodland with dry heath understorey. It also occurs as an emergent in dry or wet heathland. Often where this species occurs, it is a community dominant. In the Kurri Kurri area, <i>E. parramattensis</i> subsp. <i>decadens</i> is a characteristic species of 'Kurri Sand Swamp Woodland and in the Tomago Sandbeds area, the species is usually associated with the 'Tomago Swamp Woodland'. | <b>Known</b><br>Tree species recorded at several locations within or in close proximity to the REZ. Found within areas of Kurri Sand Heathy Woodland. |
| <i>Eucalyptus pumila</i><br>Pokolbin Mallee                                 | V                 | V                    | Yes | Present as a mid-canopy species to a height of 6 m within dry sclerophyll woodland which has a canopy comprising <i>Eucalyptus fibrosa</i> , <i>Callitris endlicheri</i> and, to a lesser extent, <i>Corymbia maculata</i> . Very restricted distribution. Occupies a north-west facing slope of the Broken Back Range west of Pokolbin.                                                                                                                                                                                                                          | <b>Unlikely</b><br>No suitable habitat was present.                                                                                                   |
| <i>Melaleuca biconvexa</i><br>Biconvex Paperbark                            | V                 | V                    | No  | Only found in NSW, with scattered and dispersed populations found in the Jervis Bay area in the south and the Gosford-Wyong area in the north. Grows in damp places, often near streams; coastal districts and adjacent tablelands from Jervis Bay north to the Port Macquarie district.                                                                                                                                                                                                                                                                          | <b>Low</b><br>Only marginal habitat was present around low drainage areas in the far east of the REZ.                                                 |
| <i>Melaleuca groveana</i><br>Grove's Paperbark                              | V                 |                      | No  | Widespread, scattered populations in coastal districts north of Yengo National Park to southeast Queensland. Grows in heath and shrubland, often in exposed sites, in low coastal hills, escarpment ranges and tablelands on outcropping granite, rhyolite and sandstone on rocky outcrops and cliffs. It also occurs in dry shrubby open forest and woodlands                                                                                                                                                                                                    | <b>Unlikely</b><br>No suitable habitat was present. Outside known distribution.                                                                       |
| <i>Rhodamnia rubescens</i><br>Scrub Turpentine                              | E4A               | CE                   | Yes | Occurs in coastal districts north from Batemans Bay in New South Wales, approximately 280 km south of Sydney, to areas inland of Bundaberg in Queensland. Found in littoral, warm temperate and subtropical rainforest and wet sclerophyll forest usually on volcanic and sedimentary soils.                                                                                                                                                                                                                                                                      | <b>Unlikely</b><br>No suitable habitat was present.                                                                                                   |
| <i>Rhodomyrtus psidioides</i><br>Native Guava                               | E4A               | CE                   | Yes | Occurs from Broken Bay New South Wales to Maryborough in Queensland. Pioneer species found in littoral, warm temperate and subtropical rainforest and wet sclerophyll forest often near creeks and drainage lines.                                                                                                                                                                                                                                                                                                                                                | <b>Unlikely</b><br>No suitable habitat was present.                                                                                                   |
| <i>Syzygium paniculatum</i><br>Magenta Lilly Pilly                          | E1                | V                    | No  | Occurs in a narrow coastal distribution in rainforests on sandy soils or stabilised coastal dunes from Jervis Bay to Bulahdelah in NSW.                                                                                                                                                                                                                                                                                                                                                                                                                           | <b>Unlikely</b><br>No suitable habitat was present.                                                                                                   |



| SPECIES                                                                              | STATUS            |                      |     | HABITAT DESCRIPTION AND LOCALLY KNOWN POPULATIONS                                                                                                                                                                                                                                                                                                                                                                                                                                                                        | LIKELIHOOD OF OCCURRENCE<br>WITHIN THE REZ SITE                                                                                                       |
|--------------------------------------------------------------------------------------|-------------------|----------------------|-----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                                      | BC<br>Act<br>2016 | EPB<br>C Act<br>1999 | SAI |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |                                                                                                                                                       |
| <i>Euphrasia arguta</i><br>Eyebright                                                 | E4A               | CE                   | Yes | Found within the Nundle area reported from eucalypt forest with a mixed grass and shrub understorey; here, plants were most dense in an open disturbed area and along the roadside, indicating the species had regenerated following disturbance.                                                                                                                                                                                                                                                                        | <b>Unlikely</b><br>No suitable habitat was present.                                                                                                   |
| <i>Persicaria elatior</i><br>Tall Knotweed                                           | V                 | V                    | No  | Recorded in south-eastern NSW (Mt Dromedary (an old record), Moruya State Forest near Turlinjah, the Upper Avon River catchment north of Robertsocaleyin, Bermagui, and Picton Lakes. In northern NSW it is known from Raymond Terrace (near Newcastle) and the Grafton area (Cherry Tree and Gibberagee State Forests). grows in damp places, especially beside streams and lakes. Occasionally in swamp forest or associated with disturbance.                                                                         | <b>Low</b><br>Only marginal habitat was present around low drainage areas in the far east of the REZ.                                                 |
| <i>Banksia penicillata</i>                                                           |                   | E                    | No  | Endemic to NSW, found on sandstone-derived soils along ridges, among rocks or near cliffs, and occasionally on lower slopes and in plateau valleys, in the northern Blue Mountains. Occurs in sclerophyll forest or woodland (occasionally heaths) on sandstone soils.                                                                                                                                                                                                                                                   | <b>Unlikely</b><br>No suitable habitat was present.                                                                                                   |
| <i>Grevillea parviflora</i><br>subsp. <i>parviflora</i><br>Small-flower<br>Grevillea | V                 | V                    | No  | Grows in sandy or light clay soils usually over thin shales. Occurs in a range of vegetation types from heath and shrubby woodland to open forest and is found over a range of altitudes from flat, low-lying areas to upper slopes and ridge crests. Common canopy species vary greatly with community type but generally are species that favour soils with a strong lateritic influence including <i>Eucalyptus fibrosa</i> , <i>E. parramattensis</i> , <i>Angophora bakeri</i> and <i>Eucalyptus sclerophylla</i> . | <b>Known</b><br>Tree species recorded at several locations within or in close proximity to the REZ. Found within areas of Kurri Sand Heathy Woodland. |
| <i>Olearia cordata</i>                                                               | V                 | V                    | No  | Scattered distribution generally restricted to the south-western Hunter Plateau, eastern Colo Plateau, and the far north-west of the Hornsby Plateau near Wisemans Ferry east of Maroota. Grows in dry open sclerophyll forest and open shrubland, on sandstone ridges.                                                                                                                                                                                                                                                  | <b>Unlikely</b><br>No suitable habitat was present.                                                                                                   |
| <i>Persoonia hirsuta</i><br>Hairy Geebung                                            | E1                | E                    | Yes | Has a scattered distribution around Sydney. The species is distributed from Singleton in the north, along the east coast to Bargo in the south and the Blue Mountains to the west. Found in sandy soils in dry sclerophyll open forest, woodland and heath on sandstone.                                                                                                                                                                                                                                                 | <b>Unlikely</b><br>No suitable habitat was present. Outside known distribution.                                                                       |
| <i>Persoonia pauciflora</i><br>North Rothbury<br>Persoonia                           | E4A               | CE                   | Yes | Found only in small area near North Rothbury in the Hunter Valley. Occurs in open forests and woodlands with a shrubby understorey on clay soils derived from silty sandstones.                                                                                                                                                                                                                                                                                                                                          | <b>Low-Moderate</b><br>Suitable habitat was present within the vicinity of Rothbury.                                                                  |
| <i>Androcalva procumbens</i>                                                         |                   | V                    | No  | In sandy sites mainly confined to the Dubbo;-Mendooran;-and Gilgandra region.                                                                                                                                                                                                                                                                                                                                                                                                                                            | <b>Unlikely</b><br>No suitable habitat was present. Outside known distribution.                                                                       |
| <i>Pomaderris brunnea</i><br>Brown Pomaderris                                        | E1                | V                    | No  | Found in a very limited area around the Colo, Nepean and Hawkesbury Rivers, including the Bargo area and near Camden. It also occurs near Walcha on the New                                                                                                                                                                                                                                                                                                                                                              | <b>Unlikely</b><br>No suitable habitat was present. Outside known distribution.                                                                       |

| SPECIES                                                 | STATUS            |                      |     | HABITAT DESCRIPTION AND LOCALLY KNOWN POPULATIONS                                                                                                                                                                                                                                                                                                                                                          | LIKELIHOOD OF OCCURRENCE<br>WITHIN THE REZ SITE                                                                                     |
|---------------------------------------------------------|-------------------|----------------------|-----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------|
|                                                         | BC<br>Act<br>2016 | EPB<br>C Act<br>1999 | SAI |                                                                                                                                                                                                                                                                                                                                                                                                            |                                                                                                                                     |
|                                                         |                   |                      |     | England tablelands and in far eastern Gippsland in Victoria. Grows in moist woodland or forest on clay and alluvial soils of flood plains and creek lines.                                                                                                                                                                                                                                                 |                                                                                                                                     |
| <i>Pomaderris queenslandica</i><br>Scant Pomaderris     | E1                |                      | No  | Known from several locations on the NSW north coast and a few locations on the New England Tablelands and North West Slopes, including near Torrington and Coolata. Found in moist eucalypt forest or sheltered woodlands with a shrubby understorey, and occasionally along creeks.                                                                                                                       | <b>Unlikely</b><br>No suitable habitat was present.<br>Outside known distribution.                                                  |
| <i>Asperula asthenes</i><br>Trailing Woodruff           | V                 | V                    | No  | Occurs only in NSW. It is found in scattered locations from Bulahdelah north to near Kempsey, with several records from the Port Stephens/Wallis Lakes area/Forster (including Myall Lakes NP, New England NP, Wallingat NP and Darawnk NR). Occurs in damp sites, often along riverbanks.                                                                                                                 | <b>Unlikely</b><br>No suitable habitat was present.<br>Outside known distribution.                                                  |
| <i>Homoranthus darwinoides</i>                          | V                 | V                    | No  | Rare in the central tablelands and western slopes of NSW, occurring from Putty to the Dubbo district. It is found west of Muswellbrook between Merriwa and Bylong, and north of Muswellbrook to Goonoo SCA. Grows in shrubby woodland, usually in deep sandy soils over sandstone.                                                                                                                         | <b>Unlikely</b><br>No suitable habitat was present.<br>Outside known distribution.                                                  |
| <i>Leionema lamprophyllum</i><br>subsp. <i>fractum</i>  | E4A               | CE                   | Yes | Currently known only from the Broken Back Range near Cessnock, with a historical collection from Munghorn Gap Nature Reserve near Wollar.                                                                                                                                                                                                                                                                  | <b>Unlikely</b><br>No suitable habitat was present.<br>Outside known distribution.                                                  |
| <i>Thesium australe</i><br>Austral Toadflax             | V                 | V                    | No  | Grows in grassland or woodland, often in damp sites.                                                                                                                                                                                                                                                                                                                                                       | <b>Low</b><br>Marginal habitat was present.                                                                                         |
| <i>Bertya mollissima</i>                                | E1                | E                    | No  | Endemic to north-eastern New South Wales (NSW), known to occur historically from Mount Kaputar, Warrumbungle and Liverpool Ranges to the Scone and Singleton districts. Occurs on steep hillsides and mountain summits in shallow sandy or gravelly soil in rock cracks and among boulders. Within heath or open woodland communities surrounded by Eucalyptus spp. between 500 to 1500 m above sea level. | <b>Unlikely</b><br>No suitable habitat was present.                                                                                 |
| <i>Maundia triglochoides</i>                            | V                 |                      | No  | Restricted to coastal NSW and extending into southern Queensland. The current southern limit is Wyong. Grows in swamps or shallow freshwater in heavy clay.                                                                                                                                                                                                                                                | <b>Unlikely</b><br>No suitable habitat was present.                                                                                 |
| <i>Zannichellia palustris</i>                           | E1                |                      | No  | In NSW, known from the lower Hunter and in Sydney Olympic Park. Grows in fresh or slightly saline stationary or slowly flowing water.                                                                                                                                                                                                                                                                      | <b>Low</b><br>Suitable habitat was confined to areas of aquatic habitat in the far east of the REZ site such as within Swamp Creek. |
| <b>FAUNA - AMPHIBIANS</b>                               |                   |                      |     |                                                                                                                                                                                                                                                                                                                                                                                                            |                                                                                                                                     |
| <i>Heleioporus australiacus</i><br>Giant Burrowing Frog | V                 | V                    | No  | Banks of semi-permanent to ephemeral sand or rock-based streams and has also been identified in dams, drainage ditches and roadside culverts.                                                                                                                                                                                                                                                              | <b>Unlikely</b><br>No suitable habitat was present.<br>Outside known distribution.                                                  |



| SPECIES                                                 | STATUS            |                      |     | HABITAT DESCRIPTION AND LOCALLY KNOWN POPULATIONS                                                                                                                                                                                                                                                                                                                                                                 | LIKELIHOOD OF OCCURRENCE<br>WITHIN THE REZ SITE                                                                                                                         |
|---------------------------------------------------------|-------------------|----------------------|-----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                         | BC<br>Act<br>2016 | EPB<br>C Act<br>1999 | SAI |                                                                                                                                                                                                                                                                                                                                                                                                                   |                                                                                                                                                                         |
| <i>Litoria aurea</i><br>Green and Golden Bell Frog      | E1                | V                    | No  | Inhabits swamps, lagoons, streams and ponds as well as dams, drains and storm water basins.                                                                                                                                                                                                                                                                                                                       | <b>Low</b><br>Suitable habitat was present within dams, drainage lines and wetlands along the REZ route. Few recent records. Most recent records are around Ravensworth |
| <i>Litoria booroolongensis</i><br>Booroolong Frog       | E1                | E                    | No  | Restricted to NSW and north-eastern Victoria, predominantly along the western-flowing streams of the Great Dividing Range. Lives along permanent streams with some fringing vegetation cover such as ferns, sedges, or grasses. Adults occur on or near cobble banks and other rock structures within stream margins.                                                                                             | <b>Unlikely</b><br>No suitable habitat was present. Outside known distribution.                                                                                         |
| <i>Litoria brevipalmata</i><br>Green-thighed Frog       | V                 |                      | No  | Occurs in isolated localities along the coast and ranges from just north of Wollongong to south-east Queensland. Occurs in a range of habitats from rainforest and moist eucalypt forest to dry eucalypt forest and heath, typically in areas where surface water gathers after rain. It prefers wetter forests in the south of its range but extends into drier forests in northern NSW and southern Queensland. | <b>Low</b><br>Marginal habitat confined to aquatic areas in the far east of the REZ route.                                                                              |
| <i>Litoria littlejohni</i><br>Littlejohn's Tree Frog    | E1                | E                    | No  | Habitats include wet and dry sclerophyll forest, coastal woodland and heath. Associated characteristics include rocky streams and sandstone outcrops, semi-permanent dams and slow flowing streams. The water quality required for breeding is usually tannic (pH 6.2) and contains detritus which is used as anchors for egg clusters.                                                                           | <b>Low</b><br>Marginal habitat confined to aquatic areas in the far east of the REZ route.                                                                              |
| <i>Mixophyes balbus</i><br>Stuttering Frog              | E1                | V                    | Yes | Occurs in wet forest regions of south-eastern Queensland, Eastern NSW and Victoria. In late spring, eggs are deposited among leaf litter on the banks of streams and subsequently are washed into the water during heavy rain.                                                                                                                                                                                    | <b>Unlikely</b><br>No suitable habitat was present. Outside known distribution.                                                                                         |
| <i>Mixophyes iteratus</i><br>Giant Barred Frog          | V                 | V                    | No  | Distributed from Doongul Creek, Wongi State Forest, near Maryborough in south-eastern Queensland (Hines 2003), south to Warrimoo in the Blue Mountains, New South Wales. Occurs in rainforests and wet sclerophyll forests in upper to lower catchment areas (Ingram & McDonald 1993).                                                                                                                            | <b>Unlikely</b><br>No suitable habitat was present. Outside known distribution.                                                                                         |
| <i>Pseudophryne australis</i><br>Red-crowned Toadlet    | V                 |                      | No  | Distribution within sandstone of the Sydney Basin. This frog species may utilise areas below the first cliff line for breeding and the adjacent ridge tops and escarpments for foraging and refuge. Usually found along "feeder creeks" which feed run-off water from ridges into perennial creeks.                                                                                                               | <b>Unlikely</b><br>No suitable habitat was present. Outside known distribution.                                                                                         |
| <b>FAUNA - REPTILES</b>                                 |                   |                      |     |                                                                                                                                                                                                                                                                                                                                                                                                                   |                                                                                                                                                                         |
| <i>Aprasia parapulchella</i><br>Pink-tailed Worm-lizard | V                 | V                    | No  | Is distributed along the western foothills of the Great Dividing Range between Bendigo in Victoria and Gunnedah in northern New South Wales. Generally, occupies sites with a grassy ground layer particularly those dominated by Kangaroo Grass with little or no                                                                                                                                                | <b>Low</b><br>Marginal habitat was present.                                                                                                                             |

| SPECIES                                                | STATUS            |                      |     | HABITAT DESCRIPTION AND LOCALLY KNOWN POPULATIONS                                                                                                                                                                                                                                                                                  | LIKELIHOOD OF OCCURRENCE<br>WITHIN THE REZ SITE                                                                                                                  |
|--------------------------------------------------------|-------------------|----------------------|-----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                        | BC<br>Act<br>2016 | EPB<br>C Act<br>1999 | SAI |                                                                                                                                                                                                                                                                                                                                    |                                                                                                                                                                  |
|                                                        |                   |                      |     | leaf litter, and relatively low tree and shrub cover. Sites are typically well-drained, with rocky outcrops or scattered, partially buried rocks.                                                                                                                                                                                  |                                                                                                                                                                  |
| <i>Delma vescolineata</i><br>Hunter Valley Delma       | E1                | E                    | No  | Is known almost entirely from a 25 km wide corridor in the Hunter Valley NSW, between Maitland and Muswellbrook. The sites where the species has been detected include rehabilitated mine sites and pastoral land used for cattle grazing.                                                                                         | <b>High</b><br>Known records within proximity to the Muswellbrook STS site and easement to the NE. Less likely to be recorded in the eastern portion of the REZ. |
| <i>Varanus rosenbergi</i><br>Rosenberg's Goanna        | V                 |                      | No  | Occurs on the Sydney Sandstone in Wollemi National Park to the north-west of Sydney, in the Goulburn and ACT regions and near Cooma in the south. Found in heath, open forest and woodland. Associated with termites, the mounds of which this species nests in; termite mounds are a critical habitat component.                  | <b>Low</b><br>Suitable habitat was present. No known local records.                                                                                              |
| <i>Hoplocephalus bungaroides</i><br>Broad-headed Snake | E1                | E                    | Yes | Found under large slabs of rock or crevices on sandstone outcrops. Is largely confined to Triassic and Permian sandstones, including the Hawkesbury, Narrabeen and Shoalhaven groups, within the coast and ranges in an area within approximately 250 km of Sydney.                                                                | <b>Unlikely</b><br>No suitable habitat was present. Outside known distribution.                                                                                  |
| <b>FAUNA - BIRDS</b>                                   |                   |                      |     |                                                                                                                                                                                                                                                                                                                                    |                                                                                                                                                                  |
| <i>Calidris acuminata</i><br>Sharp-tailed Sandpiper    |                   | V & M                | No  | Widespread in both inland and coastal locations and in both freshwater and saline habitats.                                                                                                                                                                                                                                        | <b>Low</b><br>Suitable habitat was present.                                                                                                                      |
| <i>Calidris ferruginea</i><br>Curlew Sandpiper         | E4A               | CE                   | Yes | Tidal mudflats; saltmarsh; fresh, brackish or saline wetlands; sewage ponds.                                                                                                                                                                                                                                                       | <b>Unlikely</b><br>No suitable habitat was present.                                                                                                              |
| <i>Tringa nebularia</i><br>Common Greenshank           | E1                | E & M                | No  | Inhabits a wide variety of inland permanent and temporary wetlands and sheltered coastal habitats of varying salinity.                                                                                                                                                                                                             | <b>Low</b><br>Suitable habitat was present.                                                                                                                      |
| <i>Numenius madagascariensis</i><br>Eastern Curlew     | E4A               | CE<br>M              | No  | Estuaries, tidal mudflats, sandspits, saltmarshes, mangroves; occasionally fresh or brackish lakes.                                                                                                                                                                                                                                | <b>Unlikely</b><br>No suitable habitat was present.                                                                                                              |
| <i>Charadrius leschenaultii</i><br>Greater Sand-plover | V                 |                      | No  | In NSW, the species has been recorded between the northern rivers and the Illawarra, with most records coming from the Clarence and Richmond estuaries. Almost entirely restricted to coastal areas in NSW, occurring mainly on sheltered sandy, shelly or muddy beaches or estuaries with large intertidal mudflats or sandbanks. | <b>Unlikely</b><br>No suitable habitat was present.                                                                                                              |
| <i>Gallinago hardwickii</i><br>Latham's Snipe          | V                 | V & M                | No  | Utilises a variety of habitat, such as soft wet ground or shallow water with tussock and other green and dead vegetation, and scrub or open wetland from sea-level to alpine bogs.                                                                                                                                                 | <b>Moderate</b><br>Suitable areas of aquatic habitat were present along the REZ route.                                                                           |
| <i>Rostratula australis</i>                            | E1                | E                    | No  | Margins of swamps and streams, chiefly those covered with low and stunted vegetation.                                                                                                                                                                                                                                              | <b>Low</b>                                                                                                                                                       |



| SPECIES                                                 | STATUS            |                      |     | HABITAT DESCRIPTION AND LOCALLY KNOWN POPULATIONS                                                                                                                                                                                                                                                                                                                                                                          | LIKELIHOOD OF OCCURRENCE<br>WITHIN THE REZ SITE                                              |
|---------------------------------------------------------|-------------------|----------------------|-----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------|
|                                                         | BC<br>Act<br>2016 | EPB<br>C Act<br>1999 | SAI |                                                                                                                                                                                                                                                                                                                                                                                                                            |                                                                                              |
| Australian Painted snipe                                |                   |                      |     |                                                                                                                                                                                                                                                                                                                                                                                                                            | Suitable areas of aquatic habitat were present along the REZ route.                          |
| <i>Botaurus poiciloptilus</i><br>Australasian Bittern   | E1                | E                    | No  | The Australasian Bittern lives alone or in loose groups and favours permanent fresh waters dominated by sedges, rushes, reeds or cutting grasses (e.g. Phragmites, Scirpus, Eleocharis, Juncus, Typha, Baumea and Gahnia) and feeds on insects, small fish, eels, frogs and other aquatic life, sometimes in rice fields.                                                                                                  | <b>Low</b><br>Suitable areas of aquatic habitat were present along the REZ route.            |
| <i>Ixobrychus flavicollis</i><br>Black Bittern          | V                 |                      | No  | Inhabits both terrestrial and estuarine wetlands, generally in areas of permanent water and dense vegetation. Where permanent water is present, the species may occur in flooded grassland, forest, woodland, rainforest and mangroves.                                                                                                                                                                                    | <b>Low</b><br>Suitable areas of aquatic habitat were present along the REZ route.            |
| <i>Ephippiorhynchus asiaticus</i><br>Black-necked Stork | E1                |                      | No  | Widespread in coastal and subcoastal northern and eastern Australia, as far south as central NSW. Breeding has been recorded as far south as Tomago NSW.                                                                                                                                                                                                                                                                   | <b>Low</b><br>Suitable larger areas of aquatic habitat were present along the REZ route.     |
| <i>Anseranas semipalmata</i><br>Magpie Goose            | V                 |                      | No  | Relatively common in the Australian northern tropics. Records in central and northern NSW. Vagrants can follow food sources to south-eastern NSW. Mainly found in shallow wetlands (less than 1 m deep) with dense growth of rushes or sedges.                                                                                                                                                                             | <b>Low</b><br>Suitable areas of aquatic habitat were present along the REZ route.            |
| <i>Oxyura australis</i><br>Blue-billed Duck             | V                 |                      | No  | Endemic to south-eastern and south-western Australia. It is widespread in NSW. Most common in the southern Murray-Darling Basin area. Prefers deep water in large permanent wetlands and swamps with dense aquatic vegetation.                                                                                                                                                                                             | <b>Low</b><br>Suitable areas of aquatic habitat were present along the REZ route.            |
| <i>Stictonetta naevosa</i><br>Freckled Duck             | V                 |                      | No  | Found primarily in south-eastern and south-western Australia, occurring as a vagrant elsewhere. It breeds in large temporary swamps created by floods in the Bulloo and Lake Eyre basins and the Murray-Darling system. The species may also occur as far as coastal NSW and Victoria during extensive inland droughts.                                                                                                    | <b>Low</b><br>Suitable areas of aquatic habitat were present along the REZ route.            |
| <i>Irediparra gallinacea</i><br>Comb-crested Jacana     | V                 |                      | No  | Occurs in northern and eastern Australia, mainly in coastal and subcoastal regions, from the north-eastern Kimberley Division of Western Australia to Cape York Peninsula then south along the east coast to the Hunter region of NSW. Inhabits permanent freshwater wetlands, either still or slow-flowing, with a good surface cover of floating vegetation, especially waterlilies, or fringing and aquatic vegetation. | <b>Unlikely</b><br>No suitable habitat was present.                                          |
| <i>Burhinus grallarius</i><br>Bush Stone Curlew         | E1                |                      | No  | Found throughout Austroriparian Australia except for the central southern coast and inland, the far south-east corner, and Tasmania. Inhabits open forests and woodlands with a sparse grassy ground layer and fallen timber.                                                                                                                                                                                              | <b>Low</b><br>Suitable areas of habitat were present along the REZ route.                    |
| <i>Turnix maculosus</i><br>Red-backed Button-quail      | V                 |                      | No  | Recorded only infrequently in NSW, with most records from the North Coast Bioregion. In NSW, said to occur in grasslands, heath and crops. Said to prefer sites close to water, especially when breeding.                                                                                                                                                                                                                  | <b>Low</b><br>Suitable areas of habitat were present along the REZ route. Few local records. |
| <i>Sterna albifrons</i><br>Little Tern                  | E1                |                      | No  | Migrating from eastern Asia, the Little Tern is found on the north, east and south-east Australian coasts. Exclusively coastal, nests in small, scattered colonies in low dunes                                                                                                                                                                                                                                            | <b>Unlikely</b><br>No suitable habitat was present.                                          |

| SPECIES                                                  | STATUS            |                      |     | HABITAT DESCRIPTION AND LOCALLY KNOWN POPULATIONS                                                                                                                                                                                                            | LIKELIHOOD OF OCCURRENCE<br>WITHIN THE REZ SITE                               |
|----------------------------------------------------------|-------------------|----------------------|-----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------|
|                                                          | BC<br>Act<br>2016 | EPB<br>C Act<br>1999 | SAI |                                                                                                                                                                                                                                                              |                                                                               |
|                                                          |                   |                      |     | or on sandy beaches just above high tide mark near estuary mouths or adjacent to coastal lakes and islands.                                                                                                                                                  |                                                                               |
| <i>Ptilinopus magnificus</i><br>Wompoo Fruit-Dove        | V                 |                      | No  | Occurs along the coast and coastal ranges from the Hunter River in NSW to Cape York Peninsula. It is rare south of Coffs Harbour. Found in, or near rainforest, low elevation moist eucalypt forest and brush box forests.                                   | <b>Unlikely</b><br>No suitable habitat was present.                           |
| <i>Ptilinopus regina</i><br>Rose-crowned Fruit-Dove      | V                 |                      | No  | Coast and ranges of eastern NSW and Queensland, from Newcastle to Cape York. Occur mainly in sub-tropical and dry rainforest and occasionally in moist eucalypt forest and swamp forest, where fruit is plentiful.                                           | <b>Unlikely</b><br>No suitable habitat was present.                           |
| <i>Calyptorhynchus lathamii</i><br>Glossy Black-Cockatoo | V                 | V                    | No  | Lowland coastal forests, dense mountain forests, semi-arid woodland and trees bordering watercourses, with (Allo)Casuarina trees for foraging.                                                                                                               | <b>Low</b><br>Suitable habitat was present.                                   |
| <i>Callocephalon fimbriatum</i><br>Gang Gang Cockatoo    | E1                | E                    | No  | Tall montane forests and woodlands in mature wet sclerophyll forests. Requires hollows in which to breed between October and January.                                                                                                                        | <b>Low</b><br>Suitable habitat was present.                                   |
| <i>Lathamus discolor</i><br>Swift Parrot                 | E1                | CE                   | Yes | Open Forest to Woodland, also street trees and in parks and gardens, winter flowering eucalypts for feeding. This species nests in Tasmania during the summer months.                                                                                        | <b>Moderate</b><br>Seasonal foraging habitat was present.                     |
| <i>Neophema chrysostoma</i><br>Blue-winged Parrot        | V                 | V                    | No  | Found in western NSW. They favour grasslands and grassy woodlands. They are often found near wetlands both near the coast and in semi-arid zones. Blue-winged Parrots can also be seen in altered environments such as airfields, golf-courses and paddocks. | <b>Low</b><br>Suitable habitat areas were present.                            |
| <i>Neophema pulchella</i><br>Turquoise Parrot            | V                 |                      | No  | Lives on the edges of Eucalypt woodland adjoining clearings and on timbered ridges and creeks in farmland. It has also been recorded utilising roadside verges and orchards. Nests in small hollow branches of Eucalypts.                                    | <b>Low</b><br>Suitable habitat areas were present.                            |
| <i>Polytelis swainsonii</i><br>Superb Parrot             | V                 | V                    | No  | Found throughout eastern inland NSW. They inhabit Box-Gum, Box-Cypress-pine and Boree woodlands and River Red Gum Forest.                                                                                                                                    | <b>Low</b><br>Marginal habitat areas were present in the far west of the REZ. |
| <i>Glossopsitta pusilla</i><br>Little Lorikeet           | V                 |                      | No  | Tall Open Forests, woodlands, orchards, parks and street trees.                                                                                                                                                                                              | <b>High</b><br>Suitable habitat areas were present along the REZ route.       |
| <i>Aphelocephala leucopsis</i>                           | V                 | V                    | No  | Open woodlands and shrublands with understorey of grasses or shrubs, or both.                                                                                                                                                                                | <b>Low</b>                                                                    |



| SPECIES                                                                     | STATUS            |                      |     | HABITAT DESCRIPTION AND LOCALLY KNOWN POPULATIONS                                                                                                                                                                                                                                                                                                                                                                                                                                                                        | LIKELIHOOD OF OCCURRENCE<br>WITHIN THE REZ SITE                                                                                                    |
|-----------------------------------------------------------------------------|-------------------|----------------------|-----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                             | BC<br>Act<br>2016 | EPB<br>C Act<br>1999 | SAI |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |                                                                                                                                                    |
| Southern Whiteface                                                          |                   |                      |     |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          | Suitable habitat areas were present along the REZ route.                                                                                           |
| <i>Hirundapus caudacutus</i><br>White-throated Needletail                   | V                 | V & M                | No  | Inhabits the airspace above forests, woodlands, farmlands, plains, lakes, coasts and towns.                                                                                                                                                                                                                                                                                                                                                                                                                              | <b>Moderate</b><br>Due to the non-specific habitat requirements of this species habitat was considered to be present.                              |
| <i>Artamus cyanopterus cyanopterus</i><br>Dusky Woodswallow                 | V                 |                      | No  | The Dusky Woodswallow is found in open forests and woodlands and may be seen along roadsides and on golf courses.                                                                                                                                                                                                                                                                                                                                                                                                        | <b>Moderate-High</b><br>Suitable habitat areas were present along the REZ route.                                                                   |
| <i>Pycnoptilus floccosus</i><br>Pilotbird                                   | V                 | V                    |     | Found in wet forested areas and heathland in eastern Victoria and south-eastern New South Wales                                                                                                                                                                                                                                                                                                                                                                                                                          | <b>Unlikely</b><br>No suitable habitat was present.                                                                                                |
| <i>Epthianura albifrons</i><br>White-fronted Chat                           | V                 |                      | No  | In NSW, it occurs mostly in the southern half of the state, in damp open habitats along the coast, and near waterways in the western part of the state. Along the coastline, it is found predominantly in saltmarsh vegetation but also in open grasslands and sometimes in low shrubs bordering wetland areas. Gregarious species usually found foraging on bare or grassy ground in wetland areas, singly or in pairs. They are insectivorous, feeding mainly on flies and beetles caught from or close to the ground. | <b>Unlikely</b><br>No suitable habitat was present.                                                                                                |
| <i>Melanodryas cucullata cucullata</i><br>Hooded Robin (south-eastern form) | E1                | E                    | No  | Eucalypt woodlands, Acacia scrublands, Banksia dominated coastal scrubs and open forests.                                                                                                                                                                                                                                                                                                                                                                                                                                | <b>Low-Moderate</b><br>Suitable habitat areas were present along the REZ route.                                                                    |
| <i>Petroica boodang</i><br>Scarlet Robin                                    | V                 |                      | No  | Primarily a resident in forests and woodlands, but some adults and young birds disperse to more open habitats after breeding. This species lives in dry eucalypt forests and woodlands. The understorey is usually open and grassy with few scattered shrubs. Habitat usually contains abundant logs and fallen timber and these are important components of its habitat.                                                                                                                                                | <b>Low-Moderate</b><br>Suitable habitat areas were present along the REZ route.                                                                    |
| <i>Petroica phoenicea</i><br>Flame Robin                                    | V                 |                      | No  | Breeds in upland tall moist eucalypt forests and woodlands, often on ridges and slopes. Endemic to south-eastern Australia, and ranges from near the Queensland border to southeast South Australia and also in Tasmania.                                                                                                                                                                                                                                                                                                | <b>Low-Moderate</b><br>Suitable habitat areas were present along the REZ route.                                                                    |
| <i>Climacteris picumnus victoriae</i><br>Brown Treecreeper                  | V                 | V                    | No  | This species is a medium sized insectivorous bird that occupies Eucalypt woodlands, particularly open woodlands lacking a dense understorey, River Red Gums on watercourses and around lakeshores. It is sedentary and nests in tree hollows within permanent territories.                                                                                                                                                                                                                                               | <b>Known</b><br>Recorded within proximity to easement NE of Muswellbrook STS. Suitable habitat areas were also present along the entire REZ route. |
| <i>Stagonopleura guttata</i>                                                | V                 | V                    | No  | Inhabits areas with a grassy, shrubby understorey including Eucalypt woodlands, forests, Acacia scrubs and mallee.                                                                                                                                                                                                                                                                                                                                                                                                       | <b>Moderate</b>                                                                                                                                    |

| SPECIES                                                                         | STATUS            |                      |     | HABITAT DESCRIPTION AND LOCALLY KNOWN POPULATIONS                                                                                                                                                                                                                                                                                                                    | LIKELIHOOD OF OCCURRENCE<br>WITHIN THE REZ SITE                                  |
|---------------------------------------------------------------------------------|-------------------|----------------------|-----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------|
|                                                                                 | BC<br>Act<br>2016 | EPB<br>C Act<br>1999 | SAI |                                                                                                                                                                                                                                                                                                                                                                      |                                                                                  |
| Diamond Firetail                                                                |                   |                      |     |                                                                                                                                                                                                                                                                                                                                                                      | Suitable habitat areas were present along the REZ route.                         |
| <i>Pomatostomus temporalis</i> subsp. <i>temporalis</i><br>Grey-crowned Babbler | V                 |                      | No  | Open forest, woodland, scrubland, farmland and outer suburbs. Prefers woodlands with regenerating trees, tall shrubs, and an intact ground cover of grass and forbs.                                                                                                                                                                                                 | <b>Known</b><br>Recorded along a number of locations along the REZ route.        |
| <i>Chthonicola sagittata</i><br>Speckled Warbler                                | V                 |                      | No  | Speckled Warblers live in a wide range of eucalypt-dominated vegetation that has a grassy understorey, often on rocky ridges or in gullies. It builds a domed nest of grass, bark shreds and moss, lined with fur on the ground.                                                                                                                                     | <b>Moderate-High</b><br>Suitable habitat areas were present along the REZ route. |
| <i>Anthochaera phrygia</i><br>Regent Honeyeater                                 | E4A               | CE                   | Yes | Temperate woodlands and open forest, including forest edges, preferring to forage on large-flowered Eucalypts.                                                                                                                                                                                                                                                       | <b>Low</b><br>Suitable habitat areas were present along the REZ route.           |
| <i>Certhionyx variegatus</i><br>Pied Honeyeater                                 | V                 |                      |     | Occasionally occurs on the slopes and plains and the Hunter Valley, typically during periods of drought. Inhabits wattle shrub mallee, spinifex and eucalypt woodlands, usually when shrubs are flowering.                                                                                                                                                           | <b>Low</b><br>Suitable habitat areas were present along the REZ route.           |
| <i>Grantiella picta</i><br>Painted Honeyeater                                   | V                 | V                    | No  | Nomadic, within a range of generally drier forested areas with mistletoes.                                                                                                                                                                                                                                                                                           | <b>Low</b><br>Suitable habitat areas were present along the REZ route.           |
| <i>Melithreptus gularis</i><br>Black-chinned Honeyeater (eastern subspecies)    | V                 |                      | No  | Usually found on the western side of the Great Dividing Range in dry sclerophyll forests and woodlands containing box-ironbark associations and River Red Gum. In the Hunter Valley this species is known to utilise drier coastal woodlands. Usually found in open woodlands.                                                                                       | <b>Low-Moderate</b><br>Suitable habitat areas were present along the REZ route.  |
| <i>Daphoenositta chrysoptera</i><br>Varied Sittella                             | V                 |                      | No  | Open eucalypt woodland/forest, mallee, inland acacia, coastal tea-tree scrubs, golf courses, orchards and parks.                                                                                                                                                                                                                                                     | <b>Moderate</b><br>Suitable habitat areas were present along the REZ route.      |
| <i>Circus assimilis</i><br>Spotted Harrier                                      | V                 |                      | No  | Occurs throughout the Australian mainland, except in densely forested or wooded habitats of the coast, escarpment and ranges, and rarely in Tasmania. Found in grassy open woodland including Acacia and mallee remnants, inland riparian woodland, grassland and shrub steppe. It is found most commonly in native grassland, but also occurs in agricultural land. | <b>Low-Moderate</b><br>Suitable habitat areas were present along the REZ route.  |
| <i>Pandion cristatus</i><br>Eastern Osprey                                      | V                 |                      | No  | Found right around the Australian coastline. Favour coastal areas, especially the mouths of large rivers, lagoons and lakes. Feeds on fish over open waters.                                                                                                                                                                                                         | <b>Unlikely</b><br>No suitable habitat was present.                              |



| SPECIES                                                   | STATUS            |                      |      | HABITAT DESCRIPTION AND LOCALLY KNOWN POPULATIONS                                                                                                                                                                                                                                                                                                                                                                                 | LIKELIHOOD OF OCCURRENCE<br>WITHIN THE REZ SITE                                 |
|-----------------------------------------------------------|-------------------|----------------------|------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------|
|                                                           | BC<br>Act<br>2016 | EPB<br>C Act<br>1999 | SAIL |                                                                                                                                                                                                                                                                                                                                                                                                                                   |                                                                                 |
| <i>Lophoictinia isura</i><br>Square-tailed Kite           | V                 |                      | No   | Inhabits open forests and woodlands, particularly those on fertile soils with abundant passerines.                                                                                                                                                                                                                                                                                                                                | <b>Low-Moderate</b><br>Suitable habitat areas were present along the REZ route. |
| <i>Erythrotriorchis radiatus</i><br>Red Goshawk           | E1                | E                    | Yes  | The species is very rare in NSW, extending south to about 30°S, with most records north of this, in the Clarence River Catchment, and a few around the lower Richmond and Tweed Rivers. Formerly, it was at least occasionally reported as far south as Port Stephens. In NSW, preferred habitats include mixed subtropical rainforest, Melaleuca swamp forest and riparian Eucalyptus Forest of coastal rivers.                  | <b>Unlikely</b><br>No suitable habitat was present.                             |
| <i>Hieraaetus morphnoides</i><br>Little Eagle             | V                 |                      | No   | Is found throughout the Australian mainland excepting the most densely forested parts of the Dividing Range escarpment. It occurs as a single population throughout NSW. Occupies open eucalypt forest, woodland or open woodland. Sheoak or acacia woodlands and riparian woodlands of interior NSW are also used.                                                                                                               | <b>Moderate</b><br>Suitable habitat areas were present along the REZ route.     |
| <i>Haliaeetus leucogaster</i><br>White-bellied Sea-Eagle  | V                 | M                    | No   | Occupies habitat characterised by the presence of large areas of open water and feeds opportunistically on a variety of fish, birds, reptiles, mammals and crustaceans. The nests are built in a variety of sites including tall trees, bushes, mangroves, cliffs, rocky outcrops, caves, crevices, on the ground or even in artificial structures.                                                                               | <b>Low-Moderate</b><br>Suitable habitat areas were present along the REZ route. |
| <i>Hamirostra melanosternon</i><br>Black-breasted Buzzard | V                 |                      | No   | Found sparsely in areas of less than 500mm rainfall, from north-western NSW and north-eastern South Australia to the east coast at about Rockhampton. Lives in a range of inland habitats, especially along timbered watercourses which is the preferred breeding habitat. Hunts over grasslands and sparsely timbered woodlands.                                                                                                 | <b>Low-Moderate</b><br>Suitable habitat areas were present along the REZ route. |
| <i>Falco hypoleucos</i><br>Grey Falcon                    | V                 | V                    | No   | Sparsely distributed in NSW, chiefly throughout the Murray-Darling Basin, with the occasional vagrant east of the Great Dividing Range. Generally restricted to shrubland, grassland and wooded watercourses of arid and semi-arid regions, although it is occasionally found in open woodlands near the coast.                                                                                                                   | <b>Low</b><br>Suitable habitat areas were present along the REZ route.          |
| <i>Falco subniger</i><br>Black Falcon                     | V                 |                      | No   | Widely, but sparsely, distributed in New South Wales, mostly occurring in inland regions.                                                                                                                                                                                                                                                                                                                                         | <b>Low-Moderate</b><br>Suitable habitat areas were present along the REZ route. |
| <i>Ninox connivens</i><br>Barking Owl                     | V                 |                      | No   | Is found in forest and woodland, encountered most commonly in savanna and paperbark woodlands. It sometimes roosts in rainforests, but it requires the more open country for hunting and hollow Eucalypts for breeding.                                                                                                                                                                                                           | <b>Low-Moderate</b><br>Suitable habitat areas were present along the REZ route. |
| <i>Ninox strenua</i><br>Powerful Owl                      | V                 |                      | No   | Inhabits a wide range of vegetation types from wet Eucalypt forests with a Rainforest understorey to Dry Open Forests and Woodlands. The species has been recorded utilising disturbed habitats such as exotic pine plantations and large trees in parks and gardens. Powerful Owls nest in a slight depression in the wood-mould on the base of a cavity in a large old tree, sometimes in excess of 25 metres above the ground. | <b>Low-Moderate</b><br>Suitable habitat areas were present along the REZ route. |
| <i>Tyto longimembris</i><br>Eastern Grass Owl             | V                 |                      | No   | Recorded occasionally in all mainland states of Australia but are most common in northern and north-eastern Australia. Found in areas of tall grass, including grass                                                                                                                                                                                                                                                              | <b>Unlikely</b><br>No suitable habitat was present.                             |

| SPECIES                                                                     | STATUS            |                      |     | HABITAT DESCRIPTION AND LOCALLY KNOWN POPULATIONS                                                                                                                                                                                                                                                                                                                                                     | LIKELIHOOD OF OCCURRENCE<br>WITHIN THE REZ SITE                                                            |
|-----------------------------------------------------------------------------|-------------------|----------------------|-----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------|
|                                                                             | BC<br>Act<br>2016 | EPB<br>C Act<br>1999 | SAI |                                                                                                                                                                                                                                                                                                                                                                                                       |                                                                                                            |
|                                                                             |                   |                      |     | tussocks, in swampy areas, grassy plains, swampy heath, and in cane grass or sedges on flood plains.                                                                                                                                                                                                                                                                                                  |                                                                                                            |
| <i>Tyto novaehollandiae</i><br>Masked Owl                                   | V                 |                      | No  | A range of wooded habitats that contain mature trees with large hollows for roosting and nesting, and more open areas for hunting.                                                                                                                                                                                                                                                                    | <b>Low</b><br>Suitable habitat areas were present along the REZ route.                                     |
| <i>Tyto tenebricosa</i><br>Sooty Owl                                        | V                 |                      | Yes | Prefers dense dimly lit forests, inhabiting pockets of rainforest and wet sclerophyll forest mainly in mountainous areas, often in south-east facing gullies.                                                                                                                                                                                                                                         | <b>Unlikely</b><br>No suitable habitat was present.                                                        |
| <b>FAUNA – MAMMALS</b>                                                      |                   |                      |     |                                                                                                                                                                                                                                                                                                                                                                                                       |                                                                                                            |
| <i>Dasyurus maculatus</i><br>ssp. <i>maculatus</i><br>Spotted-tailed Quoll  | V                 | E                    | No  | Inhabits sclerophyll forests, rainforests and coastal woodlands. Nests are made in rock caves and hollow logs or trees, and basking sites are usually found nearby.                                                                                                                                                                                                                                   | <b>Unlikely</b><br>No suitable habitat was present.                                                        |
| <i>Phascogale tapoatafa</i><br>Brush-tailed<br>Phascogale                   | V                 |                      | No  | Prefer dry sclerophyll open forest with sparse groundcover of herbs, grasses, shrubs or leaf litter.                                                                                                                                                                                                                                                                                                  | <b>Low</b><br>Suitable habitat areas containing open forest and woodland were present along the REZ route. |
| <i>Phascolarctos cinereus</i><br>Koala                                      | E1                | E                    | No  | Coastal woodland and open forest containing suitable food trees.                                                                                                                                                                                                                                                                                                                                      | <b>Low</b><br>Areas containing habitat were present along the REZ route.                                   |
| <i>Petrogale penicillata</i><br>Brush-tailed Rock-wallaby                   | E1                | V                    | Yes | Found in steep rocky sites in sclerophyll forests with a grassy understorey.                                                                                                                                                                                                                                                                                                                          | <b>Unlikely</b><br>No suitable habitat was present.                                                        |
| <i>Notamacropus parma</i><br>Parma Wallaby                                  | V                 | V                    | No  | Range is now confined to the coast and ranges of central and northern NSW from the Gosford district to south of the Bruxner Highway between Tenterfield and Casino. Preferred habitat is moist eucalypt forest with thick, shrubby understorey, often with nearby grassy areas, rainforest margins and occasionally drier eucalypt forest.                                                            | <b>Unlikely</b><br>No suitable habitat was likely present.                                                 |
| <i>Potorous tridactylus</i><br>sp. <i>tridactylus</i><br>Long-nosed Potoroo | V                 | V                    | No  | This species is known from a variety of habitats, including Rainforest, Open Forests and Woodlands with dense groundcover, and dense, wet coastal heathlands. Soft (often sandy) substrates are preferred by this species.                                                                                                                                                                            | <b>Low</b><br>A small number of areas containing marginal were present along the REZ route.                |
| <i>Cercartetus nanus</i><br>Eastern Pygmy-possum                            | V                 |                      | No  | In NSW it extends from the coast inland as far as the Pilliga, Dubbo, Parkes and Wagga Wagga on the western slopes. Found in a broad range of habitats from rainforest through sclerophyll (including Box-Ironbark) forest and woodland to heath, but in most areas woodlands and heath appear to be preferred, except in north-eastern NSW where they are most frequently encountered in rainforest. | <b>Low</b><br>A small number of areas containing marginal were present along the REZ route.                |
| <i>Petaurus australis</i><br>Yellow-bellied Glider                          | V                 | V                    | No  | Occurs in tall mature eucalypt forest generally in areas with high rainfall and nutrient rich soils. Forest type preferences vary with latitude and elevation; mixed coastal forests to                                                                                                                                                                                                               | <b>Low</b>                                                                                                 |



| SPECIES                                                           | STATUS            |                      |     | HABITAT DESCRIPTION AND LOCALLY KNOWN POPULATIONS                                                                                                                                                                                                                                                                                                                                                                                                                           | LIKELIHOOD OF OCCURRENCE<br>WITHIN THE REZ SITE                                                                                                                                                                                        |
|-------------------------------------------------------------------|-------------------|----------------------|-----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                   | BC<br>Act<br>2016 | EPB<br>C Act<br>1999 | SAI |                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |                                                                                                                                                                                                                                        |
|                                                                   |                   |                      |     | dry escarpment forests in the north; moist coastal gullies and creek flats to tall montane forests in the south. Is found along the eastern coast to the western slopes of the Great Dividing Range, from southern Queensland to Victoria.                                                                                                                                                                                                                                  | A small number of areas containing marginal were present along the REZ route.                                                                                                                                                          |
| <i>Petaurus norfolcensis</i><br>Squirrel Glider                   | V                 |                      | No  | Dry sclerophyll forests and woodlands with exudates for foraging and hollows for nesting.                                                                                                                                                                                                                                                                                                                                                                                   | <b>High</b><br>Suitable habitat areas containing open forest and woodland were present along the REZ route. Has been recorded within a number of locations within proximity to the REZ according to BioNet records (NSW DCCEEW, 2024). |
| <i>Petauroides volans</i><br>Greater Glider                       | E1                | E                    | No  | Eucalypt-dominated low open forests on the coast to tall forests in the ranges and low woodland west of Great Dividing Range. Not found within rainforests.                                                                                                                                                                                                                                                                                                                 | <b>Low</b><br>A small number of areas containing marginal habitat were present along the REZ route. No known records.                                                                                                                  |
| <i>Pseudomys novaehollandiae</i><br>New Holland Mouse             |                   | V                    | No  | Known to inhabit open heathlands, open woodlands with a heathland understorey and vegetated sand dunes.                                                                                                                                                                                                                                                                                                                                                                     | <b>Low</b><br>A small number of areas containing suitable habitat were present along the REZ route.                                                                                                                                    |
| <i>Pseudomys oralis</i><br>Hastings River Mouse                   | E1                | E                    | No  | A patchy distribution spanning the Great Dividing Range from the Hunter Valley, south of Mt Royal, north to the Bunya Mountains near Kingaroy in south-east Queensland, at elevations between 300 m and 1100 m. Found in a variety of dry open forest types with dense, low ground cover and a diverse mixture of ferns, grass, sedges and herbs. Access to seepage zones, creeks and gullies is important, as is permanent shelter such as rocky outcrops and fallen logs. | <b>Unlikely</b><br>No suitable habitat was likely present.                                                                                                                                                                             |
| <i>Pteropus poliocephalus</i><br>Grey-headed Flying-Fox           | V                 | V                    | No  | Wet and Dry Sclerophyll Forests, Rainforest, Mangroves and Paperbark swamps and Banksia Woodlands.                                                                                                                                                                                                                                                                                                                                                                          | <b>High</b><br>Suitable seasonal foraging habitat was present along the entire length of the REZ.                                                                                                                                      |
| <i>Saccolaimus flaviventris</i><br>Yellow-bellied Sheath-tail-bat | V                 |                      | No  | Has been reported from a wide variety of habitats. Roosts in tree hollows, animal burrows, dry clay cracks, under rock slabs and in abandoned Sugar Glider nests.                                                                                                                                                                                                                                                                                                           | <b>Moderate</b><br>Suitable habitat areas containing hunting and roosting habitat were present along the REZ route.                                                                                                                    |
| <i>Micronomus norfolkensis</i><br>Eastern Freetail-bat            | V                 |                      | No  | Appears to live in sclerophyll forests and woodland. Roosts in tree hollows or under loose bark.                                                                                                                                                                                                                                                                                                                                                                            | <b>Moderate - High</b><br>Suitable habitat areas containing hunting and roosting habitat were present along the REZ route.                                                                                                             |

| SPECIES                                                        | STATUS            |                      |     | HABITAT DESCRIPTION AND LOCALLY KNOWN POPULATIONS                                                                                                                                                                                                                                                                                                                                                                        | LIKELIHOOD OF OCCURRENCE<br>WITHIN THE REZ SITE                                                                                                          |
|----------------------------------------------------------------|-------------------|----------------------|-----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                | BC<br>Act<br>2016 | EPB<br>C Act<br>1999 | SAI |                                                                                                                                                                                                                                                                                                                                                                                                                          |                                                                                                                                                          |
| <i>Falsistrellus tasmaniensis</i><br>Eastern False Pipistrelle | V                 |                      | No  | Inhabits sclerophyll forests and has been observed roosting in holes and hollow trunks of Eucalypts.                                                                                                                                                                                                                                                                                                                     | <b>Moderate</b><br>Suitable habitat areas containing hunting and roosting habitat were present along the REZ route.                                      |
| <i>Miniopterus australis</i><br>Little Bentwing-bat            | V                 |                      | Yes | Tropical rainforest to warm-temperate wet and dry sclerophyll forest; caves or similar structures for roosting.                                                                                                                                                                                                                                                                                                          | <b>Moderate - High</b><br>Suitable habitat areas containing hunting habitat were present along the REZ route. No preferred roosting habitat was present. |
| <i>Miniopterus orianae oceanensis</i><br>Large Bentwing-bat    | V                 |                      | Yes | Wet and dry tall open forest, rainforest, monsoon forest, open woodland, paperbark forests and open grasslands, caves or similar structures for roosting. It occasionally uses tree hollows.                                                                                                                                                                                                                             | <b>Moderate - High</b><br>Suitable habitat areas containing hunting habitat were present along the REZ route. No preferred roosting habitat was present. |
| <i>Myotis macropus</i><br>Southern Myotis                      | V                 |                      | No  | Various habitats of the coast and adjacent ranges with suitable waterbodies for hunting; caves or similar structures for roosting. It occasionally uses tree hollows.                                                                                                                                                                                                                                                    | <b>Moderate - High</b><br>Suitable habitat areas containing hunting and roosting habitat were present along the REZ route.                               |
| <i>Nyctophilus corbeni</i><br>Corben's Long-eared Bat          | V                 | V                    | No  | Inhabits a variety of vegetation types, including mallee, bull oak <i>Allocasuarina leuhmanni</i> and box eucalypt dominated communities, but it is distinctly more common in box/ironbark/cypress-pine vegetation that occurs in a north-south belt along the western slopes and plains of NSW and southern Queensland. Roosts in tree hollows, crevices, and under loose bark.                                         | <b>Low - Moderate</b><br>Suitable habitat areas containing hunting and roosting habitat were present along the REZ route.                                |
| <i>Scoteanax rueppellii</i><br>Greater Broad-nosed Bat         | V                 |                      | No  | Tree-lined creeks, woodland/clearing ecotones and rainforest creeks, roosting mainly in tree hollows.                                                                                                                                                                                                                                                                                                                    | <b>Moderate - High</b><br>Suitable habitat areas containing hunting and roosting habitat were present along the REZ route.                               |
| <i>Chalinolobus dwyeri</i><br>Large Pied Bat                   | E1                | E                    | Yes | Occupies dry sclerophyll forest and woodland. Roosts in caves, abandoned mud-nests of Fairy Martins and mine tunnels.                                                                                                                                                                                                                                                                                                    | <b>Low</b><br>Suitable habitat areas containing hunting habitat were present along the REZ route. No preferred roosting habitat was present.             |
| <i>Vespadelus troughtoni</i><br>Eastern Cave Bat               | V                 |                      | Yes | The Eastern Cave Bat is found in a broad band on both sides of the Great Dividing Range from Cape York to Kempsey, with records from the New England Tablelands and the upper north coast of NSW. The western limit appears to be the Warrumbungle Range, and there is a single record from southern NSW, east of the ACT. A cave-roosting species that is usually found in dry open forest and woodland, near cliffs or | <b>Low - Moderate</b><br>Suitable habitat areas containing hunting habitat were present along the REZ route. No preferred roosting habitat was present.  |



| SPECIES | STATUS            |                      |     | HABITAT DESCRIPTION AND LOCALLY KNOWN POPULATIONS                                                                        | LIKELIHOOD OF OCCURRENCE<br>WITHIN THE REZ SITE |
|---------|-------------------|----------------------|-----|--------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------|
|         | BC<br>Act<br>2016 | EPB<br>C Act<br>1999 | SAI |                                                                                                                          |                                                 |
|         |                   |                      |     | rocky overhangs; has been recorded roosting in disused mine workings, occasionally in colonies of up to 500 individuals. |                                                 |

## 5.4 FAUNA APPRAISAL RESULTS

### 5.4.1 DIURNAL SURVEYS

#### Amphibians

Suitable habitat was found to be present for a number of species of amphibian species in the form of drainage lines, creeks, rivers, dams and wetlands. Species recorded during fieldwork were *Crinia signifera* (Common Eastern Froglet), *Limnodynastes tasmaniensis* (Spotted Grass Frog) and *Litoria fallax* (Dwarf Green Tree Frog).

No threatened amphibian species listed as threatened under either the NSW BC Act 2016 and National EPBC Act 1999 were recorded.

#### Reptiles

Suitable habitat was found to be present for a number of species of reptile within the REZ site. Reptile species recorded during fieldwork included *Lampropholis delicata* (Grass Skink), *Ctenotus robustus* (Striped Skink), *Carlia tetradactyla* (Rainbow Skink), *Anomalopus leuckartii* (Two-clawed worm-skink), *Cryptoblepharus pulcher* (Fence Skink), *Pogona barbata* (Eastern Bearded Dragon) and *Varanus varius* (Lace Monitor).

No threatened reptile species listed as threatened under either the NSW BC Act 2016 and National EPBC Act 1999 were recorded. One newly described threatened reptile species *Delma vescolineata* (Hunter Valley Delma), listed as Endangered under both the BC Act 2026 and National EPBC Act 1999 has been recorded within the vicinity of the Muswellbrook STS (Wildthing Environmental Consultants, 2024 & WPS, 2021) within similar habitat in the form of woodland/derived native grassland.

#### Avifauna

An array of avifauna species was present within the REZ impact area. Common species recorded were *Gymnorhina tibicen* (Australian Magpie), *Trichoglossus moluccanus* (Rainbow Lorikeet), *Caligavis chrysops* (Yellow-faced Honeyeater), *Malurus cyaneus* (Superb Blue Wren); *Philemon corniculatus* (Noisy Friarbird), *Manorina melanocephala* (Noisy Miner), *Grallina cyanoleuca* (Magpie Lark), *Platycercus eximius* (Eastern Rosella) and *Corvus coronoides* (Australian Raven).

Three threatened avifauna species *Pomatostomus temporalis temporalis* (Grey-crowned Babbler), *Climacteris picumnus victoriae* (Brown Treecreeper) and *Haliaeetus leucogaster* (White-bellied Sea-Eagle) were recorded during fieldwork. Groups of *P. temporalis temporalis* were recorded at a number of locations together with their distinct spherical nests/roosts. *Climacteris picumnus victoriae* was recorded along the easement to the north-east of the Muswellbrook STS. One specimen of *H. leucogaster* was observed flying over the proposed site for the Eastern Hub near the northern shore of Lake Liddell.



These three species have been further assessed under the BC Act 2016 in Section 7.0. As *C. picumnus victoricae* is also listed as a threatened species under the EPBC Act 1999 the impact on this species has also been assessed in Section 9.0.

## Mammals

Native mammals recorded during fieldwork were *Macropus giganteus* (Eastern Grey Kangaroo) and *Notamacropus rufogriseus* (Red-necked Wallaby). Scats and digging of *Vombatus ursinus* (Common Wombat) were also noted.

A number of introduced mammals were observed within the REZ land and included *Oryctolagus cuniculus* (European Rabbit), *Lepus capensis* (European Hare) and *Mus musculus* (House Mouse). Secondary evidence such as diggings and foot prints also identified the presence of *Sus scrofa* (Feral Pig) and *Cervus elaphus* (Red Deer) were common in some areas.

None of these mammal species are listed as threatened under either the NSW BC Act 2016 and National EPBC Act 1999.

## **5.5 SURVEY LIMITATIONS**

As with all reports of this type the main survey limitation is considered to be the very short period of time in which the fieldwork was carried out. Limitations to the likelihood of detecting certain subject species were also encountered during this survey. Such limitations were generally related to the seasonal occurrence of species, be it as a result of known flowering periods for flora or migratory movements by fauna.

These limitations have been overcome by applying the precautionary principle in all cases where the survey methodology may have given a false negative result. This precautionary principle was achieved by recognising that most threatened species are rare and therefore unlikely to be encountered during a survey even if they may utilise the site at other times. These species have been assessed on the basis of the presence of their habitat and the likely significance of that habitat to a viable local population.

## **6.0 IMPACT ASSESSMENT**

### **6.1 AVOIDANCE AND MINIMISATION OF IMPACTS**

The majority of the 132kV transmission line will be contained within an existing easement which will avoid impacting large areas of native vegetation. As the easement is required to be widened in some sections, native vegetation occurring within these areas will be impacted. These impacts will be limited to tree branch trimming and some tree removal. Areas impacted by the Muswellbrook STS upgrade and new sections of easement particularly at Lower Belford along the New England Highway, along the Golden Highway (Mitchell Line of Road) and between the Kurri STS and Hunter Expressway will require removal of native vegetation. The area of the Eastern Hub near Lake Liddell will be situated within an already cleared grassy area.

### **6.2 DIRECT IMPACT**

The proposal will result in the following direct and potential impacts/losses:

- Impact to areas of native vegetation along the 132kV transmission line will impact 11 Plant Community Types (PCT's):
  - PCT 3631 Kurri Sand Clay Woodland
  - PCT 3433 Hunter Coast Foothills Spotted Gum-Ironbark Forest
  - PCT 3634 Quorrobolong Sand Flats Forest
  - PCT 3444 Lower Hunter Spotted Gum-Ironbark Forest
  - PCT 3446 Lower North Foothills Spotted Gum-Ironbark Grassy Forest
  - PCT 4023 Coastal Valleys Riparian Forest
  - PCT 3315 Central Hunter Ironbark-Spotted Gum Forest
  - PCT 4015 Central Hunter Swamp Oak Riparian Forest
  - PCT 4073 Lower North Hinterland River Oak Forest
  - PCT 3431 Central Hunter Ironbark Grassy Woodland
  - PCT 3525 Upper Hunter Box-Blakely's Red Gum Grassy Forest
- Impact to 7 Ecological Communities listed under the BC Act 2016:
  - Kurri Sand Swamp Woodland in the Sydney Basin Bioregion;
  - Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW;
  - Hunter Lowland Redgum Forest in the Sydney Basin and New South Wales North Coast Bioregions;
  - Central Hunter Ironbark-Spotted Gum-Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions;
  - Central Hunter Grey Box – Ironbark Woodland in the NSW North Coast and Sydney Basin Bioregions
  - Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East;
  - White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grassland
- Impact to 4 Nationally Listed Threatened Ecological Communities:
  - Kurri sand swamp woodland of the Sydney Basin bioregion;



Central Hunter Valley eucalypt forest and woodland;

Swamp Oak Floodplain Forest;

White Box – Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland

- Impact up to 32 specimens of the state and nationally threatened *Eucalyptus parramattensis* subsp. *decadens* (Drooping Red Gum);
- Impact to a small number of specimens of *Eucalyptus glaucina* (Slaty Red Gum), a state and nationally threatened species;
- Impact to a small number of specimens of *Acacia pendula* (Weeping Myall) – population in the Hunter Catchment;
- Impact to up 35 significant (larger trees) including up to 16 hollow-bearing trees;
- Impacts to habitat for the BC Act listed species threatened fauna species *Pomatostomus temporalis* subsp. *temporalis* (Grey-crowned Babbler);
- Impacts to habitat for the BC Act listed species threatened fauna species *Haliaeetus leucogaster* (White-bellied Sea-Eagle);
- Impacts to the BC Act 2026 & national EPBC Act 1999 listed species *Climacteris picumnus victoriae* (Brown Treecreeper);
- Removal/modification of a suitable habitat for a number of the addressed threatened flora and fauna species;

Details of PCT areas impacted by the proposal is shown in Table 6.1.

### 6.3 INDIRECT IMPACTS

The proposal may result in the following indirect and potential impacts:

- Trees and other Native Vegetation
- Potential unintentional impact to a small number of specimens of *Diuris tricolor* (Pine Donkey Orchid) and *Grevillea parviflora* subsp. *parvifolia* (Small-flowered Grevillea) within the existing easement;
- Impact on native fauna during vegetation removal:
- Erosion and sedimentation, particularly in relation to vegetation removal along streambanks;
- Introduction of priority, other weed species into other areas.

**Table 6.1: Details of impacts to Plant Community Types (PCT's).**

| PCT                                                                | Condition                              | BC Act 2016                                                                                                              | EPBC Act 1999                                                                            | Total Area (ha) |
|--------------------------------------------------------------------|----------------------------------------|--------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------|-----------------|
| PCT 3315 Central Hunter Ironbark-Spotted Gum Forest                | Existing Easement (Managed Vegetation) |                                                                                                                          |                                                                                          | 42.96           |
| PCT 3328 Lower Hunter Red Gum-Paperbark Riverflat Forest           | Existing Easement (Managed Vegetation) |                                                                                                                          |                                                                                          | 0.07            |
| PCT 3431 Central Hunter Ironbark Grassy Woodland                   | Existing Easement (Managed Vegetation) |                                                                                                                          |                                                                                          | 17.53           |
| PCT 3433 Hunter Coast Foothills Spotted Gum-Ironbark Grassy Forest | Existing Easement (Managed Vegetation) | EEC - Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW North Coast Bioregions                        |                                                                                          | 3.6             |
| PCT 3444 Lower Hunter Spotted Gum-Ironbark Forest                  | Existing Easement (Managed Vegetation) | EEC - Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW North Coast Bioregions                        |                                                                                          | 0.88            |
| PCT 3446 Lower North Foothills Ironbark-Box-Gum Grassy Forest      | Existing Easement (Managed Vegetation) | Hunter Lowland Redgum Forest in the Sydney Basin and New South Wales North Coast Bioregions                              |                                                                                          | 0.29            |
| PCT 3525 Upper Hunter Box-Blakelys Red Gum Grassy Forest           | Existing Easement (Managed Vegetation) |                                                                                                                          |                                                                                          | 9.78            |
| PCT 3631 Kurri Sand Clay Woodland                                  | Existing Easement (Managed Vegetation) | EEC - Kurri Sand Swamp Woodland in the Sydney Basin Bioregion                                                            | EEC - Kurri sand swamp woodland of the Sydney Basin bioregion                            | 2.15            |
| PCT 3634 Quorrobolong Sand Flats Forest                            | Existing Easement (Managed Vegetation) | EEC - Hunter Lowland Redgum Forest in the Sydney Basin and New South Wales North Coast Bioregions                        |                                                                                          | 0.79            |
| PCT 4015 Central Hunter Swamp Oak Riparian Forest                  | Existing Easement (Managed Vegetation) |                                                                                                                          |                                                                                          | 0.44            |
| PCT 4023 Coastal Valley Riparian Forest                            | Existing Easement (Managed Vegetation) |                                                                                                                          |                                                                                          | 0.74            |
| PCT 4073 Lower North Hinterland River Oak Forest                   | Existing Easement (Managed Vegetation) |                                                                                                                          |                                                                                          | 4.14            |
| PCT 3315 Central Hunter Ironbark-Spotted Gum Forest                | Derived Grassland                      | EEC - Central Hunter Ironbark-Spotted Gum-Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions |                                                                                          | 0.95            |
| PCT 3431 Central Hunter Ironbark Grassy Woodland                   | Derived Grassland                      |                                                                                                                          |                                                                                          | 2.4             |
| PCT 3315 Central Hunter Ironbark-Spotted Gum Forest                | Forested                               | EEC - Central Hunter Ironbark-Spotted Gum-Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions | Central Hunter Valley Eucalypt Forest and Woodland                                       | 13.27           |
| PCT 3431 Central Hunter Ironbark Grassy Woodland                   | Forested                               | EEC - Central Hunter Grey Box-Ironbark Woodland in the New South Wales North Coast and Sydney Basin Bioregions           | Central Hunter Valley Eucalypt Forest and Woodland                                       | 4.61            |
| PCT 3433 Hunter Coast Foothills Spotted Gum-Ironbark Grassy Forest | Forested                               | EEC - Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW North Coast Bioregions                        |                                                                                          | 0.31            |
| PCT 3444 Lower Hunter Spotted Gum-Ironbark Forest                  | Forested                               | EEC - Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW North Coast Bioregions                        |                                                                                          | 0.22            |
| PCT 3525 Upper Hunter Box-Blakelys Red Gum Grassy Forest           | Forested                               | White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grasslands                                 | White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grasslands | 0.59            |
| PCT 3631 Kurri Sand Clay Woodland                                  | Forested                               | EEC - Kurri Sand Swamp Woodland in the Sydney Basin Bioregion                                                            | EEC - Kurri sand swamp woodland of the Sydney Basin bioregion                            | 0.86            |
| PCT 3634 Quorrobolong Sand Flats Forest                            | Forested                               |                                                                                                                          |                                                                                          | 0.17            |

| PCT                                               | Condition                                                                     | BC Act 2016                                                                                                         | EPBC Act 1999                    | Total Area (ha) |
|---------------------------------------------------|-------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------|----------------------------------|-----------------|
| PCT 4015 Central Hunter Swamp Oak Riparian Forest | Forested                                                                      | Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions       | Swamp Oak Floodplain Forest      | 0.1             |
| PCT 4023 Coastal Valley Riparian Forest           | Forested                                                                      | EEC - Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions | EEC- Swamp Oak Floodplain Forest | 0.08            |
| PCT 4073 Lower North Hinterland River Oak Forest  | Forested                                                                      | EEC - Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions |                                  | 0.73            |
| Introduced <i>Juncus acutus</i>                   |                                                                               |                                                                                                                     |                                  | 0.46            |
| Native Regeneration                               |                                                                               |                                                                                                                     |                                  | 6.68            |
| Planted native trees                              |                                                                               |                                                                                                                     |                                  | 0.55            |
| Planted native vegetation                         |                                                                               |                                                                                                                     |                                  | 0.49            |
|                                                   | Total native vegetation impacted by proposal outside managed easements.       |                                                                                                                     |                                  | 32.47           |
|                                                   | Total native vegetation with existing easement vegetation (under management). |                                                                                                                     |                                  | 83.37           |



## 6.4 MITIGATION MEASURES

Mitigation measures have been specified to minimise the impact of the vegetation clearance to protect biodiversity values. The measures will include:

### Trees and other Native Vegetation

Trees and native vegetation within the subject land are to be avoided wherever possible within the scope of the proposal. Works should avoid any impact to native vegetation outside the impact area. Where unavoidable, works should minimise impacts to trees as follows:

- clearing limits will be clearly marked on the ground to prevent unnecessary clearing beyond the extent of the proposal;
- trees to be removed or trimmed are to be clearly marked to prevent any unintentional impact on trees that are to remain untouched;
- the clearing or trimming of any trees should be undertaken in a manner that avoids damaging adjacent vegetation;
- tree removal and branch trimming will be carried out in accordance with Ausgrid's Tree Safety Management Plan;
- all material stockpiles, vehicle parking and machinery storage will be located outside areas of native vegetation.

### Threatened Flora species

Threatened flora species such as *Diuris tricolor* (Pine Donkey Orchid) and *Grevillea parviflora* subsp. *parvifolia* (Small-flowered Grevillea) occurring within the existing easement are to be avoided during the construction of the transmission lines. The locations of these species should be located and marked, allowing them to be avoided.

### *Acacia pendula* (Weeping Myall) – population in the Hunter Catchment:

A fire had relatively recently burnt through all the specimens of *A. pendula* within a small area at Warkworth (first half of 2026) leaving burnt stems. As the non-pendulous form of *A. pendula* readily suckers from the roots (Umwelt 2006), these plants are likely to reshoot. It is thought that any suckering specimens of *A. pendula* within the new easement at Warkworth could remain *in-situ* and be managed to not impact the overhead wires. The area of *A. pendula* at Warkworth will be clearly marked on the ground to prevent unnecessary impacts.

### Erosion

Temporary erosion and sediment controls (e.g., silt fences) are to be installed to avoid disturbance and degradation of soils and nearby features. These controls should conform to the specifications in Soils and Construction 'Blue Book' (Landcom, 2004) and should be maintained throughout the construction process until soil is successfully stabilised.

Stumps of cut trees required to be removed along waterways such as Glennies and Bowmans Creek will be left *in-situ* to continue to hold the stream bank together.

### Weeds & Pathogens

Before entering the impact areas:

- All movement is to stay within the impact area,
- Use routes as instructed by the manager in the plan for the works
- Ensure machinery, vehicles clothing and equipment have been cleaned and inspected prior to entering new areas to prevent the spread of new plant propagules (seeds and fragments of plants), – Areas to inspect and clean within vehicles include tyres, wheel arches, under carriages, radiator grills, floor mats, load areas, boots, socks or anywhere that seeds and soil can lodge.

### Native Fauna

The removal of hollow-bearing trees is to be supervised by a suitably qualified ecologist from Ausgrid or contract ecologist to reduce the impact on any fauna which may be present. Hollow-bearing trees will be required to be flagged with tape or paint prior to works to ensure they are removed under the supervision of the ecologist.

In relation to the Grey-crowned Babbler the following is to be undertaken:

- Pre-clearance searches are to be undertaken for Grey-crowned Babbler nests prior to vegetation removal;
- Removal of nests is to be avoided where possible within the scope of the proposal;
- Where the removal of nests is unavoidable, nests are to be assessed for usage. If nests are found to be active, nests are not to be removed until all young have fledged the nest;
- If removal of a nest is unavoidable then the nest is to be removed by a suitably qualified ecologist and relocated outside the impact area a short distance away.

## 7.0 CONSIDERATIONS UNDER SECTION 7.3 OF THE BC ACT 2016

A summary of the impacts for the REZ under *Section 7.3* of the BC Act (2016) for the concerned threatened ecological communities and species is given below. The threatened communities and species dealt with are those identified during the fieldwork and those identified as having potential habitat available on site in Table 4.3. Individual assessments for threatened communities and species recorded within the REZ or close proximity under *Section 7.3* of the BC Act (2016) are contained in Appendix C.

Seven Threatened Ecological Communities were found to be present within the REZ. These were;

- Kurri Sand Swamp Woodland in the Sydney Basin Bioregion;
- Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW;
- Hunter Lowland Redgum Forest in the Sydney Basin and New South Wales North Coast Bioregions;
- Central Hunter Ironbark-Spotted Gum-Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions;
- Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East;
- Central Hunter Grey Box-Ironbark Woodland in the New South Wales North Coast and Sydney Basin Bioregions
- White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grassland

The proposal will result in an incremental reduction in each of these threatened ecological communities, particularly areas of Central Hunter Ironbark-Spotted Gum-Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions; however, it is considered unlikely to have a significant impact which might lead to extinction locally.

Four threatened flora species were recorded during fieldwork, these were

- *Eucalyptus glaucina* (Slaty Gum)
- *Eucalyptus parramattensis* subsp. *decadens* (Drooping Red Gum)
- *Grevillea parviflora* subsp. *parviflora* (Small-flowered Grevillea)
- *Diuris tricolor* (Pine Donkey Orchid)

The proposal will directly impact a small number of specimens of *Eucalyptus glaucina* (Slaty Gum) and *Eucalyptus parramattensis* subsp. *decadens* (Drooping Red Gum). Specimens of *Grevillea parviflora* subsp. *parviflora* (Small-flowered Grevillea) and *Diuris tricolor* (Pine Donkey Orchid) were recorded within the existing easement and are not required to be impacted by the proposal. However, there is a chance these flora species will be inadvertently impacted during the construction of the transmission line. Considerations under Section 7.3 of the BC Act (2016) (contained within Appendix D) found the REZ is unlikely to significantly affect the life cycle of any of these flora species or place any viable local populations of these species at risk of extinction.



No additional flora species were recorded within the survey area during fieldwork. Suitable habitat of varying quality was considered to be present for 14 of the 53 threatened flora species listed in Table

4.3. These flora species were:

- *Corybas dowlingii* (Red Helmet Orchid);
- *Prasophyllum* sp. Wybong (A Leek Orchid)
- *Pterostylis chaetophora* (Tall Rustyhood)
- *Rhizanthella slateri* (Eastern Underground Orchid);
- *Dichanthium setosum* (Blue Grass)
- *Brachyscome brownii*
- *Ozothamnus tessellatus*
- *Rutidosia heterogama* (Heath Wrinklewort)
- *Acacia bynoeana* (Bynoe's Wattle)
- *Callistemon linearifolius* (Netted Bottle Brush);
- *Persicaria elatior* (Tall Knotweed)
- *Persoonia pauciflora* (North Rothbury Persoonia)
- *Thesium australe* (Austral Toadflax)
- *Zannichellia palustris*

According to BioNet records (NSW DCCEEW, 2026a) and presence of suitable habitat the most likely of these flora species to be present were *Rutidosia heterogama* (Heath Wrinklewort), *Callistemon linearifolius* (Netted Bottle Brush), *Acacia bynoeana* (Bynoe's Wattle) and *Ozothamnus tessellatus*. The proposal will result in an incremental reduction in habitat, however considering the relatively small impact, the absence of these species during fieldwork and areas of similar surrounding habitat it is considered unlikely that it would significantly affect the life cycle of any of these flora species or place any viable local populations of these species at risk of extinction.

### Threatened Fauna

Three threatened fauna species were recorded during fieldwork. These species were:

- *Pomatostomus temporalis* subsp. *temporalis* (Grey-crowned Babbler);
- *Climacteris picumnus victoriae* (Brown Treecreeper);
- *Haliaeetus leucogaster* (White-bellied Sea-Eagle).

The proposal will impact suitable habitat for *Pomatostomus temporalis* subsp. *temporalis* (Grey-crowned Babbler) and *Climacteris picumnus victoriae* (Brown Treecreeper). A small number of nests/roosts of *P. temporalis* subsp. *temporalis* may also be impacted. No suitable habitat is likely to be impacted for *Haliaeetus leucogaster* (White-bellied Sea-Eagle). Considerations under Section 7.3 of the BC Act (2016) (contained within Appendix D) found the REZ is unlikely to significantly affect the life cycle of any of these fauna species or place any viable local populations of these species at risk of extinction.

No additional threatened fauna species were recorded during fieldwork. Of the 90 addressed threatened fauna species, the REZ was considered to contain suitable habitat for 68 species:

- *Litoria aurea* (Green and Golden Bell Frog);
- *Litoria brevipalmata* (Green-thighed Frog);
- *Litoria littlejohni* (Littlejohn's Tree Frog);
- *Aprasia parapulchella* (Pink-tailed Worm-lizard);

- *Delma vescolineata* (Hunter Valley Delma)
- *Botaurus poiciloptilus* (Australasian Bittern);
- *Varanus rosenbergi* (Rosenberg's Goanna)
- *Calidris acuminata* (Sharp-tailed Sandpiper);
- *Tringa nebularia* (Common Greenshank)
- *Gallinago hardwickii* (Latham's Snipe)
- *Rostratula australis* (Australian Painted Snipe);
- *Botaurus poiciloptilus* (Australasian Bittern)
- *Ixobrychus flavicollis* (Black Bittern);
- *Ephippiorhynchus asiaticus* (Black-necked Stork)
- *Anseranas semipalmata* (Magpie Goose);
- *Oxyura australis* (Blue-billed Duck);
- *Stictonetta naevosa* (Freckled Duck)
- *Burhinus grallarius* (Bush Stone Curlew)
- *Turnix maculosus* (Red-backed Button-quail);
- *Calyptorhynchus lathami* (Glossy Black-Cockatoo);
- *Callocephalon fimbriatum* (Gang Gang Cockatoo);
- *Lathamus discolor* (Swift Parrot);
- *Neophema chrysostoma* (Blue-winged Parrot)
- *Neophema pulchella* (Turquoise Parrot)
- *Polytelis swainsonii* (Superb Parrot)
- *Glossopsitta pusilla* (Little Lorikeet);
- *Aphelocephala leucopsis* (Southern Whiteface);
- *Hirundapus caudacutus* (White-throated Needle-tail);
- *Artamus cyanopterus cyanopterus* (Dusky Woodswallow)
- *Melanodryas cucullata cucullata* (Hooded Robin) (south-eastern form)
- *Petroica boodang* (Scarlet Robin);
- *Petroica phoenicea* (Flame Robin);
- *Stagonopleura guttata* (Diamond Firetail);
- *Chthonicola sagittata* (Speckled Warbler);
- *Anthochaera phrygia* (Regent Honeyeater);
- *Certhionyx variegatus* (Pied Honeyeater);
- *Grantiella picta* (Painted Honeyeater);
- *Melithreptus gularis gularis* (Black-chinned Honeyeater);
- *Daphoenositta chrysoptera* (Varied Sittella);
- *Circus assimilis* (Spotted Harrier);
- *Lophoictinia isura* (Square-tailed Kite);
- *Hieraaetus morphnoides* (Little Eagle);
- *Hamirostra melanosternon* (Black-breasted Buzzard);
- *Falco hypoleucos* (Grey Falcon);
- *Falco subniger* (Black Falcon);
- *Ninox connivens* (Barking Owl);
- *Ninox strenua* (Powerful Owl);
- *Tyto novaehollandiae* (Masked Owl);
- *Dasyurus maculatus* subsp. *maculatus* (Spotted-tailed Quoll);
- *Phascogale tapoatafa* (Brush-tailed Phascogale);
- *Phascolarctos cinereus* (Koala)
- *Potorous tridactylus* sp. *tridactylus* (Long-nosed Potoroo)
- *Cercartetus nanus* (Eastern Pygmy-possum);

- *Petaurus australis* (Yellow-bellied Glider);
- *Petaurus norfolcensis* (Squirrel Glider);
- *Petauroides volans* (Greater Glider);
- *Pseudomys novaehollandiae* (New Holland Mouse)
- *Pteropus poliocephalus* (Grey-headed Flying-Fox);
- *Saccolaimus flaviventris* (Yellow-bellied Sheath-tail-bat);
- *Micronomus norfolkensis* (Eastern Freetail-bat);
- *Falsistrellus tasmaniensis* (Eastern False Pipistrelle);
- *Miniopterus australis* (Little Bentwing-bat);
- *Miniopterus schreibersii oceanensis* (Large Bentwing-bat);
- *Myotis macropus* (Southern Myotis)
- *Nyctophilus corbeni* (Corben's Long-eared Bat)
- *Scoteanax rueppellii* (Greater Broad-nosed Bat);
- *Chalinolobus dwyeri* (Large Pied Bat);
- *Vespadelus troughtoni* (Eastern Cave Bat).

According to the BioNet Atlas records (NSW DCCEEW, 2026a) threatened fauna species found within proximity of the REZ included *Delma vescolineata* (Hunter Valley Delma), *Daphoenositta chrysoptera* (Varied Sittella), *Chthonicola sagittata* (Speckled Warbler), *Glossopsitta pusilla* (Little Lorikeet), *Artamus cyanopterus cyanopterus* (Dusky Woodswallow), *Petaurus norfolcensis* (Squirrel Glider), *Phascogale tapoatafa* (Brush-tailed Phascogale), *Pteropus poliocephalus* (Grey-headed Flying-fox) and addressed microchiropteran bats. The proposal will result in a small incremental reduction habitat for the above species. Given the relatively small impact and recommendations it is unlikely that the proposal will have a significant impact on these threatened fauna species such that a local extinction would occur.

### Key Threatening Processes'

The Key Threatening Processes currently listed under Schedule 4 of the BC Act 2016 that are relevant to the REZ area have been listed in Table 7.1.

**Table 7.1: Key Threatening Processes.**

| Key Threatening Process                                           | Applicability in regards to the subject site                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|-------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Clearing of Native Vegetation.                                    | The proposal will result in the removal of native vegetation and may be viewed as being part of this Key Threatening Process. The action is unlikely to be responsible for the significant loss of this EEC provided that recommendations for impact minimisation as listed within Section 6.4 are undertaken.                                                                                                                                                                                                                                                             |
| Loss of Hollow-bearing Trees                                      | Up to 16 hollow-bearing trees are proposed to be removed.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Bushrock Removal                                                  | No significant areas of bushrock are likely to be impacted.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| Invasion of native plant communities by exotic perennial grasses. | Specimens of exotic perennial grasses such as <i>Chloris gayana</i> (Rhodes Grass), <i>Eragrostis curvula</i> (African Love Grass), <i>Andropogon virginicus</i> (Whisky Grass) and <i>Hyparrhenia hirta</i> (Coolatai Grass) were present. These exotic grasses compete with species within this EEC for space, light, and other resources and may threaten its persistence at several sites. The action is unlikely to be responsible for the significant loss of this EEC provided recommendations for impact minimisation as listed within Section 6.4 are undertaken. |



| Key Threatening Process                                                                                                                                                                                                                                     | Applicability in regards to the subject site                                                                                                                                                                                                                    |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Predation by the <i>Felis catus</i> (Feral Cat)                                                                                                                                                                                                             | The Feral Cat was not recorded on site at the time of the survey however this species would be considered to have an impact on native fauna in the local area. The proposal is not likely to result in an increase in feral numbers of this introduced species. |
| Predation by the <i>Vulpes vulpes</i> (Red Fox)                                                                                                                                                                                                             | The Red Fox was not recorded on site at the time of the survey however this species would be considered to have an impact on native fauna in the local area. The proposal is not likely to result in an increase in numbers of this introduced species.         |
| Predation by the <i>Gambusia holbrooki</i> (Plague Minnow)                                                                                                                                                                                                  | The Plague Minnow was observed within a number of streams and waterbodies along the REZ. The proposal is not likely to result in an increase in numbers of this introduced species.                                                                             |
| Tracks of Feral Deer were recorded within several sections of the REZ. Feral Deer would be considered to have an impact on native flora in the local area. The proposal is not likely to result in an increase in feral numbers of this introduced species. | Tracks of Feral Deer were recorded within several sections of the REZ. Feral Deer would be considered to have an impact on native flora in the local area. The proposal is not likely to result in an increase in feral numbers of this introduced species.     |
| Predation, habitat degradation, competition and disease transmission by <i>Sus scrofa</i> (Feral Pigs)                                                                                                                                                      | Signs of Feral Pigs were noted along parts of the REZ route. The proposal is not likely to result in an increase in feral numbers of this introduced species.                                                                                                   |
| Infection by Psittacine Circoviral (beak and feather) Disease affecting endangered psittacine species and populations                                                                                                                                       | No evidence of the disease was observed on psittacine species. No endangered Psittacine species were seen on site. The proposal is unlikely to increase infection by this disease.                                                                              |
| Aggressive exclusion of birds by noisy miners ( <i>Manorina melanocephala</i> )                                                                                                                                                                             | Noisy miners were not recorded within the REZ during fieldwork. The proposal is unlikely to increase the impacts associated with this species.                                                                                                                  |
| Introduction and establishment of Exotic Rust Fungi of the order Pucciniales pathogenic on plants of the family Myrtaceae.                                                                                                                                  | No evidence of the fungi was observed during fieldwork.                                                                                                                                                                                                         |
| Competition and grazing by the feral European rabbit, <i>Oryctolagus cuniculus</i> .                                                                                                                                                                        | Observations and secondary indications of the European Rabbit such as scats and diggings were noted along parts of the REZ. The proposal is not likely to result in an increase in feral numbers of this introduced species.                                    |
| High frequency fire resulting in the disruption of life cycle processes in plants and animals and loss of vegetation structure and composition.                                                                                                             | It is unknown what impact fire has had within the site.                                                                                                                                                                                                         |
| Herbivory and environmental degradation caused by feral deer.                                                                                                                                                                                               | Tracks of Feral Deer were recorded within several sections of the REZ. Feral Deer would be considered to have an impact on native flora in the local area. The proposal is not likely to result in an increase in feral numbers of this introduced species.     |
| Infection of frogs by amphibian chytrid causing the disease chytridiomycosis                                                                                                                                                                                | No evidence of chytrid was observed during fieldwork.                                                                                                                                                                                                           |
| Infection of native plants by <i>Phytophthora cinnamomi</i>                                                                                                                                                                                                 | No evidence of <i>Phytophthora cinnamomi</i> was observed during site visits. The proposal is unlikely to cause the spread of this fungus.                                                                                                                      |
| Invasion and establishment of exotic vines and scramblers                                                                                                                                                                                                   | Exotic vines and scramblers such as <i>Cardiospermum grandiflorum</i> (Balloon vine) were common in Riparian area along the REZ.                                                                                                                                |
| Invasion of native plant communities by African Olive <i>Olea europaea</i> subsp. <i>cuspidata</i>                                                                                                                                                          | The African Olive was recorded along parts of the REZ route. The proposal is unlikely to cause further spread of this introduced tree.                                                                                                                          |
| Invasion, establishment and spread of Lantana ( <i>Lantana camara</i> L. sens. lat)                                                                                                                                                                         | Lantana was recorded within the site.                                                                                                                                                                                                                           |
| Removal of dead wood and dead trees.                                                                                                                                                                                                                        | Dead wood and trees were recorded along small sections of the REZ route. Dead wood and trees such be avoided wherever possible.                                                                                                                                 |

## 8.0 ASSESSMENT OF SERIOUS AND IRREVERSIBLE IMPACTS

Under the BC Act 2016, a determination of whether an impact is serious and irreversible (SAIL) must be made in accordance with the principles prescribed in section 6.7 of the BC Regulation.

The “*Guidance to assist a decision maker to determine a serious and irreversible impact*, 2017, sets out those potential SAIL species and ecological communities (known as “potential SAIL entities”).

The principles for determining serious and irreversible impacts in the Biodiversity Conservation Regulation, 2017 are:

- *will cause a further decline of a species or ecological community that is currently observed, estimated, inferred or reasonably suspected to be in a rapid rate of decline, or*
- *will further reduce the population of a species or ecological community that is currently observed, estimated, inferred, or reasonably suspected to have a very small population size, or*
- *are impacts on the habitat of a species or area of ecological community that is currently observed, estimated, inferred or reasonably suspected to have a very limited geographic distribution, or*
- *are impacts on a species or ecological community is unlikely to respond to measures to improve habitat and vegetation integrity and is therefore irreplaceable.*

### 8.1 POTENTIAL SAIL ENTITIES

In this case all potential SAIL entities are derived from Appendix 2 of the Guide, and are within the BioNet search area (DPE, 2023). The approval authority must take those impacts into consideration and determine whether there are any additional and appropriate measures that will minimise those impacts if approval is to be granted. An Impact evaluation is shown in Table 9.1. Entities include:

- Central Hunter Valley Eucalypt Forest and Woodland
- White Box Yellow Box Blakely's Red Gum Woodland
- *Rhizanthella slateri* (Eastern Underground Orchid);
- *Lathamus discolor* (Swift Parrot);
- *Anthochaera phrygia* (Regent Honeyeater);
- *Miniopterus australis* (Little Bentwing-bat);
- *Chalinolobus dwyeri* (Large Pied Bat);
- *Vespadelus troughtoni* (Eastern Cave Bat);

**Table 8.1: SAIL impact evaluation**

| Potential SAIL Entities                                   | Impact Evaluation                  | Impact Thresholds                                  | Serious and Irreversible Impact? |
|-----------------------------------------------------------|------------------------------------|----------------------------------------------------|----------------------------------|
| Central Hunter Valley Eucalypt Forest and Woodland        | Present within the subject land    |                                                    | No                               |
| White Box Yellow Box Blakely's Red Gum Woodland           | Present within the subject land    | No threshold identified.                           | No                               |
| <i>Rhizanthella slateri</i><br>Eastern Underground Orchid | Only marginal habitat was present. |                                                    | No                               |
| <i>Lathamus discolor</i><br>Swift Parrot                  | Suitable habitat was present.      | Not within a mapped BAM Important Area (DPE, 2023) | No                               |

| Potential SAI Entities                              | Impact Evaluation                                                                                  | Impact Thresholds                                  | Serious and Irreversible Impact? |
|-----------------------------------------------------|----------------------------------------------------------------------------------------------------|----------------------------------------------------|----------------------------------|
| <i>Anthochaera phrygia</i><br>Regent Honeyeater     | Seasonal foraging habitat was present.                                                             | Not within a mapped BAM Important Area (DPE, 2023) | No                               |
| <i>Miniopterus australis</i><br>Little Bentwing-bat | Suitable hunting habitat was present. Preferred roosting habitat was absent.                       |                                                    | No                               |
| <i>Chalinolobus dwyeri</i><br>Large Pied Bat        | Suitable hunting habitat was present. Preferred roosting habitat was absent.                       |                                                    | No                               |
| <i>Vespadelus troughtoni</i><br>Eastern Cave Bat    | Suitable hunting habitat was present. No preferred roosting habitat was available within the site. |                                                    | No                               |

Two SAI entities the TEC – Central Hunter Valley Eucalypt Forest and Woodland and White Box - Yellow Box - Blakely's Red Gum Woodland (Box Gum Woodland TEC) were identified within the REZ.



## 9.0 CONSIDERATIONS UNDER THE COMMONWEALTH ENVIRONMENT PROTECTION AND BIODIVERSITY CONSERVATION ACT 1999

Considerations have been made to the Commonwealth Environment Protection and Biodiversity Conservation (EPBC) Act 1999. Assessments have been made to determine whether or not the proposal or activity has, will have, or is likely to have a significant impact on a matter of National Environmental Significance. The matters of National Environmental Significance and the appropriate responses are listed below:

- *World Heritage properties;*

The REZ is unlikely to impact any World Heritage Properties.

- *wetlands recognised under the Ramsar convention as having international significance;*

The subject land is located upstream from the Hunter Estuary. The proposed works are not likely to have a significant impact to any Ramsar Wetlands.

- *listed threatened species and communities;*

Four nationally threatened ecological communities were recorded on the DCCEEW database as having potential to occur within 10km of the site, these being:

- Central Hunter Valley eucalypt forest and woodland;
- White Box-Yellow Box-Blakely's Red Gum Grassy Woodland and Derived Native Grassland
- Kurri sand swamp woodland of the Sydney Basin bioregion;
- Swamp Oak Floodplain Forest;

The assessment undertaken in Sections 10.1 – 10.4 found the proposal is unlikely to significantly impact any of these four communities.

A total of 52 nationally threatened species were recorded on the DCCEEW database as occurring or having potential habitat available within 10km of the site (excluding pelagic species), these being:

|                                        |                            |
|----------------------------------------|----------------------------|
| <i>Anthochaera phrygia</i>             | Regent Honeyeater          |
| <i>Calidris ferruginea</i>             | Curlew Sandpiper           |
| <i>Lathamus discolor</i>               | Swift Parrot               |
| <i>Callocephalon fimbriatum</i>        | Gang-gang Cockatoo         |
| <i>Melanodryas cucullata cucullata</i> | South-eastern Hooded Robin |
| <i>Botaurus poiciloptilus</i>          | Australasian Bittern       |
| <i>Rostratula australis</i>            | Australian Painted Snipe   |
| <i>Stagonopleura guttata</i>           | Diamond Firetail           |
| <i>Pycnoptilus floccosus</i>           | Pilotbird                  |
| <i>Polytelis swainsonii</i>            | Superb Parrot              |
| <i>Aphelocephala leucopsis</i>         | Southern Whiteface         |
| <i>Grantiella picta</i>                | Painted Honeyeater         |
| <i>Falco hypoleucos</i>                | Grey Falcon                |

|                                                                 |                                       |
|-----------------------------------------------------------------|---------------------------------------|
| <i>Hirundapus caudacutus</i>                                    | White-throated Needletail             |
| <i>Calidris acuminata</i>                                       | Sharp-tailed Sandpiper                |
| <i>Gallinago hardwickii</i>                                     | Latham's Snipe                        |
| <i>Calyptorhynchus lathamii lathamii</i>                        | South-eastern Glossy Black-Cockatoo   |
| <i>Climacteris picumnus victoriae</i>                           | Brown Treecreeper (south-eastern)     |
| <i>Neophema chrysostoma</i>                                     | Blue-winged Parrot                    |
| <i>Litoria booroolongensis</i>                                  | Booroolong Frog                       |
| <i>Heleioporus australiacus</i>                                 | Giant Burrowing Frog                  |
| <i>Phascolarctos cinereus</i>                                   | Koala                                 |
| <i>Chalinolobus dwyeri</i>                                      | Large-eared Pied Bat                  |
| <i>Dasyurus maculatus maculatus</i> (SE mainland population)    | Spot-tailed Quoll                     |
| <i>Petauroides volans</i>                                       | Greater Glider (southern and central) |
| <i>Petrogale penicillata</i>                                    | Brush-tailed Rock-wallaby             |
| <i>Pseudomys novaehollandiae</i>                                | New Holland Mouse                     |
| <i>Pteropus poliocephalus</i>                                   | Grey-headed Flying-fox                |
| <i>Petaurus australis australis</i>                             | Yellow-bellied Glider (south-eastern) |
| <i>Nyctophilus corbeni</i>                                      | Corben's Long-eared Bat               |
| <i>Euphrasia arguta</i>                                         |                                       |
| <i>Prasophyllum</i> sp. Wybong (C.Phelps ORG 5269)              | a leek-orchid                         |
| <i>Wollemia nobilis</i>                                         | Wollemi Pine                          |
| <i>Hibbertia acaulothrix</i>                                    |                                       |
| <i>Androcalva rosea</i>                                         | Sandy Hollow Commersonia              |
| <i>Persoonia hirsuta</i>                                        | Hairy Geebung                         |
| <i>Pomaderris cotoneaster</i>                                   | Cotoneaster Pomaderris                |
| <i>Cynanchum elegans</i>                                        | White-flowered Wax Plant              |
| <i>Vincetoxicum forsteri</i>                                    |                                       |
| <i>Prostanthera discolor</i>                                    |                                       |
| <i>Lepidium aschersonii</i>                                     | Spiny Peppercross                     |
| <i>Dichanthium setosum</i>                                      | bluegrass                             |
| <i>Ozothamnus tessellatus</i>                                   |                                       |
| <i>Prostanthera cryptandroides</i> subsp. <i>cryptandroides</i> | Wollemi Mint-bush                     |
| <i>Pomaderris brunnea</i>                                       | Rufous Pomaderris                     |
| <i>Homoranthus darwinioides</i>                                 |                                       |
| <i>Swainsona murrayana</i>                                      | Slender Darling-pea                   |
| <i>Androcalva procumbens</i>                                    |                                       |
| <i>Thesium australe</i>                                         | Austral Toadflax                      |
| <i>Hoplocephalus bungaroides</i>                                | Broad-headed Snake                    |
| <i>Delma vescolineata</i>                                       | Hunter Valley Delma                   |
| <i>Aprasia parapulchella</i>                                    | Pink-tailed Worm-lizard               |

One nationally threatened species; *Climacteris picumnus victoriae* (Brown Treecreeper) was recorded during fieldwork. Of the listed national threatened species, those most likely to occur within the site and close proximity would include the flora species such as *Pteropus poliocephalus* (Grey-headed Flying-fox), *Delma vescolineata* (Hunter Valley Delma), listed woodland birds, and microchiropteran bats. Potential habitat for all of the listed nationally threatened species has been assessed in Table 5.5. The action will result in a small incremental loss/modification of habitat within the locality for these species; however, it is not likely to have a significant impact on any of these addressed species.

- *migratory species protected under international agreements.*

Ten nationally listed migratory species were recorded on the DCCEEW on-line database as occurring or having potential habitat available within 10km of the study area, these being:

Migratory Terrestrial Species:

- *Hirundapus caudacutus* (White-throated Needletail)

- *Monarcha melanopsis* (Black-faced Monarch)
- *Motacilla flava* (Yellow Wagtail)
- *Myiagra cyanoleuca* (Satin Flycatcher)
- *Rhipidura rufifrons* (Rufous Fantail)

Migratory Wetland Species:

- *Actitis hypoleucos* (Common Sandpiper)
- *Calidris ferruginea* (Curlew Sandpiper)
- *Calidris melanotos* (Pectoral Sandpiper)
- *Gallinago hardwickii* (Latham's Snipe)
- *Numenius madagascariensis* (Eastern Curlew)

Migratory Marine Birds

- *Apus pacificus* (Fork-tailed Swift)

Considering the relatively small impact on habitat in the locality it is unlikely that these species or any of the listed migratory species would be significantly affected by the proposal.

8. *nuclear activities;*

The proposal does not involve any type of nuclear activity.

9. *the Commonwealth marine environment;*

The proposal does not involve the modification of the Commonwealth marine environment



### 9.1 CENTRAL HUNTER VALLEY EUCALYPT FOREST AND WOODLAND

The vegetation must be consistent with the criteria outlined in the 'Approved Conservation Advice (including listing advice) for the Central Hunter Valley eucalypt forest and woodland ecological community' (DotE 2015) (Table 9.1).

In order to be considered a Matter of National Environmental Significance under the EPBC Act, areas of the ecological community must meet:

1. the key diagnostic characteristics (in Section 1.5.1; DotE 2015) (Table 10.1); AND  
Areas of (PCT) 3431 - Central Hunter Ironbark Grassy Woodland and PCT 3315 Central Hunter Ironbark-Spotted Gum Forest met the key diagnostic characteristics of Central Hunter Valley eucalypt forest. An exception would be the areas of derived woodland which do not met the key diagnostic characteristics.

2. at least the minimum condition thresholds for species composition and patch sizes (DotE 2015).

#### Minimum Condition Thresholds

Minimum condition thresholds for species composition and patch sizes (DotE 2015).

(1) Is the patch at least 0.5 ha in size? -

Yes



(2) Is at least 50% of the perennial understorey vegetative cover of the patch (due to) native plants?

Yes, for a number of areas.



(3) Are there at least 12 native understorey species in the whole patch?

This would be true for a number of areas.



**The patch is part of the protected nationally listed ecological community**

Areas of (PCT) 3431 - Central Hunter Ironbark Grassy Woodland and PCT 3315 Central Hunter Ironbark-Spotted Gum Forest would be consistent with the Critically Endangered Ecological Community Central Hunter Valley eucalypt forest and woodland. An assessment Central Hunter Valley eucalypt forest and woodland was conducted and found that the proposal is unlikely to have a significant impact.

**Table 9.1: Assessment of EPBC Act Central Hunter Valley eucalypt forest and woodland. Key diagnostic characteristics against Plant Community Type (PCT)**

| Key diagnostic characteristics as described in in Section 1.5.1 of DotE 2015                                                                                                                                                                                                                                                                                                                                                                                                                                                          | Impacted Area<br>(PCT) 3431 with canopy (4.61ha)<br>(PCT) 3315 with canopy (13.24ha)                                                                                          | Impacted Area – 3ha<br>(PCT) 3431 (Derived Grassland) 2.4ha<br>(PCT) 3315 (Derived Grassland) 0.95ha |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------|
| It occurs in the Hunter River catchment (typically called the Hunter Valley region)                                                                                                                                                                                                                                                                                                                                                                                                                                                   | Yes                                                                                                                                                                           | Yes                                                                                                  |
| It typically occurs on lower hillslopes and low ridges, or valley floors in undulating country; on soils derived from Permian sedimentary rocks                                                                                                                                                                                                                                                                                                                                                                                       | Yes                                                                                                                                                                           | Yes                                                                                                  |
| It does not occur on alluvial flats, river terraces, aeolian sands, Triassic sediments, or escarpments                                                                                                                                                                                                                                                                                                                                                                                                                                | Yes                                                                                                                                                                           | Yes                                                                                                  |
| The canopy of the ecological community is dominated by one or more of the following four eucalypt species: <i>Eucalyptus crebra</i> (narrow-leaved ironbark), <i>Corymbia aculata</i> (syn. <i>E. maculata</i> ) (spotted gum), <i>E. dawsonii</i> (slaty gum) and <i>E. moluccana</i> (grey box) OR a fifth species, <i>Allocasuarina luehmanna</i> (bulloak, buloke) dominates in combination with one or more of the above four eucalypt species, in sites previously dominated by one or more of the above four eucalypt species. | Areas dominated by either <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark). Some specimens of <i>Eucalyptus moluccana</i> (Grey Box) | No canopy present                                                                                    |
| <i>Allocasuarina torulosa</i> (forest oak/ she-oak, rose she-oak/oak), <i>Eucalyptus acmenoides</i> (white mahogany) and <i>E. fibrosa</i> (red/broad-leaved ironbark) are largely absent from the canopy of a patch.                                                                                                                                                                                                                                                                                                                 | Yes                                                                                                                                                                           | Yes                                                                                                  |
| A ground layer is present (although it may vary in development and composition), as a sparse to thick layer of native grasses and other native herbs and/or native shrubs                                                                                                                                                                                                                                                                                                                                                             | Yes                                                                                                                                                                           | Yes                                                                                                  |
| <b>Does this PCT/condition zone meet the EPBC Act listing criterion</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                               | <b>Yes</b>                                                                                                                                                                    | <b>No</b>                                                                                            |

An action is likely to have a significant impact on a critically endangered or endangered ecological community if there is a real chance or possibility that it will:

- ***reduce the extent of an ecological community***

The proposed new easements along the southern side of the New England Highway at Lower Belford, Mitchell Line of Road at Mount Thorley and parts of Putty Road will require the direct removal of 13.27ha of native woodland vegetation (PCT 3315) consistent with the CEEC Central Hunter Valley eucalypt forest. An additional 4.61ha (PCT 3431) is required to be removed for the Muswellbrook STS and associated easements.

A large portion of the trees that will be impacted for the new proposed easement along the southern side of the New England Highway at Lower Belford were planted as part of rehabilitation for the road duplication undertaken in the early 1990's. This was evidenced by the remains of plastic tree guard around the bases of a number of trees.

- ***fragment or increase fragmentation of an ecological community, for example by clearing vegetation for roads or transmission lines***

The project involves the development of an area for the STS and easements for the overhead lines. As the relatively narrow proposal will not impact groundcovers and small shrubs within the easement it is unlikely to increase fragmentation between habitat areas due to the presence of similar adjoining habitat.

- ***adversely affect habitat critical to the survival of an ecological community***

The project is unlikely to adversely affect habitat considered to be critical to the survival of the community.

- ***modify or destroy abiotic (non-living) factors (such as water, nutrients, or soil) necessary for an ecological community's survival, including reduction of groundwater levels, or substantial alteration of surface water drainage patterns***

The project is unlikely to affect any abiotic processes necessary for the community's survival.

- ***cause a substantial change in the species composition of an occurrence of an ecological community, including causing a decline or loss of functionally important species, for example through regular burning or flora or fauna harvesting the following criteria:***

The proposal is unlikely to result in substantial change in the species composition of this community locally.

- ***cause a substantial reduction in the quality or integrity of an occurrence of an ecological community, including, but not limited to:***  
-assisting invasive species, that are harmful to the listed ecological community, to become established, or  
– causing regular mobilisation of fertilisers, herbicides or other chemicals or pollutants into the ecological community which kill or inhibit the growth of species in the ecological community, or

Taking the mitigations measures into consideration, the proposal is unlikely to assist any invasive species, nor cause any increased mobilisation of fertilisers which would kill or inhibit the growth of species in the ecological community.

- ***interfere with the recovery of an ecological community.***

The proposal is unlikely to interfere with any current recovery programs in the area.



### Conclusion

A large portion of the REZ through existing easements largely avoids the removal of CEEC Central Hunter Valley eucalypt forest. However, the REZ will impact up to 17.88ha of CEEC Central Hunter Valley eucalypt forest, largely along new sections of narrow new easement along the New England Highway at Lower Belford, Mitchell Line of Road and Putty Road at Mount Thorley and at the Muswellbrook STS.

Areas of CEEC Central Hunter Valley Eucalypt Forest ranged from areas of low to good condition. Some sections along the Mitchell Line of Road at Mount Thorley were in relatively good condition. A large portion of the trees that will be impacted for the new proposed easement along the southern side of the New England Highway at Lower Belford were planted as part of rehabilitation for the road duplication undertaken in the early 1990's. This was evidenced by the remains of plastic tree guard around the bases of a number of trees.

The proposal will result in an incremental reduction in area and quality of Central Hunter Valley Eucalypt Forest. Due to the narrowness of CEEC Central Hunter Valley Eucalypt Forest to be impacted, and taking into consideration the recommendations given in this report it is considered less likely that the proposed action would have an adverse effect on the extent of the TEC or substantially and adversely modify the composition of the TEC such that its local occurrence is likely to be placed at risk of extinction. Given these factors a referral is unlikely to be required.

### **9.2 WHITE BOX -YELLOW BOX- BLAKELYS RED GUM GRASSY WOODLAND AND DERIVED NATIVE GRASSLAND**

To be considered consistent with the Critically Endangered listing under the EPBC Act, the vegetation must be consistent with the criteria outlined in the EPBC Act policy statement 3.5 – White box – Yellow box – Blakely's red gum grassy woodlands and derived native grasslands (Department of the Environment and Heritage, 2006) and as summarised in Table 9.2.

The assessment concluded that the vegetation did meet the condition criteria of the EPBC Act listing for White Box – Yellow Box Blakely's Red Gum Grassy Woodland and Derived Native Grasslands. An assessment under the Significant Impact Criteria for this community has been undertaken below.

An action is likely to have a significant impact on a critically endangered or endangered ecological community if there is a real chance or possibility that it will:

- ***reduce the extent of an ecological community***

Areas mapped as PCT 3525 Upper Hunter Box-Blakelys Red Gum Grassy Forest were found to be consistent with the CEEC; White Box – Yellow Box Blakely's Red Gum Grassy Woodland and Derived Native Grasslands. This CEEC was present along a section of existing easement to the north-east of the Muswellbrook STS. This CEEC largely occurred within areas of existing easement, composed of low maintained grown covers with a varying diversity (derived grassland). Better quality areas of White Box – Yellow Box Blakely's Red Gum Grassy Woodland and Derived Native Grasslands were generally present on the edges of the existing easement. The proposal will result in the removal/modification of a relatively small amount (0.59ha) of generally disturbed White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grassland and a small number of trees resulting in an incremental reduction in the local area; however, it is considered unlikely to have a significant impact.

- ***fragment or increase fragmentation of an ecological community, for example by clearing vegetation for roads or transmission lines***

As the proposal will only involve impacts to limited areas of White Box – Yellow Box Blakely's Red Gum Grassy Woodland and Derived Native Grasslands on the edges of an existing easement no areas will be further fragmented.

- ***adversely affect habitat critical to the survival of an ecological community***

The project is unlikely to adversely affect habitat considered to be critical to the survival of the community.

- ***modify or destroy abiotic (non-living) factors (such as water, nutrients, or soil) necessary for an ecological community's survival, including reduction of groundwater levels, or substantial alteration of surface water drainage patterns***

The project is unlikely to affect any abiotic processes necessary for the community's survival.

- ***cause a substantial change in the species composition of an occurrence of an ecological community, including causing a decline or loss of functionally important species, for example through regular burning or flora or fauna harvesting the following criteria:***

The proposal is unlikely to result in substantial change in the species composition of this community locally.

- ***cause a substantial reduction in the quality or integrity of an occurrence of an ecological community, including, but not limited to:***  
-assisting invasive species, that are harmful to the listed ecological community, to become established, or  
– causing regular mobilisation of fertilisers, herbicides or other chemicals or pollutants into the ecological community which kill or inhibit the growth of species in the ecological community, or

Taking the mitigations measures into consideration, the proposal is unlikely to assist any invasive species, nor cause any increased mobilisation of fertilisers which would kill or inhibit the growth of species in the ecological community.

- ***interfere with the recovery of an ecological community.***

The proposal is unlikely to interfere with any current recovery programs in the area.

#### Conclusion

The project is unlikely to significantly impact this community

**Table 9.2: Assessment of EPBC Act White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland. Key diagnostic characteristics against Plant Community Type (PCT)**

| Key diagnostic characteristics as described in in Section 1.5.1 of DoE 2015                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              | Key Diagnostics Met                                                                 |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------|
| The ecological community occurs in the following bioregions (IBRA, DoE 2012): Brigalow Belt South, Murray Darling Depression, Nandewar, New England Tableland, Threatened Species Scientific Committee Page 14 of 121 White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland Conservation Advice NSW North Coast, NSW South Western Slopes, Riverina, South Eastern Queensland, South East Corner, South East Coastal Plain, South Eastern Highlands, Southern Volcanic Plain, Sydney Basin and Victorian Midlands (TSSC 2006; DECCW 2011; DCCEE 2022). | Yes, occurs within the Sydney Basin IBRA Bioregion.                                 |
| It has, or previously had, an overstorey dominated or co-dominated by: <ul style="list-style-type: none"> <li><i>Eucalyptus albens</i> (white box) and/or <i>E. melliodora</i> (yellow box) and/or <i>E. blakelyi</i> (Blakely's red gum) (applicable across the entire range of the ecological community); or,</li> <li>in the Nandewar bioregion (IBRA, DoE 2012), any of the above three species and/or <i>E. microcarpa</i> (western grey box) and/or <i>E. moluccana</i> (grey box, coastal grey box);</li> </ul>                                                                   | Yes, has an overstorey dominated by <i>Eucalyptus blakelyi</i> (Blakely's Red Gum). |
| It has a predominantly native ground layer                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               | Yes                                                                                 |
| Tussock grasses are conspicuous in the ground layer (except in some situations, such as under dense groves of shrubs or regenerating trees), usually with several native species from some the following genera: <i>Austrostipa</i> , <i>Bothriochloa</i> , <i>Chloris</i> , <i>Cymbopogon</i> , <i>Dichanthium</i> , <i>Microlaena</i> , <i>Poa</i> , <i>Themeda</i> , <i>Rytidosperma</i> or <i>Sorghum</i> .                                                                                                                                                                          | Yes                                                                                 |
| Amongst the grass tussocks and sometimes in swathes, a range of broad-leaved forbs and petaloid monocots (e.g. lilies sens. lat.) may be a major component of the plant diversity.                                                                                                                                                                                                                                                                                                                                                                                                       | Yes                                                                                 |
| While shrubs may be dominant locally within areas of the ecological community, areas of native vegetation with a more continuous shrub layer, in which the average shrub cover of the whole patch is greater than 30%, is considered to be a shrubby woodland and so is not part of the listed ecological community. In assessing this, the effects of disturbance need to be considered, for example where heavy grazing may result in high densities of shrubs during a recovery phase                                                                                                 | Yes                                                                                 |
| <b>Does this PCT meet the EPBC Act listing criterion</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 | <b>Yes</b>                                                                          |



### 9.3 KURRI SAND SWAMP WOODLAND OF THE SYDNEY BASIN BIOREGION

To be considered consistent with the Endangered listing under the EPBC Act, the vegetation must be consistent with the criteria outlined in the Approved Conservation Advice for the Kurri sand swamp woodland of the Sydney Basin bioregion (DCCEE, 2022) (Section 2.1) and as summarised in Table 9.3. The assessment concluded that the vegetation did meet the condition criteria of the EPBC Act listing for Kurri sand swamp woodland of the Sydney Basin bioregion. An assessment under the Significant Impact Criteria for this community has been undertaken below.

***An action is likely to have a significant impact on a critically endangered or endangered ecological community if there is a real chance or possibility that it will:***

- ***reduce the extent of an ecological community***

The REZ will impact 0.86ha Kurri Sand Swamp Woodland for approximately 150m of a new section of easement between the Kurri STS and Hunter Expressway (0.2ha) and another section on the other side of the Hunter Expressway (0.66ha). The REZ is will only have a small impact on Kurri Sand Swamp Woodland within the existing easement west of Hart Road. Areas of impacted Kurri Sand Swamp Woodland were connected to much larger areas of this EEC.

- ***fragment or increase fragmentation of an ecological community, for example by clearing vegetation for roads or transmission lines***

No areas are likely to become further fragmented or isolated from other areas that are important to the long-term survival of Kurri Sand Swamp Woodland in the locality.

- ***adversely affect habitat critical to the survival of an ecological community***

The project is unlikely to adversely affect habitat considered to be critical to the survival of the community.

- ***modify or destroy abiotic (non-living) factors (such as water, nutrients, or soil) necessary for an ecological community's survival, including reduction of groundwater levels, or substantial alteration of surface water drainage patterns***

The project is unlikely to affect any abiotic processes necessary for the community's survival.

- ***cause a substantial change in the species composition of an occurrence of an ecological community, including causing a decline or loss of functionally important species, for example through regular burning or flora or fauna harvesting the following criteria:***

The proposal is unlikely to result in substantial change in the species composition of this community locally.

- ***cause a substantial reduction in the quality or integrity of an occurrence of an ecological community, including, but not limited to:***  
 -assisting invasive species, that are harmful to the listed ecological community, to become established, or  
 – causing regular mobilisation of fertilisers, herbicides or other chemicals or pollutants into the ecological community which kill or inhibit the growth of species in the ecological community, or

Taking the mitigations measures into consideration, the proposal is unlikely to assist any invasive species, nor cause any increased mobilisation of fertilisers which would kill or inhibit the growth of species in the ecological community.

**Table 9.3: Assessment of EPBC Act Kurri sand swamp woodland of the Sydney Basin bioregion. Key diagnostic characteristics against Plant Community Type (PCT)**

| Key diagnostic characteristics as described in in Section 2.1 Approved Conservation Advice for the Kurri sand swamp woodland of the Sydney Basin bioregion (DCCEEW, 2022).                                                                                                                                                                                                                                                                                                                                                                                                     | Key Diagnostics Met |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------|
| Mainly occurs within the Hunter subregion of the Sydney Basin Bioregion, with some patches extending into the Wyong subregion (DoE 2013).                                                                                                                                                                                                                                                                                                                                                                                                                                      | Yes                 |
| The vegetation structure is typically a heathland to low sclerophyll woodland.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 | Yes                 |
| Where a eucalypt canopy layer is evident, <i>Angophora bakeri</i> , <i>Eucalyptus parramattensis</i> subsp. <i>decadens</i> and/or <i>Eucalyptus</i> sp. aff. <i>agglomerata</i> are the typical dominant to codominant tree species of the canopy. Other eucalypts, for example <i>E. fibrosa</i> , may also be present and be locally dominant, but not as dominant canopy species throughout a patch. This also applies to hybrids of these trees. In areas where the eucalypt canopy is sparse, <i>Melaleuca nodosa</i> can dominate the mid-storey to tree canopy layers. | Yes                 |
| The mid-layer, where present, is typically characterised by a mid-dense to dense shrub layer of sclerophyllous species, commonly; <i>Acacia elongata</i> , <i>A. longifolia</i> , <i>Banksia collina</i> , <i>Bossiaea</i> spp. <i>Callistemon linearis</i> , <i>Dillwynia retorta</i> , <i>Hakea sericea</i> , <i>Isopogon anemonifolius</i> , <i>Lambertia formosa</i> , <i>Leptospermum</i> spp. <i>Leucopogon virgatus</i> , <i>Melaleuca nodosa</i> , <i>M. thymifolia</i> (thyme honey-myrtle), <i>Monotoca scoparia</i> and <i>Xanthorrhoea glauca</i> (grasstree).     | Yes                 |
| The ground layer is typically a patchy sparse to mid-dense cover of low shrubs and/or grasses, sedges, and forbs, including <i>Anisopogon avenaceus</i> , <i>Aristida</i> spp., <i>Entolasia stricta</i> , and <i>Lomandra</i> spp.                                                                                                                                                                                                                                                                                                                                            | Yes                 |
| <b>Does this PCT meet the EPBC Act listing criterion</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       | <b>Yes</b>          |

- ***interfere with the recovery of an ecological community.***

The proposal is unlikely to interfere with any current recovery programs in the area.

#### Conclusion

The project is unlikely to significantly impact this community

### **9.4 COASTAL SWAMP OAK (CASUARINA GLAUCA) FOREST OF THE NEW SOUTH WALES AND SOUTH-EAST QUEENSLAND ECOLOGICAL COMMUNITY**

To be considered consistent with the Endangered listing under the EPBC Act, the vegetation must be consistent with the criteria outlined in the Approved Conservation Advice for the Coastal Swamp Oak (Casuarina glauca) Forest of New South Wales and South East Queensland ecological community (DCCEEW, 2018) (Section 3.1) and as summarised in Table 9.4. The assessment concluded that the Areas mapped as PCT 4015 Central Hunter Riparian Forest and PCT 4023 Coastal Valley Riparian Forest did meet the condition criteria of the EPBC Act listing for Coastal Swamp Oak (Casuarina glauca) Forest. An assessment under the Significant Impact Criteria for this community has been undertaken below.

An action is likely to have a significant impact on a critically endangered or endangered ecological community if there is a real chance or possibility that it will:

- ***reduce the extent of an ecological community***

The proposal will result in a narrow impact to areas of Coastal Swamp Oak Forest, along drainage lines crossed by the proposed drainage lines.

- ***fragment or increase fragmentation of an ecological community, for example by clearing vegetation for roads or transmission lines***

The project involves the development of an easement for the overhead lines. As the relatively narrow proposal will not impact groundcovers and small shrubs it is unlikely to impede any genetic transfer between areas.

- ***adversely affect habitat critical to the survival of an ecological community***

The project is unlikely to adversely affect habitat considered to be critical to the survival of the community.

- ***modify or destroy abiotic (non-living) factors (such as water, nutrients, or soil) necessary for an ecological community's survival, including reduction of groundwater levels, or substantial alteration of surface water drainage patterns***

The project is unlikely to affect any abiotic processes necessary for the community's survival.



**Table 9.4: Assessment of EPBC Act Coastal Swamp Oak (*Casuarina glauca*) Forest of New South Wales and South East Queensland ecological community. Key diagnostic characteristics against Plant Community Type (PCT)**

| Key diagnostic characteristics as described in in Section 3.1 Approved Conservation Advice for Coastal Swamp Oak ( <i>Casuarina glauca</i> ) Forest (DCCEEW, 2018).                                                                                                                                                                                                            | Key Diagnostics Met<br>PCT 4015 Central Hunter<br>Riparian Forest | Key Diagnostics Met<br>PCT 4023 Coastal Valley<br>Riparian Forest |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------|-------------------------------------------------------------------|
| Occurs from south-east Queensland to southern NSW within the South Eastern Queensland, NSW North Coast, Sydney Basin, or South East Corner bioregions                                                                                                                                                                                                                          | Yes                                                               | Yes                                                               |
| Occurs in coastal catchments at elevations up to 50 m ASL, typically less than 20 m ASL, on coastal flats, floodplains, drainage lines, lake margins, wetlands and estuarine fringes where soils are at least occasionally saturated, water-logged or inundated. There are also minor occurrences on coastal dune swales or flats, particularly deflated dunes and dune soaks. | Yes                                                               | Yes                                                               |
| Occurs on soils derived from unconsolidated sediments (including alluvium), typically hydrosols (grey-black clay-loam and/or sandy loam soils) and sometimes organosols (peaty soils). It may occur in transitional soils (or catenas) where shallow unconsolidated sediments border lithic substrates.                                                                        | Yes                                                               | Yes                                                               |
| Has an open woodland, woodland, forest, or closed forest structure, with a tree canopy that has a total crown cover <sup>1</sup> of at least 10 per cent.                                                                                                                                                                                                                      | Yes                                                               | Yes                                                               |
| Has a canopy of trees dominated by <i>Casuarina glauca</i> (swamp-oak, swamp she-oak).                                                                                                                                                                                                                                                                                         | Yes                                                               | Yes                                                               |
| <b>Does this PCT meet the EPBC Act listing criterion</b>                                                                                                                                                                                                                                                                                                                       | <b>Yes</b>                                                        | <b>Yes</b>                                                        |

- ***cause a substantial change in the species composition of an occurrence of an ecological community, including causing a decline or loss of functionally important species, for example through regular burning or flora or fauna harvesting the following criteria:***

The proposal is unlikely to result in substantial change in the species composition of this community locally.

- **cause a substantial reduction in the quality or integrity of an occurrence of an ecological community, including, but not limited to:**  
 -assisting invasive species, that are harmful to the listed ecological community, to become established, or  
 – causing regular mobilisation of fertilisers, herbicides or other chemicals or pollutants into the ecological community which kill or inhibit the growth of species in the ecological community, or

Taking the mitigations measures into consideration, the proposal is unlikely to assist any invasive species, nor cause any increased mobilisation of fertilisers which would kill or inhibit the growth of species in the ecological community.

- ***interfere with the recovery of an ecological community.***

The proposal is unlikely to interfere with any current recovery programs in the area.

#### Conclusion

The project is unlikely to significantly impact this community

### **9.5 BROWN TREECREEPER**

An action is likely to have a significant impact on a critically endangered or endangered species if there is a real chance or possibility that it will:

- **lead to a long-term decrease in the size of a population**

The proposal resulting in the modification of a relatively narrow area of suitable habitat for the Brown Treecreeper which is connected to substantial areas of habitat is unlikely to lead to any long-term decrease in the size of a population.

- **reduce the area of occupancy of the species**

The removal/modification of habitat is unlikely to reduce the area of occupancy of the mobile Brown Treecreeper.

- **fragment an existing population into two or more populations**

The project will not fragment an existing population of this mobile bird species.

- **adversely affect habitat critical to the survival of a species**

No area critical to the survival of the Brown Treecreeper will be adversely impacted.

- **disrupt the breeding cycle of a population**

The project is unlikely to disrupt any breeding.

- **modify, destroy, remove, isolate or decrease the availability or quality of habitat to the extent that the species is likely to decline**

The proposal is unlikely to remove or modify habitat such that the Brown Treecreeper is likely to decline.

- **result in invasive species that are harmful to a critically endangered or endangered species becoming established in the endangered or critically endangered species' habitat**

The proposal is unlikely to result in the introduction of any invasive species.

- **introduce disease that may cause the species to decline, or**

The proposal is unlikely to result in the introduction of any disease that may affect the Brown Treecreeper.

- **interfere with the recovery of the species.**

The project will not interfere with the recovery of the species.

### Conclusion

The proposal is unlikely to have a significant impact on the Brown Treecreeper as a result of the relatively small area impacted and presence of extensive adjoining habitat.



## **10.0 CONCLUSION**

It is believed that the ecological assessment has been undertaken in accordance with the requirements of the Environmental Planning and Assessment Amendment Act 2017 (EP&A Act 2017), the Biodiversity Conservation Act 2016 (BC Act 2016) and the Commonwealth Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act 1999).

The ecological assessment found the proposed REZ Project and modifications will result in an incremental reduction/modification of habitat, however taking into consideration the mitigation measures the proposal is unlikely to have a significant impact on any addressed threatened species, endangered population or threatened ecological community.

## 11.0 BIBLIOGRAPHY

Churchill, S. (2008). Australian Bats (2nd edn). Allen & Unwin Australia.

Cogger, H.G. (2014). Reptiles and Amphibians of Australia (7th edn.). CSIRO Publishing.

Cropper, S. (1993). *Management of Endangered Plants*. CSIRO Publications, East Melbourne.

DCCEEW (Department of Climate Change, Energy, the Environment and Water) (2022). Approved Conservation Advice for the Kurri sand swamp woodland of the Sydney Basin bioregion.

DCCEEW (Department of Climate Change, Energy, the Environment and Water) (2024a). EPBC Act Protected Matters Report for a 10 Kilometre radius search from the centre of the subject land. Department of the Environment, Commonwealth of Australia. Accessed via, <https://www.environment.gov.au/cgi-bin/sprat/public/sprat.pl> >.

DCCEEW (Department of Climate Change, Energy, the Environment and Water) (2024b). *Commonwealth species profiles and threats database (SPRAT)*. Department of the Environment, Commonwealth of Australia.

Department of Environment, Climate Change and Water NSW. 2010. National Recovery Plan for White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland. Department of Environment, Climate Change and Water NSW, Sydney.

DAWE (Department of Agriculture, Water and the Environment) (2016). Interim Biogeographic Regionalisation for Australia (IBRA), Version 7 (Regions). Accessed via <<https://datasets.seed.nsw.gov.au/dataset/interim-biogeographic-regionalisation-for-australia-ibra-version-7-regions> />

DEH (Department of the Environment and Heritage) (2016). White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grasslands – EPBC Act Policy Statement. Sourced from: <https://www.dcceew.gov.au/sites/default/files/documents/box-gum.pdf>

DoE (Department of the Environment) (2013). *Matters of National Significance. Significant Impact Guidelines 1.1* Environment Protection and Biodiversity Conservation Act 1999.

DoEC (Department of Environment and Conservation (NSW)) (2007). Declaration of critical habitat for the Wollemi Pine (*Wollemia nobilis*)

DPE (Department of Planning and Environment) (2021). White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions- endangered ecological community listing. Sourced from: <https://www.environment.nsw.gov.au/topics/animals-and-plants/threatened-species/nsw-threatened-species-scientific-committee/determinations/final-determinations/2020/white-box-yellow-box-critically-endangered-ecological-community-listing>

DPE (Department of Planning and Environment), (2024a). BioNet Atlas NSW Wildlife Database selected area. (Data extracted 6 F 2023).

DPE (Department of Planning and Environment), (2024b). NSW Biodiversity Values Map accessed via <<https://www.lmbc.nsw.gov.au/Maps/index.html?viewer=BVMap>>.

DPE (Department of Planning and Environment), (2024c) BioNet vegetation classification database. Accessed via < <http://www.environment.nsw.gov.au/research/Visclassification.htm>>

DPE (Department of Planning and Environment), (2024d). Profiles of threatened species, population, and ecological communities. Accessed via <<http://www.environment.nsw.gov.au/threatenedspeciesapp/>>

DPIE (Department of Planning, Industry and Environment), (2011). *Soil and Land Resources of the Hunter Region, version 1.5, NSW Department of Planning, Industry and Environment, Parramatta.*

DPIE (Department of Planning, Industry and Environment) (2017) BioNet NSW (Mitchell) Landscapes – Version 3.1 accessed via < <https://datasets.seed.nsw.gov.au/dataset/nsw-mitchell-landscapes-version-3-1>>.

Ecological Australia (2023). Muswellbrook Solar Farm Biodiversity Development Assessment Report

Harden, G (1991-2000). *Flora of New South Wales*. Vols 1-4. NSW University Press.

Landcom (2004). *Managing Urban Stormwater: Soils and Construction*. National Library of Australia, Canberra, Australia

Hunter Eco (2019). Maxwell Project. Biodiversity Development Assessment Report. Prepared for Malabar Coal.

Leonard, G. (2007). *Eucalypts: A Bushwalker's Guide. Second Edition* New South Wales University Press, Sydney.

LPI (Land and Property Information NSW), (2023). NSW Imagery Web Services.

Mahony, S.V., Cutajar, T. and Rowley, J. J.L. (2022). A new species of *Delma* Gray 1831 (Squamata: Pygopodidae) from the Hunter Valley and Liverpool Plains of New South Wales

Nearmap (2026)

NSW Government (2018) *Australia's IBRA Bioregions and sub-bioregions*. Accessed via <<https://www.seed.nsw.gov.au/>> (Data extracted 1 April 2019).

NSW Government (2024). NSW SEED Mapping. Accessed via <<https://www.seed.nsw.gov.au/>>

NSW DCCEE (NSW Department of Climate Change, Energy, the Environment and Water) (2026a). BioNet Atlas NSW Wildlife Database.

OEH (Office of Environment and Heritage) (2018). *Threatened Species Test of Significance Guidelines*. State of New South Wales and Office of Environment and Heritage 2019.

Pizzey, G. & Knight, F. (2012). *The field guide to the birds of Australia*. (9th edn). HarperCollins Publishers, Australia.

PlantNET (The NSW Plant Information Network System) (2026) *Royal Botanic Gardens and Domain Trust, Sydney*. <http://plantnet.rbgsyd.nsw.gov.au>.

Robinson, L. (2003). *Field Guide to the Native Plants of Sydney*. 3rd edn. Kangaroo Press Pty. Ltd., New South Wales.

Tattersall Lander (2024a). Bushfire Threat Report, Lot 12 in DP 839233, 20-24 Sandy Creek Road Muswellbrook NSW. Ausgrid.

Tattersall Lander (2024b). Bushfire Threat Report, Lot 9 in DP 250890 Hebdon Road, Muswellbrook NSW. Ausgrid. September 2024.

Tattersall Lander (2024c). Bushfire Threat Report, Kurri Kurri. STS Ausgrid.

Triggs, B. (1996). *Mammal Tracks and Signs-A Field Guide for South-eastern Australia*. Oxford University Press, Melbourne.



Umwelt (Australia) Pty Limited (2006), 'Status of the endangered ecological community, Hunter Valley Weeping Myall Woodland', unpublished report to Department of Environment and Conservation, Sydney.

Van Dyck, S. & Strahan, R. (Ed) (2008). *The Mammals of Australia*. 3rd edn. New Holland Publishers, Australia.

Wildthing Environmental Consultants (2024). Biodiversity Development Assessment Report for Stages 6 and 7 of a proposed subdivision at Lot 446 DP 1278895 Northview Estate, Muswellbrook NSW.

WSP (2021). New England Highway – Muswellbrook bypass Biodiversity Assessment Report. August 2021. Transport for NSW.

**APPENDIX A**

**SURVEY POINTS ALONG  
REZ ROUTE**

**Table A1 Survey Point Details**

| Point No. | Location |          | Point Description & Impact                                                                                                                                                                                                                                                                                                                                                                                                                  | Date       | Photo and/or Photo No.  |
|-----------|----------|----------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|-------------------------|
|           | Easting  | Northing |                                                                                                                                                                                                                                                                                                                                                                                                                                             |            |                         |
| 1         | 359498   | 6369179  | From Kurri Kurri STS<br>Maintained lawn around Kurri Kurri STS<br><i>Cynodon dactylon</i> (Couch), <i>Hypochaeris radicata</i> (Flatweed), <i>Richardia humistrata</i> (White Eye)                                                                                                                                                                                                                                                          | 22/08/2024 | #9121                   |
| 2         | 359435   | 6369207  | No existing easement<br>PCT 3631 Kurri Sand-Clay Woodland<br><i>Melaleuca decora</i> (White-feathered Honey-myrtle), <i>Melaleuca nodosa</i> (Ball Honey-myrtle), <i>Acacia longifolia</i> (Sydney Golden Wattle), <i>Grevillea montana</i> .<br>Deer prints on ground.                                                                                                                                                                     | 22/08/2024 | #9135<br>#9134          |
| 3         | 359411   | 6369233  | No existing easement, within drainage line.<br>PCT 3631 Kurri Sand-Clay Woodland<br><i>Eucalyptus parramattensis subsp. decadens</i> (Drooping Red Gum), <i>Melaleuca decora</i> (White-feathered Honey-myrtle), <i>Melaleuca decora</i> (White-feathered Honey-myrtle),<br>Ground: <i>Carex appressa</i> (Saw Sedge), <i>Hypolepis muelleri</i> (Harsh Ground Fern).                                                                       | 22/08/2024 | #9142<br>#9141          |
| 4         | 359324   | 6369289  | Ground: <i>Imperata cylindrica</i> (Blady Grass), <i>Eragrostis curvula</i> (African Lovegrass).                                                                                                                                                                                                                                                                                                                                            | 22/08/2024 | #9179<br>#9180<br>#9181 |
| 5         | 385748   | 6369533  | PCT 3433<br>No new clearing along this section. The new line will be within an already maintained easement.                                                                                                                                                                                                                                                                                                                                 | 29/05/2024 | #8120<br>#8119          |
| 6         | 357765   | 6370033  | <i>Angophora floribunda</i> (Rough-barked Apple), <i>Acacia parvipinnula</i> , <i>Pteridium esculentum</i> (Bracken Fern), <i>Leptospermum polygalifolium</i> (Tantoon), Weeds: <i>Ehrharta erecta</i> (Panic Veldtgrass), <i>Anthoxanthum odoratum</i> (Sweet Vernal Grass),                                                                                                                                                               | 29/05/2024 |                         |
| 7         | 357734   | 6370068  | <i>Acacia parvipinnula</i> , <i>Acacia ulicifolia</i> (Prickly Moses), <i>Pteridium esculentum</i> (Bracken Fern).                                                                                                                                                                                                                                                                                                                          | 29/05/2024 |                         |
| 8         | 357701   | 6370106  | Water crossing <i>Typha orientalis</i> (Cumbungi), <i>Melaleuca sieberi</i> (Sebers Paperback), <i>Hypolepis muelleri</i> (Harsh Ground Fern), <i>Melaleuca nodosa</i> (Ball Honey-myrtle), <i>Imperata cylindrica</i> (Blady Grass), Weeds: <i>Hyparrhenia hirta</i> (Coolatai Grass), <i>Coreopsis lanceolata</i> (Coreopsis).                                                                                                            | 29/05/2024 | #8136                   |
| 9         | 357688   | 6870134  | Scales Road<br>Overhanging branches require removal. <i>Angophora floribunda</i> (Rough-barked Apple), <i>Eucalyptus punctata</i> (Grey Gum), <i>Melaleuca nodosa</i> (Ball Honey-myrtle), <i>Acacia parvipinnula</i> , <i>Angophora bakeri</i> (Narrow-leaved Apple), <i>Bursaria spinosa</i> (Blackthorn). Weeds: <i>Lantana camara</i> (Lantana), <i>Cestrum parqui</i> (Green Cestrum), <i>Asparagus asparagoides</i> (Bridal Creeper). | 29/05/2024 | #8140<br>#8141          |
| 10        | 357630   | 6370185  | Grassy open area. <i>Lomandra longifolia</i> (Spiny Mat Rush), Weeds: <i>Anthoxanthum odoratum</i> (Sweet Vernal Grass), <i>Andropogon virginicus</i> (Whiskey Grass), <i>Sida rhombifolia</i> (Paddys Lucerne), <i>Coreopsis lanceolata</i> (Coreopsis), <i>Ambrosia artemisiifolia</i> (Annual Ragweed),                                                                                                                                  | 29/05/2024 | #8147                   |
| 11        | 357536   | 6370307  | Scales Road<br>Introduced grassy west side of road. Small number of specimens of <i>Eucalyptus tereticornis</i> (Forest Red Gum) may require branch trimming.                                                                                                                                                                                                                                                                               | 29/05/2024 |                         |
| 12        | 357496   | 6370355  | Over Hart Road<br>20m of existing easement already present. Couple of large trees in south may require branch trimming.                                                                                                                                                                                                                                                                                                                     | 29/05/2024 | #8165                   |
| 13        | 357788   | 6370024  | Outside corner Scales Road. <i>Acacia parvipinnula</i> regrowth. Weeds <i>Lantana camara</i> (Lantana), <i>Cenchrus clandestinus</i> (Kikuyu), <i>Ehrharta erecta</i> (Panic Veldtgrass).                                                                                                                                                                                                                                                   | 29/05/2024 | #8166                   |
| 14        | 357834   | 6369979  | Existing easement off bend on Scale Road                                                                                                                                                                                                                                                                                                                                                                                                    | 29/05/2024 | #8168                   |



| Point No. | Location |          | Point Description & Impact                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   | Date       | Photo and/or Photo No.  |
|-----------|----------|----------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|-------------------------|
|           | Eastings | Northing |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |            |                         |
|           |          |          | <i>Cynodon dactylon</i> (Couch), <i>Microlaena stipoides</i> (Weeping Meadow Grass), <i>Cheilanthes sieberi</i> (Mulga Fern), Weeds: <i>Anthoxanthum odoratum</i> (Sweet Vernal Grass). Either side of easement PCT 4039 Hunter Range Creekflat Apple-Red Gum Forest <i>Eucalyptus amplifolia</i> (Cabbage Gum), <i>Pteridium esculentum</i> (Bracken Fern).                                                                                                                                                                                                                                                                                                                                 |            |                         |
| 15        | 357906   | 6369919  | Next to Swamp Creek. <i>Eucalyptus amplifolia</i> (Cabbage Gum), <i>Angophora floribunda</i> (Rough-barked Apple), <i>Lomandra longifolia</i> (Spiny Mat Rush), <i>Phragmites australis</i> (Australian Reed) on edge of creek. Introduced species <i>Ligustrum sinense</i> (Small-leaved Privet), <i>Megathyrsus maximus</i> (Guinea Grass). Only need to remove <i>Acacia parvipinnula</i> on the edge of the easement (likely two trees) Possible branch trimming of <i>Angophora floribunda</i> .                                                                                                                                                                                        | 29/05/2024 | #8173                   |
| 16        | 334160   | 6373     | Bishops Bridge off Old Maitland Road<br>Approximately 3m of additional clearing on edge of clearing.<br><i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus fibrosa</i> (Broad-leaved Ironbark), <i>Eucalyptus moluccana</i> (Grey Box), <i>Melaleuca styphelioides</i> (Prickly-leaved Paperbark).<br>In easement <i>Dodonaea triquetra</i> (Common Hop Bush), <i>Notelaea longifolia</i> (Mock Olive), <i>Breynia oblongifolia</i> (Breynia), <i>Leucopogon juniperinus</i> (Prickly Bearded Heath). <i>Bursaria spinosa</i> (Blackthorn), <i>Macrozamia flexuosa</i> , <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Lobelia purpurescens</i> (Whiteroot). <i>Gahnia aspera</i> . | 29/05/2024 | #8180 to #8185          |
| 17        | 353908   | 6373909  | Three (3) <i>Eucalyptus moluccana</i> (Grey Box) require likely removal. Introduced <i>Chloris gayana</i> (Rhodes Grass) around base of trees.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               | 29/05/2024 | #8196                   |
| 18        | 353837   | 6373944  | <i>Eucalyptus moluccana</i> (Grey Box), <i>Dianella caerulea</i> var. <i>caerulea</i> (Blue Flax Lily). <i>Chloris gayana</i> (Rhodes Grass), <i>Bidens pilosa</i> (Bidens).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 | 29/05/2024 | #8202                   |
| 19        | 353780   | 6373978  | <i>Eucalyptus moluccana</i> (Grey Box), <i>Corymbia maculata</i> (Spotted Gum).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              | 29/05/2024 | #8207<br>#8208          |
| 20        | 353655   | 6374041  | <i>Eucalyptus moluccana</i> (Grey Box), <i>Corymbia maculata</i> (Spotted Gum).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              | 29/05/2024 | #8213                   |
| 21        | 353503   | 6374172  | <i>Acacia parvipinnula</i> , <i>Pultenaea spinosa</i> , <i>Dichondra repens</i> (Kidneyweed). Weeds: <i>Paspalum urvillei</i> (Vasey Grass).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 | 29/05/2024 | #8223<br>#8224          |
| 22        | 353398   | 6374211  | Drainage Crossing: <i>Melaleuca styphelioides</i> (Prickly-leaved Paperbark), <i>Acacia fimbriata</i> , <i>Adiantum aethiopicum</i> (Maidenhair Fern).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       | 29/05/2024 | #8229<br>#8230          |
| 23        | 353222   | 6374317  | Existing Easement. Complies, only minor branch trimming. <i>Eucalyptus moluccana</i> (Grey Box).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             | 29/05/2024 | #8246                   |
| 24        | 351636   | 6375102  | Private Property, Existing creek crossing will need some work, area approx. 5x5m for access for cranes and EWP (heavy vehicle access) Mostly branch removal of up to 2 trees.. PCT 4023 Coastal Valley Riparian Forest – <i>Casuarina glauca</i> (Swamp Oak), <i>Melaleuca styphelioides</i> (Prickly-leaved Paperbark), <i>Alectryon tomentosus</i> (Hairy Bird's Eye), <i>Juncus usitatus</i> (Common Rush), <i>Gahnia aspera</i> , <i>Dichondra repens</i> (Kidneyweed).<br>Fauna Brown Thornbill. <i>Crinia signifera</i> . (Common Eastern Froglet)                                                                                                                                     | 28/05/2024 | #8253<br>#8254<br>#8263 |
| 25        | 351632   | 6375115  | Area of tree to the side of the existing easement require removal. <i>Casuarina glauca</i> (Swamp Oak), <i>Melaleuca styphelioides</i> (Prickly-leaved Paperbark), <i>Bursaria spinosa</i> (Blackthorn). Weeds: <i>Cenchrus clandestinus</i> (Kikuyu).                                                                                                                                                                                                                                                                                                                                                                                                                                       | 28/05/2024 | #8256<br>#8257          |
| 26        | 351453   | 6375216  | Existing 20m easement, only minor branch trimming. PCT 3446 Lower North Foothills Ironbark-Box-Gum Grassy Forest<br><i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Angophora floribunda</i> (Rough-barked Apple), <i>Pultenaea spinosa</i> , <i>Bursaria spinosa</i> (Blackthorn).                                                                                                                                                                                                                                                                                                                                                                                                    | 28/05/2024 | #8267                   |
| 27        | 351324   | 6375282  | Open Grassland. NNW distance, existing easement largely conforms. Only minor branch trimming if required.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | 28/05/2024 | #8268<br>#8269          |
| 28        | 349981   | 6376096  | Creek crossing- PCT 4023 Coastal Valley Riparian Forest <i>Casuarina glauca</i> (Swamp Oak), <i>Melaleuca styphelioides</i> (Prickly-leaved Paperbark), <i>Gahnia aspera</i> , <i>Juncus usitatus</i> (Common Rush), <i>Microlaena stipoides</i> (Weeping Meadow Grass). Weeds <i>Ehrharta erecta</i> (Panic Veldtgrass). <i>Cheilanthes sieberi</i> (Mulga Fern),                                                                                                                                                                                                                                                                                                                           | 28/05/2024 | #8275<br>#8276          |
| 29        | 349965   | 6376149  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              | 28/05/2024 | #8281                   |
| 30        | 349586   | 6376906  | Lovedale                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     | 28/05/2024 | #8282                   |

| Point No. | Location |          | Point Description & Impact                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      | Date       | Photo and/or Photo No.  |
|-----------|----------|----------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|-------------------------|
|           | Eastings | Northing |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |            |                         |
|           |          |          | Possible removal of three (3) trees. <i>Eucalyptus tereticornis</i> (Forest Red Gum).                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |            | #8284                   |
| 31        | 349332   | 6377525  | Lovedale Road, Lovedale.<br>PCT 3315 Central Hunter Ironbark-Spotted Gum Forest Possible removal of three (3) specimens of <i>Eucalyptus moluccana</i> (Grey Box), 1 <i>Eucalyptus fibrosa</i> (Broad-leaved Ironbark). Trees are located in the neighbouring property. Highly altered groundcover. Weeds <i>Hyparrhenia hirta</i> (Coolatai Grass).                                                                                                                                                                                                            | 30/05/2024 | #8288                   |
| 32        | 349329   | 6377547  | Lovedale Road<br>Possible removal of three (3) smaller specimens of <i>Angophora floribunda</i> (Rough-barked Apple). Weeds: <i>Chloris gayana</i> (Rhodes Grass),                                                                                                                                                                                                                                                                                                                                                                                              | 30/05/2024 | #8289                   |
| 33        | 349342   | 6377481  | Lovedale Road<br>Trim overhanging branches. Stringybark.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        | 30/05/2024 | #8290                   |
| 34        | 349342   | 6377481  | Lovedale Road (Branch Trimming)<br>PCT 3315 Central Hunter Ironbark-Spotted Gum Forest <i>Eucalyptus moluccana</i> (Grey Box), <i>Acacia parvipinnula</i> , Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive), <i>Melinis repens</i> (Red Natal Grass), <i>Hyparrhenia hirta</i> (Coolatai Grass).<br>Threatened Grey-crowned Babblers                                                                                                                                                                                                        | 30/05/2024 | #8291                   |
| 35        | 349146   | 6377962  | Existing easement largely compliant. Some overhanging branches likely require removal. PCT 3315 Central Hunter Ironbark-Spotted Gum Forest <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus moluccana</i> (Grey Box), <i>Eucalyptus tereticornis</i> (Forest Red Gum), <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Lomandra multiflora</i> , <i>Eremophila debilis</i> (Amulla). Weeds in easement: <i>Eragrostis curvula</i> (African Lovegrass), <i>Senecio madagascariensis</i> (Fireweed), <i>Bryophyllum delagoense</i> (Mother of Millions). | 30/05/2024 | #8314<br>#8315          |
| 36        | 349105   | 6378007  | Existing easement largely compliant. Branches on eastern side require some trimming.<br>PCT 3315 Central Hunter Ironbark-Spotted Gum Forest <i>Corymbia maculata</i> (Spotted Gum), <i>Pultenaea spinosa</i> , <i>Leucopogon juniperinus</i> (Bearded Heath), <i>Ozothamnus diosmifolius</i> (Dogwood), <i>Daviesia ulicifolia</i> (Gorse Bitter Pea), <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Lomandra multiflora</i> .                                                                                                                            | 30/05/2024 | #3315                   |
| 37        | 348936   | 6378331  | PCT 3315 Central Hunter Ironbark-Spotted Gum Forest – Mainly Derived Grassland. <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus moluccana</i> (Grey Box). <i>Daviesia ulicifolia</i> (Gorse Bitter Pea), <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Themeda triandra</i> (Kangaroo Grass).                                                                                                                                                                                                                                                       | 30/05/2024 | #8326<br>#8237<br>#8238 |
| 38        | 348888   | 6378428  | PCT 3315 Central Hunter Ironbark-Spotted Gum Forest – Derived Grassland. <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Lomandra multiflora</i> , <i>Chrysocephalum apiculatum</i> (Common Everlasting), <i>Themeda triandra</i> (Kangaroo Grass).                                                                                                                                                                                                                                                                                                         | 30/05/2024 | #8330<br>#8331          |
| 39        | 348781   | 6378655  | PCT 3315 Central Hunter Ironbark-Spotted Gum Forest Derived Grassland within existing easement. Side of easement. Minor branch trimming. <i>Eucalyptus moluccana</i> (Grey Box), <i>Acacia implexa</i> (Hickory Wattle), <i>Pultenaea spinosa</i> , <i>Daviesia ulicifolia</i> (Gorse Bitter Pea), <i>Ozothamnus diosmifolius</i> (Dogwood), <i>Leucopogon juniperinus</i> (Prickly Bearded Heath), <i>Calotis cuneifolia</i> (Purple Burr-daisy).                                                                                                              | 30/05/2024 | #8335<br>#8336          |
| 40        | 348834   | 6378547  | Dam within easement. <i>Crinia signifera</i> (Common Eastern Froglet) calling.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  | 30/05/2024 | #8341                   |
| 41        | 348703   | 6378833  | Gate. Some minor branch trimming is required. <i>Eucalyptus moluccana</i> (Grey Box), <i>Eucalyptus tereticornis</i> (Forest Red Gum), <i>Goodenia rotundifolia</i> , <i>Panicum effusum</i> , <i>Aristida vagans</i> .                                                                                                                                                                                                                                                                                                                                         | 30/05/2024 | #8349<br>#8350          |
| 42        | 348613   | 6379021  | Existing easement largely compliant. Minor branch trimming. <i>Eucalyptus moluccana</i> (Grey Box), <i>Pultenaea spinosa</i> , <i>Chrysocephalum apiculatum</i> (Common Everlasting), <i>Rytidosperma</i> sp. (Wallaby Grass), <i>Hardenbergia violacea</i> (Purple Coral Pea), <i>Hypericum gramineum</i> (Native St John's Wort).                                                                                                                                                                                                                             | 30/05/2024 | #8352<br>#8353          |
| 43        | 348524   | 6379212  | <i>Eucalyptus moluccana</i> (Grey Box),                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         | 30/05/2024 | #8354<br>#8355          |

| Point No. | Location |           | Point Description & Impact                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | Date       | Photo and/or Photo No.  |
|-----------|----------|-----------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|-------------------------|
|           | Easting  | Northing  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |            |                         |
| 44        | 348475   | 6379458   | <i>Eucalyptus tereticornis</i> (Forest Red Gum), <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Ozothamnus diosmifolius</i> (Dogwood), <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Cynodon dactylon</i> (Couch). Weeds: <i>Bryophyllum delagoense</i> (Mother of Millions), <i>Paspalum dilatatum</i> (Paspalum).                                                                                                                                                                                                              | 30/05/2024 | #8365<br>#8366          |
| 45        | 348470   | 6379591   | Near Camp Road<br>Smaller Specimens of <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Angophora floribunda</i> (Rough-barked Apple), <i>Melaleuca styphelioides</i> (Prickly-leaved Paperbark), <i>Cynodon dactylon</i> (Couch). Weeds: <i>Senecio madagascariensis</i> (Fireweed).                                                                                                                                                                                                                                                    | 30/05/2024 | #8368<br>#8369          |
| 49        | 348419   | 6379950   | <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Pultenaea spinosa</i> , <i>Acacia longifolia</i> (Sydney Golden Wattle), <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Calotis lappulacea</i> (Yellow Burr-daisy).                                                                                                                                                                                                                                                                                                                | 30/05/2024 | #8383<br>#8384          |
| 50        | 348404   | 6380122   | Minor branch trimming. <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Pultenaea spinosa</i> , <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Glycine tabacina</i> . Weeds: <i>Senecio madagascariensis</i> (Fireweed).                                                                                                                                                                                                                                                                    | 30/05/2024 | #8385<br>#8386          |
| 51        | 348363   | 6380354   | Branch Trimming. Smaller trees may require removal. Minor branch trimming. <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus moluccana</i> (Grey Box), <i>Daviesia ulicifolia</i> (Gorse Bitter Pea), <i>Pultenaea spinosa</i> , <i>Ozothamnus diosmifolius</i> (Dogwood), <i>Acacia elongata</i> . Weeds: <i>Bryophyllum delagoense</i> (Mother of Millions), <i>Senecio madagascariensis</i> (Fireweed).                                                                             | 30/05/2024 | #8389<br>#8390          |
| 52        | 348290   | 6380600   | Minor branch trimming. Possible removal of small trees.<br><i>Eucalyptus moluccana</i> (Grey Box), <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Breynia oblongifolia</i> (Breynia), <i>Pultenaea spinosa</i> , <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Gahnia aspera</i> , <i>Aristida racemosa</i> (Three-awn Grass). Weeds: <i>Bryophyllum delagoense</i> (Mother of Millions), <i>Senecio madagascariensis</i> (Fireweed), <i>Lantana camara</i> (Lantana), <i>Axonopus fissifolius</i> (Narrow-leaved Carpet Grass). | 30/05/2024 | #8393<br>#8394          |
| 53        | 348220   | 6380938   | Possible removal of smaller trees. Minor branch trimming.<br><i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus fibrosa</i> (Broad-leaved Ironbark), <i>Pultenaea spinosa</i> , <i>Acacia parvipinnula</i> , <i>Aristida racemosa</i> , <i>Lomandra multiflora</i> .                                                                                                                                                                                                                                                                       | 30/05/2024 | #8397<br>#8398          |
| 54        | 348177   | 6381150   | <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus fibrosa</i> (Broad-leaved Ironbark), <i>Pultenaea spinosa</i> , <i>Acacia parvipinnula</i> , <i>Bursaria spinosa</i> (Blackthorn), <i>Aristida racemosa</i> , <i>Goodenia rotundiflora</i> , <i>Lomandra multiflora</i> .<br>Grape vines in vineyard to the east.                                                                                                                                                                                                                       | 30/05/2024 | #8400<br>#8401          |
| 55        | 348124   | 6381592   | Derived grassland between fence and highway. Trees <i>Eucalyptus tereticornis</i> (Forest Red Gum), <i>Lomandra longifolia</i> , <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Lomandra multiflora</i> . <i>Chryscephalum apiculatum</i> . Weeds: <i>Senecio madagascariensis</i> (Fireweed), <i>Setaria parviflora</i> (Slender Pigeon Grass).                                                                                                                                                                                         | 30/05/2024 | #8402<br>#8403          |
| 56        | 348048   | 6381594   | Maintained Lawn, Paddock.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     | 30/05/2024 | #8403<br>#8404          |
| 57        | 347617   | 6381634   | Tuckers Lane. Branch trimming on the corner. <i>Corymbia maculata</i> (Spotted Gum), Young <i>Eucalyptus fibrosa</i> off tuckers road to the north. <i>Acacia podalyriifolia</i> (Qld Silver Wattle), <i>Melinus repens</i> (Red Natal Grass).                                                                                                                                                                                                                                                                                                | 30/05/2024 | #8406<br>#8407          |
| 58        | 347603   | 6381823   | Young regrowth, <i>Eucalyptus fibrosa</i> . <i>Acacia podalyriifolia</i> (Qld Silver Wattle) not much on the ground.                                                                                                                                                                                                                                                                                                                                                                                                                          | 30/05/2024 | #8412<br>#84            |
| 59        | 347582   | 638198763 | Next to dam. <i>Acacia parvipinnula</i> , <i>Hakea sericea</i> (Needlebush), <i>Acacia falcata</i> (Falcate Wattle).<br><i>Crinia signifera</i> (Common Eastern Froglet) call from dam. Birds Jacky Winter, Willy Wagtail.                                                                                                                                                                                                                                                                                                                    | 30/05/2024 | #8429<br>#8430<br>#8431 |
| 60        | 347583   | 6382141   | <i>Acacia longifolia</i> (Sydney Golden Wattle) – dense area next to drainage line.<br>Brown Thornbill, Fuscus Honeyeater, Grey Fantail.                                                                                                                                                                                                                                                                                                                                                                                                      | 30/05/2024 | #8433                   |
| 61        | 347594   | 6382214   | Highly disturbed area. <i>Acacia parvipinnula</i> , <i>Breynia oblongifolia</i> , <i>Cynodon dactylon</i> (Couch), <i>Lantana camara</i> (Lantana).<br><i>Carlia tetradactyla</i> (Southern Rainbow Skink).                                                                                                                                                                                                                                                                                                                                   | 30/05/2024 | #8440<br>#8441          |



| Point No. | Location |          | Point Description & Impact                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 | Date       | Photo and/or Photo No.  |
|-----------|----------|----------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|-------------------------|
|           | Easting  | Northing |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |            |                         |
| 62        | 347622   | 6382392  | PCT 3315 Central Hunter Ironbark-Spotted Gum Forest, <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Eucalyptus punctata</i> (Grey Gum), <i>Eucalyptus tereticornis</i> (Forest Red Gum), <i>Melaleuca nodosa</i> (Ball Honey Myrtle), <i>Acacia ulicifolia</i> , <i>Persoonia linearis</i> (Narrow-leaved Geebung), <i>Breynia oblongifolia</i> , <i>Acacia parvipinnula</i> , <i>Grevillea montana</i> , <i>Dianella revoluta</i> , <i>Chrysocephalum apiculatum</i> , <i>Aristida racemosa</i> , <i>Pultenaea spinosa</i> .                                                                                                                                                                                                       | 30/05/2024 | #8445<br>#8446          |
| 63        | 347662   | 6382656  | PCT 3315 Central Hunter Ironbark-Spotted Gum Forest, Top of hill, Road off to the side. <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Corymbia maculata</i> (Spotted Gum), <i>Melaleuca nodosa</i> (Ball Honey Myrtle), <i>Persoonia linearis</i> (Narrow-leaved Geebung), <i>Breynia oblongifolia</i> , <i>Hakea sericea</i> (Needlebush), <i>Acacia longifolia</i> (Sydney Golden Wattle), <i>Grevillea montana</i> , <i>Lepidosperma laterale</i> , <i>Acacia parvipinnula</i> , <i>Lomandra multiflora</i> , <i>Phyllanthus hirtellus</i> , <i>Aristida vagans</i> , <i>Dianella revoluta</i> , <i>Chrysocephalum apiculatum</i> , <i>Cymbopogon refractus</i> (Barbed Wire Grass). Rainbow Lorikeet, Yellow-faced Honeyeater. | 03/06/2024 | #8448<br>#8449          |
| 64        | 347693   | 6382875  | PCT 3315 Central Hunter Ironbark-Spotted Gum Forest, Similar vegetation to P63. <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Corymbia maculata</i> (Spotted Gum), <i>Styphandra glauca</i> , <i>Melichrus urceolatus</i>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          | 03/06/2024 | #8457<br>#8458          |
| 65        | 347676   | 6383125  | PCT 3315 Central Hunter Ironbark-Spotted Gum Forest, <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus fibrosa</i> , <i>Melaleuca nodosa</i> (Ball Honey Myrtle), <i>Melaleuca decora</i> , <i>Persoonia linearis</i> (Narrow-leaved Geebung), <i>Acacia longifolia</i> (Sydney Golden Wattle), <i>Grevillea montana</i> .                                                                                                                                                                                                                                                                                                                                                          | 03/06/2024 | #8461<br>#8462          |
| 66        | 347614   | 6383258  | <i>Eucalyptus fibrosa</i> , <i>Corymbia maculata</i> (Spotted Gum), <i>Persoonia linearis</i> (Narrow-leaved Geebung), <i>Grevillea montana</i> , <i>Phyllanthus hirtellus</i> , <i>Daviesia ulicifolia</i> (Gorse Bitter Pea), <i>Acacia falcata</i> , <i>Aristida racemosa</i> .                                                                                                                                                                                                                                                                                                                                                                                                                                                         | 03/06/2024 | #8465<br>#8466          |
| 67        | 347458   | 6383449  | <i>Eucalyptus fibrosa</i> (Dominant), Some <i>Eucalyptus tereticornis</i> (Forest Red Gum), <i>Breynia oblongifolia</i> , <i>Aristida racemosa</i> , <i>Cymbopogon refractus</i> (Barbed Wire Grass). Intact Grey-crowned Babbler Nest in skinny Spotted Gum.                                                                                                                                                                                                                                                                                                                                                                                                                                                                              | 03/06/2024 | #8476<br>#8477          |
| 68        | 347458   | 6383449  | PCT 3328 Lower Hunter Red Gum-Paperbark Riverflat Forest - Creek Crossing, <i>Eucalyptus tereticornis</i> (Forest Red Gum), <i>Melaleuca decora</i> .                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      | 03/06/2024 | #8488                   |
| 69        | 347188   | 6383662  | Possible removal of smaller trees and branch trimming of larger trees. <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Eucalyptus punctata</i> (Grey Gum), <i>Melaleuca decora</i> , <i>Persoonia linearis</i> (Narrow-leaved Geebung), <i>Phyllanthus hirtellus</i> , <i>Entolasia stricta</i> (Wiry Panic), <i>Lomandra multiflora</i> , <i>Lobelia purpurea</i> , <i>Breynia oblongifolia</i> , <i>Dianella revoluta</i> .                                                                                                                                                                                                                                                                | 03/06/2024 | #8489<br>#8490<br>#8491 |
| 70        | 346965   | 6383829  | Creek crossing- PCT 4023 Coastal Valley Riparian Forest Casuarina glauca (Swamp Oak), <i>Eucalyptus tereticornis</i> (Forest Red Gum), <i>Melaleuca linariifolia</i> , <i>Microlaena stipoides</i> (Weeping Meadow Grass), Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive), <i>Asparagus asparagoides</i> (Bridal Creeper), <i>Juncus acutus</i> . Satin Bowerbird.                                                                                                                                                                                                                                                                                                                                                    | 03/06/2024 | #8496<br>#8497          |
| 71        | 346649   | 6384037  | <i>Eucalyptus tereticornis</i> (Forest Red Gum), <i>Eucalyptus punctata</i> (Grey Gum), <i>Melaleuca decora</i> , <i>Breynia oblongifolia</i> , <i>Persoonia linearis</i> (Narrow-leaved Geebung), <i>Acacia falcata</i> . Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive), <i>Grevillea montana</i> .                                                                                                                                                                                                                                                                                                                                                                                                                 | 03/06/2024 | #8514<br>#8515<br>#8516 |
| 72        | 346649   | 6384037  | <i>Eucalyptus tereticornis</i> (Forest Red Gum) to east <i>Eucalyptus punctata</i> (Grey Gum).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             | 03/06/2024 | #8519<br>#8520          |
| 73        | 346488   | 6384103  | Easement is relatively wide – only minor branch trimming. <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Acacia parvipinnula</i> , <i>Melaleuca nodosa</i> (Ball Honey Myrtle).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             | 03/06/2024 | #8523<br>#8524          |
| 74        | 346323   | 6384150  | <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Acacia parvipinnula</i> , <i>Melaleuca nodosa</i> (Ball Honey Myrtle), <i>Aristida racemosa</i> , <i>Pultenaea spinosa</i> , <i>Grevillea montana</i> , <i>Microlaena stipoides</i> (Weeping Meadow Grass), <i>Cheilanthes sieberi</i> (Mulga Fern), <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Bursaria spinosa</i> (Blackthorn), <i>Goodenia rotundifolia</i> .                                                                                                                                                                                                                                                                   | 03/06/2024 | #8525<br>#8526<br>#8527 |
| 75        | 346172   | 6384150  | Thick <i>Melaleuca nodosa</i> (Ball Honey Myrtle), <i>Eucalyptus fibrosa</i> , <i>Grevillea montana</i> , <i>Aristida racemosa</i> .                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       | 03/06/2024 | #8531                   |

| Point No. | Location |          | Point Description & Impact                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             | Date       | Photo and/or Photo No.  |
|-----------|----------|----------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|-------------------------|
|           | Easting  | Northing |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |            |                         |
|           |          |          |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |            | #8532                   |
| 76        | 346030   | 6384197  | Disturbed cleared area <i>Cynodon dactylon</i> (Couch), Weeds <i>Ambrosia artemisiifolia</i> (Annual Ragweed), <i>Bidens pilosa</i> (Bidens), <i>Eragrostis curvula</i> (African Lovegrass), <i>Sida rhombifolia</i> (Paddys Lucerne)                                                                                                                                                                                                                                                                                  | 03/06/2024 | #8536<br>#8537          |
| 77        | 346040   | 6384285  | Area between Road and Expressway – Disturbed easement. Only minor branch trimming.<br><i>Eucalyptus fibrosa</i> , <i>Eucalyptus moluccana</i> (Grey Box), <i>Acacia parvipinnula</i> , <i>Acacia falcata</i> , <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Dichodra repens</i> (Kidneyweed), <i>Cynodon dactylon</i> . Weeds: <i>Lycium ferocissimum</i> (African Boxthorn), <i>Chloris gayana</i> (Rhodes Grass).                                                                                             | 03/06/2024 |                         |
| 78        | 346070   | 6384505  | Corner bend in powerline. Sides of existing easement <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus fibrosa</i> , <i>Melaleuca decora</i> , <i>Grevillea montana</i> , <i>Cynodon dactylon</i> , <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Microlaena stipoides</i> (Weeping Meadow Grass). Weeds common within easement. <i>Paspalum urvillei</i> (Vasey Grass), <i>Melinis repens</i> (Red Natal Grass), <i>Bryophyllum delagoense</i> (Mother-of-millions), <i>Opuntia stricta</i> (Prickly Pear). | 03/06/2024 | #8547<br>#8548          |
| 79        | 345778   | 6384607  | Drainage line 15m to west.<br>Weedy grasses within easement, <i>Cynodon dactylon</i> , <i>Capillipedium parviflorum</i> (Scented Top Grass). <i>Casuarina glauca</i> (Swamp Oak) either side of easement.<br>Mistletoe Bird.                                                                                                                                                                                                                                                                                           | 03/06/2024 | #8566<br>#8567          |
| 80        | 345601   | 6384666  | Highly disturbed easement. <i>Eucalyptus fibrosa</i> to side. Only minor branch trimming required. Weeds <i>Verbena bonariensis</i> (Purple Top Verbena), <i>Setaria parviflora</i> .                                                                                                                                                                                                                                                                                                                                  | 03/06/2024 | #8568<br>#8569<br>#8570 |
| 81        | 345305   | 6384761  | On northern side of easement, slightly west likely removal of some trees either side<br><i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus fibrosa</i> , <i>Acacia parvipinnula</i> , <i>Leucopogon juniperinus</i> . Weeds <i>Melinis repens</i> (Red Natal Grass), <i>Eragrostis curvula</i> (African Lovegrass).                                                                                                                                                                                                 | 03/06/2024 | #8573<br>#8574          |
| 82        | 345010   | 6384850  | Some tree removal to the east.<br>Disturbed introduced grassy easement. <i>Eucalyptus fibrosa</i> , <i>Corymbia maculata</i> (Spotted Gum), <i>Acacia parvipinnula</i> , <i>Acacia falcata</i> , <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive).<br>Grey-crowned Babbler Nest                                                                                                                                                                                                                            | 03/06/2024 | #8579<br>#8580          |
| 83        | 344729   | 6384944  | Near Branxton Railway Station<br>Highly disturbed easement. <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark) small number of trees require removal approx. 50m to the west.                                                                                                                                                                                                                                                                                                   | 03/06/2024 | #8591<br>#8592          |
| 84        | 344429   | 6384981  | <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Acacia falcata</i> . Small number of smaller trees require removal & some branch trimming. <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive), <i>Melaleuca decora</i> , <i>Dianella revoluta</i> .                                                                                                                                                                                                            | 03/06/2024 | #8597<br>#8598          |
| 85        | 344227   | 6385061  | <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus moluccana</i> (Grey Box), <i>Melaleuca decora</i> , <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark). <i>Lomandra longifolia</i> , <i>Grevillea montana</i> , <i>Aristida vagans</i> , <i>Lomandra multiflora</i> , <i>Breynia oblongifolia</i> .<br>Two Grey-crowned Babbler Nests.                                                                                                                                                                            | 03/06/2024 | #8603<br>#8604          |
| 86        | 343968   | 6385126  | Branch trimming along section.<br><i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus moluccana</i> (Grey Box), <i>Acacia falcata</i> , <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Cynodon dactylon</i> , <i>Lomandra multiflora</i> , <i>Aristida racemosa</i> , <i>Lomandra longifolia</i>                                                                                                                                                                                                                | 03/06/2024 | #8607<br>#8608          |
| 88        | 343848   | 6385150  | <i>Corymbia maculata</i> (Spotted Gum), <i>Acacia falcata</i> , <i>Grevillea montana</i> .                                                                                                                                                                                                                                                                                                                                                                                                                             | 03/06/2024 | #8614<br>#8615          |
| 89        | 343787   | 6385148  | Old railway to west (cuts through). <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Acacia falcata</i> .<br>Minor branch trimming.                                                                                                                                                                                                                                                                                                                                       | 03/06/2024 | #8617<br>#8618          |
| 90        | 343484   | 6385160  | Cleared Grassy Area, <i>Aristida racemosa</i> , <i>Cynodon dactylon</i> , <i>Cymbopogon refractus</i> (Barbed Wire Grass). Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive),                                                                                                                                                                                                                                                                                                                        | 03/06/2024 | #8619<br>#8620          |

| Point No. | Location |          | Point Description & Impact                                                                                                                                                                                                                                                                           | Date       | Photo and/or Photo No.  |
|-----------|----------|----------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|-------------------------|
|           | Easting  | Northing |                                                                                                                                                                                                                                                                                                      |            |                         |
| 91        | 343268   | 6385169  | <i>Casuarina glauca</i> (Swamp Oak), <i>Carex appressa</i> , <i>Cynodon dactylon</i> , Weeds: <i>Cardiospermum grandiflorum</i> (Balloon Vine).<br>Welcome Swallow, Eastern Grey Kangaroos.                                                                                                          | 03/06/2024 | #8624<br>#8625          |
| 94        | 342342   | 6385161  | Lower grassed paddock (grasses). Area of <i>Eucalyptus tereticornis</i> (Forest Red Gum) to NE. <i>Cynodon dactylon</i> , <i>Senecio madagascariensis</i> (Fireweed).                                                                                                                                | 4/06/2024  | #8652<br>#8653          |
| 95        | 342065   | 6385108  | Low Paddock Grasses. Trees to north, <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Eucalyptus moluccana</i> (Grey Box).<br>Black-faced Cuckoo Shrike, Noisy Miner.                                                                                                                           | 4/06/2024  | #8659<br>#8660          |
| 96        | 341882   | 6385075  | Grassy Paddock. <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Eucalyptus moluccana</i> (Grey Box). <i>Cynodon dactylon</i> .<br>Minor Branch trimming.                                                                                                                                       | 4/06/2024  | #8662<br>#8663          |
| 97        | 341787   | 6385057  | New Green Tank to North. <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark) near tank.<br>Noisy Miner                                                                                                                                                                                                 | 4/06/2024  | #8668<br>#8669          |
| 98        | 341520   | 6385056  | Grassy Paddock low grasses- <i>Cynodon dactylon</i> , <i>Mentha satuireioides</i> (Native Mint), <i>Senecio madagascariensis</i> (Fireweed). Two smaller trees to NW both <i>Eucalyptus fibrosa</i> .                                                                                                | 4/06/2024  | #8668<br>#8669          |
| 99        | 341224   | 6385052  | Grassy Paddock low grasses- <i>Cynodon dactylon</i> , <i>Senecio madagascariensis</i> (Fireweed).<br>Two smaller trees to NW both <i>Eucalyptus fibrosa</i> likely to require trimming. <i>Allocasuarina luehmannii</i> (Bulloak) nearby.                                                            | 4/06/2024  | #8673<br>#8674          |
| 113       | 338837   | 6385659  | <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Eucalyptus sideroxylon</i> (Mugga Ironbark) (this Ironbark species may not be endemic to the area).                                                                                                    | 4/06/2024  | #8770<br>#8771          |
| 114       | 338771   | 6385680  | Along New England Highway, Belford.<br><i>Eucalyptus moluccana</i> (Grey Box), <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus tereticornis</i> (Forest Red Gum).<br>Quite weedy within easement – Introduced grasses <i>Megathyrsus maximus</i> (Guinea Grass).<br>Potential branch trimming. | 4/06/2024  | #8774<br>#8775          |
| 115       | 338464   | 6385724  | Along New England Highway, Belford.<br><i>Eucalyptus punctata</i> (Grey Gum), <i>Corymbia maculata</i> (Spotted Gum). Introduced grasses in easement <i>Chloris gayana</i> (Rhodes Grass), <i>Hyparrhenia hirta</i> (Coolatai Grass), <i>Gomphocarpus fruticosus</i> .                               | 4/06/2024  | #8785<br>#8786          |
| 116       | 338250   | 6385759  | Next to intersection of New England Highway and Hermitage Road.<br><i>Corymbia maculata</i> (Spotted Gum). Introduced ground covers <i>Chloris gayana</i> (Rhodes Grass), <i>Galium aparine</i> .                                                                                                    | 4/06/2024  | #8792<br>#8793          |
| 117       | 338149   | 6385776  | Minor drainage line.<br>Native: <i>Phragmites australis</i> (Australian Reed). Weeds <i>Chloris gayana</i> (Rhodes Grass).<br><i>Crinia signifera</i> (Common Eastern Froglet) heard calling.                                                                                                        | 4/06/2024  | #8799<br>#8800          |
| 118       | 338021   | 6385801  | Along New England Highway, Belford just east of <b>Jump up Creek</b> .<br>PCT 4023 Coastal Valley Riparian Forest. <i>Casuarina glauca</i> (Swamp Oak), <i>Phragmites australis</i> (Australian Reed) in creek. Specimen of <i>Eucalyptus glaucina</i> (Slaty Red Gum) to south outside easement.    | 4/06/2024  | #8805<br>#8807<br>#8808 |
| 119       | 337776   | 6385838  | Along New England Highway, Belford.<br>Weeds - <i>Megathyrsus maximus</i> (Panic) and <i>Cenchrus clandestinus</i> (Kikuyu).                                                                                                                                                                         | 4/06/2024  | #8815<br>#8816          |
| 120       | 337581   | 6385838  | Along Belford Road.<br><i>Capillipedium spicigerum</i> (Scented Top Grass).                                                                                                                                                                                                                          | 4/06/2024  | #8822<br>#8823          |
| 121       | 337322   | 6385799  | Three <i>Eucalyptus sideroxylon</i> (Mugga) trees.<br>Likely require branch trimming. Pied Currawong.                                                                                                                                                                                                | 4/06/2024  | #8828<br>#8829          |
| 122       | 337220   | 6385788  | Pole to be move slightly north will result in the removal of three specimens of <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark) (two of these trees are small).                                                                                                                                    | 4/06/2024  | #8836<br>#8837          |
| 123       | 336909   | 6385821  | Natives: <i>Bothriochloa decipiens</i> (Red Grass), <i>Cynodon dactylon</i> (Couch). Weeds: <i>Plantago lanceolata</i> (Plantain), <i>Senecio madagascariensis</i> (Fireweed).                                                                                                                       | 4/06/2024  | #8844<br>#8845          |



| Point No. | Location |          | Point Description & Impact                                                                                                                                                                                                                                                                                     | Date      | Photo and/or Photo No.  |
|-----------|----------|----------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------|-------------------------|
|           | Easting  | Northing |                                                                                                                                                                                                                                                                                                                |           |                         |
| 124       | 336705   | 6385855  | <i>Angophora floribunda</i> (Rough-barked Apple) trees on dam to the east (may require removal). <i>Eucalyptus tereticornis</i> (Forest Red Gum) to west may require branch trimming. Paddock grasses.                                                                                                         | 4/06/2024 | #8851                   |
| 125       | 336377   | 6385912  | Top of hill near Telstra Tower. Olive Trees 100m to the east. <i>Bothriochloa decipiens</i> (Red Grass), <i>Cynodon dactylon</i> (Couch). Weeds: <i>Senecio madagascariensis</i> (Fireweed), <i>Tagetes minuta</i> (Stinking Roger).                                                                           | 4/06/2024 | #8858<br>#8859          |
| 126       | 336087   | 6386003  | Paddock grasses, <i>Cynodon dactylon</i> (Couch). Weeds: <i>Lysimachia arvensis</i> (Scarlet Pimpernel).                                                                                                                                                                                                       | 4/06/2024 | #8863<br>#8864          |
| 127       | 335859   | 6386076  | Clear area, composed of paddock grasses. <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark) trees to side may require trimming.                                                                                                                                                                                 | 4/06/2024 | #8665<br>#8666          |
| 128       | 335680   | 6386164  | Natives: <i>Corymbia maculata</i> (Spotted Gum) (close trees) potential branch trimming. Weeds: <i>Senecio madagascariensis</i> (Fireweed), <i>Gomphocarpus fruticosus</i> .                                                                                                                                   | 4/06/2024 | #8868<br>#8869          |
| 129       | 335460   | 6386276  | Fence just to west. Nearby trees to NE <i>Allocasuarina luehmannii</i> (Bullock) and NW <i>Casuarina glauca</i> (Swamp Oak). Large <i>Corymbia maculata</i> (Spotted Gum) to the south.                                                                                                                        | 4/06/2024 | #8872<br>#8873          |
| 130       | 335282   | 6386367  | Within crop of <i>Avena</i> sp. (Oats). Drainage line 120m to the east. Small specimens of <i>Casuarina glauca</i> (Swamp Oak) may require removal.                                                                                                                                                            | 4/06/2024 | #8875<br>#8876          |
| 131       | 334982   | 6386514  | Within crop of <i>Avena</i> sp. (Oats). <i>Eucalyptus moluccana</i> (Grey Box) to SW likely just outside impact area. Weeds: <i>Malva parviflora</i> (Small-flowered Mallow).                                                                                                                                  | 4/06/2024 | #8878<br>#8879          |
| 132       | 334564   | 6386720  | Just east of drainage line. <i>Juncus usitatus</i> (Common Reed), <i>Carex appressa</i> (Saw Sedge), <i>Eleocharis</i> sp. Weeds: <i>Rumex crispus</i> (Curled Dock).                                                                                                                                          | 4/06/2024 | #8882<br>#8883          |
| 133       | 334303   | 6386848  | Near Golden Highway Intersection with New England Highway. Road reserve, <i>Eucalyptus tereticornis</i> (Forest Red Gum), <i>E. crebra</i> , <i>E. fibrosa</i> . Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive), <i>Ricinus communis</i> (Caster Oil Plant). <b>Grey-crowned Babblers</b> | 4/06/2024 | #8886<br>#8887          |
| 138       | 333429   | 6387071  | Entry to private property. Row of planted <i>Corymbia maculata</i> (Spotted Gum), Planted native trees just inside boundary to the north. <i>Eucalyptus tereticornis</i> (Forest Red Gum) to SW in road reserve. : <i>Chloris gayana</i> (Rhodes Grass),                                                       | 6/06/2024 | #9087<br>#9088          |
| 139       | 333224   | 6387102  | Old Mill Gate. <i>Eucalyptus</i> cut back, appearing to lean to the north. Likely specimen of <i>Eucalyptus glauca</i> (Slaty Red Gum) in paddock to the NW.                                                                                                                                                   | 6/06/2024 | #9091<br>#9092          |
| 140       | 333152   | 6387111  | <i>Eucalyptus crebra</i> & <i>E. moluccana</i> within property to the north. Branch trimming maybe required.                                                                                                                                                                                                   | 6/06/2024 | #9093<br>#9094          |
| 141       | 332986   | 6387135  | <i>Allocasuarina luehmannii</i> (Bullock) just over fence & <i>Angophora floribunda</i> may require removal or trimming.                                                                                                                                                                                       | 6/06/2024 | #9095<br>#9096          |
| 142       | 332746   | 6387161  | <i>Crinia signifera</i> (Common Eastern Froglet) heard calling from the creek.                                                                                                                                                                                                                                 | 6/06/2024 | #9098<br>#9099          |
| 143       | 332362   | 6387213  | No impact disturbed grassland.                                                                                                                                                                                                                                                                                 | 6/06/2024 | #9100<br>#9101          |
| 144       | 332074   | 6387259  | Horse head on gate. Introduced species: <i>Chloris gayana</i> (Rhodes Grass), <i>Rumex crispus</i> (Curled Dock), <i>Galium aparine</i> (Clevers).                                                                                                                                                             | 6/06/2024 | #9102<br>#9103          |
| 145       | 331967   | 6387284  | Just East of Muds Creek. <i>Casuarina glauca</i> (Swamp Oak), <i>Clematis</i> sp. <i>Austrostipa</i> sp.. <i>Phragmites australis</i> in creek. PCT 4015. Introduced: <i>Anredera cordifolia</i> (Madeira Vine) quite invasive.                                                                                | 6/06/2024 | #9105<br>#9106<br>#9107 |
| 146       | 331909   | 6387293  | <i>Phragmites australis</i> : Introduced <i>Galium aparine</i> (Clevers).                                                                                                                                                                                                                                      | 6/06/2024 | #9112<br>#9113          |
| 147       | 331801   | 6387306  | Few <i>Casuarina glauca</i> (Swamp Oaks) to east. Disturbed roadside grasses to the west.                                                                                                                                                                                                                      | 6/06/2024 |                         |
| 148       | 331068   | 6387375  | At gate-ED lodge. Tree 100m to east may require trimming. Planted trees to west in private property including <i>Grevillea robusta</i> , <i>Callistemon salignus</i> and <i>Melaleuca styphelioides</i> .                                                                                                      | 6/06/2024 | #9116<br>#9117          |
| 149       | 330865   | 6387385  | Planned trees back to east. Disturbed roadside grasses to west - <i>Chloris gayana</i> (Rhodes Grass).                                                                                                                                                                                                         | 6/06/2024 | #9118                   |

| Point No. | Location |          | Point Description & Impact                                                                                                                                                                                                                                                              | Date      | Photo and/or Photo No.           |
|-----------|----------|----------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------|----------------------------------|
|           | Easting  | Northing |                                                                                                                                                                                                                                                                                         |           |                                  |
|           |          |          |                                                                                                                                                                                                                                                                                         |           | #9119<br>#9120                   |
| 150       | 330370   | 6387498  | Thick area of <i>Allocasuarina luehmannii</i> (Bulloak) over fence. <i>Hyparrhenia hirta</i> (Coolatai Grass) in easement.                                                                                                                                                              | 6/06/2024 | #9125<br>#9126<br>#9127<br>#9128 |
| 151       | 330313   | 6387548  | End of <i>Allocasuarina luehmannii</i> to the west.                                                                                                                                                                                                                                     | 6/06/2024 | #9129<br>#9130                   |
| 152       | 330239   | 6387608  | Disturbed, natives <i>Acacia parvipinnula</i> , <i>Allocasuarina luehmannii</i> . Introduced: <i>Hyparrhenia hirta</i> (Coolatai Grass) & <i>Chloris gayana</i> (Rhodes Grass), <i>Gomphocarpus fruticosus</i> .                                                                        | 6/06/2024 | #9131<br>#9132                   |
| 153       | 330105   | 6387721  | Area of <i>Allocasuarina luehmannii</i> (Bulloak) to ENE.                                                                                                                                                                                                                               | 6/06/2024 | #9133<br>#9134                   |
| 154       | 329957   | 6387795  |                                                                                                                                                                                                                                                                                         | 6/06/2024 | #9135<br>#9136                   |
| 155       | 329714   | 6387848  | In road reserve, Military land to north. Highly disturbed introduced species.                                                                                                                                                                                                           | 6/06/2024 | #9139<br>#9140                   |
| 156       | 329022   | 6388038  | Native tree either side of the easement. <i>Eucalyptus crebra</i> . Weeds <i>Tiger Pear</i> .                                                                                                                                                                                           | 6/06/2024 | #9142<br>#9143<br>#9144<br>#9145 |
| 157       | 328860   | 6388068  | Front of Singleton Army Base. Dead Red-necked Wallaby on side of the road.                                                                                                                                                                                                              | 6/06/2024 | #9153<br>#9152                   |
| 158       | 328523   | 6388137  | High disturbance roadside grasses. Natives <i>Lobelia concolor</i> , Weeds: <i>Setaria sphacelata</i> (South African Pigeon Grass), <i>Chloris gayana</i> (Rhodes Grass). Superb Wren.                                                                                                  | 6/06/2024 | #9157<br>#9158                   |
| 159       | 328277   | 6388052  | <i>Eucalyptus moluccana</i> (Grey Box). Stick Nest to NE.#9159.                                                                                                                                                                                                                         | 6/06/2024 | #9160<br>#9161                   |
| 160       | 328102   | 6387966  | Weedy Road reserve. <i>Eucalyptus tereticornis</i> (Forest Red Gum) in Army Land.                                                                                                                                                                                                       | 6/06/2024 | #9167<br>#9168                   |
| 161       | 328014   | 6387910  | Weedy Road reserve. Likely specimen of <i>Eucalyptus glaucina</i> (Slaty Red Gum) over fence within Army Base).                                                                                                                                                                         | 6/06/2024 | #9172<br>#9173<br>#9174          |
| 162       | 327961   | 6387882  | Weedy Road reserve. Potential <i>E. glaucina</i> (labelled No. 2) on its truck. Introduced <i>Mus musculus</i> (House Mouse) under debris.                                                                                                                                              | 6/06/2024 | #9175<br>#9176<br>#9177<br>#9178 |
| 163       | 327881   | 6387841  | Weedy Road reserve                                                                                                                                                                                                                                                                      | 6/06/2024 | #9180<br>#9181                   |
| 164       | 327771   | 6387830  | Weedy Road reserve, <i>Hyparrhenia hirta</i> (Coolatai Grass).                                                                                                                                                                                                                          | 6/06/2024 | #9182<br>#9183                   |
| 165       | 327461   | 6387818  | Weedy Road reserve, <i>Hyparrhenia hirta</i> (Coolatai Grass).                                                                                                                                                                                                                          | 6/06/2024 | #9184<br>#9185                   |
| 166       | 327379   | 6387824  | On boundary of road reserve and Singleton Army Base land. <i>Eucalyptus crebra</i> , <i>Allocasuarina luehmannii</i> (Bulloak), <i>Eucalyptus tereticornis</i> (Forest Red Gum). <i>Capillipedium spicigerum</i> (Scented Top Grass). Weeds: <i>Hyparrhenia hirta</i> (Coolatai Grass). | 6/06/2024 | #9186<br>#9187                   |

| Point No. | Location |          | Point Description & Impact                                                                                                                                                                                                                                                                                                                                                                                           | Date       | Photo and/or Photo No.                                               |
|-----------|----------|----------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|----------------------------------------------------------------------|
|           | Easting  | Northing |                                                                                                                                                                                                                                                                                                                                                                                                                      |            |                                                                      |
| 169       | 319230   | 6394107  | <u>Long Point</u> – Mining buffer land. <i>Daviesia genistifolia</i> (Broom Bitter Pea), <i>Aristida racemosa</i> , <i>Acacia falcata</i> , <i>Acacia cultriformis</i> (Knife-leaf Wattle).                                                                                                                                                                                                                          | 6/06/2024  | #9205<br>#9206                                                       |
| 170       | 319086   | 6394492  | <u>Long Point</u> – Native: <i>Sporobolus creber</i> , <i>Cynodon dactylon</i> (Couch). <i>Cymbopogon refractus</i> (Barbed Wire Grass)                                                                                                                                                                                                                                                                              | 6/06/2024  | #9207<br>#9208<br>#9209                                              |
| 171       | 318917   | 6394969  | <u>Long Point</u> – Within existing easement. Natives: <i>Acacia cultriformis</i> (Knife-leaf Wattle), <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Aristida vagans</i> , Outside easement: <i>Eucalyptus crebra</i> , <i>Allocasuarina luehmannii</i> (Bullock), <i>Eucalyptus tereticornis</i> (Forest Red Gum). Deer prints present on the ground.                                                         | 6/06/2024  | #9213                                                                |
| 172       | 318674   | 6395647  | <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Melinis repens</i> (Red Natel Grass). Possible <i>E. glaucina</i> outside existing easement.                                                                                                                                                                                                                                                                     | 6/06/2024  | #9218<br>#9219<br>#9220<br>#9221                                     |
| 173       | 318564   | 6396239  | <i>Sporobolus creber</i> , Weeds: <i>Eragrostis curvula</i> (African Love Grass)                                                                                                                                                                                                                                                                                                                                     | 6/06/2024  |                                                                      |
| 175       | 319352   | 6393766  | Area where feed goes into Redbank Power Station.<br>Natives: <i>Aristida vagans</i> , , <i>Acacia falcata</i> , <i>Cymbopogon refractus</i> (Barbed Wire Grass). Bush to east <i>Allocasuarina luehmannii</i> (Bullock) & <i>Eucalyptus moluccana</i> (Grey Box).                                                                                                                                                    | 6/06/2024  | #9230<br>#9231                                                       |
| 176       | 319459   | 6393452  | Railway line 200m to south. <i>Daviesia genistifolia</i> (Broom Bitter Pea), <i>Cymbopogon refractus</i> (Barbed Wire Grass), Weeds: <i>Verbena bonariensis</i> , <i>Senecio madagascariensis</i> (Fireweed). Bush to the east: <i>Eucalyptus moluccana</i> (Grey Box), <i>Myoporum montanum</i> .                                                                                                                   | 6/06/2024  | #9235<br>#9236                                                       |
| 177       | 319484   | 6393383  | <i>Acacia salicina</i> (Cooba),                                                                                                                                                                                                                                                                                                                                                                                      | 6/06/2024  | #9238                                                                |
| 178       | 319217   | 6399017  | Near Singleton Substation. Area of woodland will be shaved off. <i>E. crebra</i> , <i>Allocasuarina luehmannii</i> (Bullock), <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Cheilanthes sieberi</i> (Mulga Fern), <i>Sporobolus creber</i> . Weeds: <i>Senecio madagascariensis</i> (Fireweed).                                                                                                                | 11/06/2024 | #9337<br>#9338<br>#9339<br>#9340<br>#9341<br>#9342                   |
| 179       | 318907   | 6398832  | <i>E. crebra</i> , <i>Allocasuarina luehmannii</i> (Bullock), <i>Acacia cultriformis</i> (Knife-leaf Wattle) (common in easement). <i>Acacia decora</i> , <i>Eremophila debilis</i> (Amulla), <i>Chloris ventricosa</i> , <i>Myoporum montanum</i> . <i>Calotis lappulacea</i> (Yellow burr-daisy). Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive), <i>Senecio madagascariensis</i> (Fireweed). | 11/06/2024 | #9342<br>#9243<br>#9344<br>#9345<br>#9346<br>#9347<br>#9348<br>#9349 |
| 180       | 319023   | 6399115  | Two specimens of <i>E. crebra</i> likely to require removal. One tree has a hollow at its base.                                                                                                                                                                                                                                                                                                                      | 11/06/2024 | #9351                                                                |
| 181       | 319052   | 6399187  | Near Maison Dieu Road. (45m easement) Trees along road One <i>E. moluccana</i> with a small hollow. Probably just outside impact area. Two <i>E. crebra</i> on southern side of the road will require removal.                                                                                                                                                                                                       | 11/06/2024 | #9353<br>#9354                                                       |
| 182       | 319059   | 6399220  | 45m easement. Area of <i>Allocasuarina luehmannii</i> (Bullock), <i>Myoporum montanum</i> , <i>Acacia cultriformis</i> (Knife-leaf Wattle) within roadside reserve require removal.<br>Unkempt Grey-crowned Babbler nest.                                                                                                                                                                                            | 11/06/2024 | #9355<br>#9356                                                       |
| 183       | 319236   | 6399959  | <i>Sporobolus creber</i> . <i>Chloris ventricosa</i> , <i>Aristida racemosa</i> . Weeds: <i>Senecio madagascariensis</i> (Fireweed).                                                                                                                                                                                                                                                                                 | 11/06/2024 | #9359<br>#9360                                                       |
| 184       | 319281   | 6400160  | In drainage line: Introduced <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive) very thick. Small <i>E. crebra</i> & <i>A. luehmannii</i> on edge. <i>Angophora floribunda</i> in creek. <i>Sporobolus creber</i> .                                                                                                                                                                                        | 11/06/2024 | #9361<br>#9362                                                       |



| Point No. | Location |          | Point Description & Impact                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         | Date       | Photo and/or Photo No.                             |
|-----------|----------|----------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|----------------------------------------------------|
|           | Easting  | Northing |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |            |                                                    |
| 185       | 319289   | 6400224  | <i>Cynodon dactylon</i> (Couch). Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive), <i>Senecio madagascariensis</i> (Fireweed), <i>Gomphocarpus fruticosus</i> .                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 | 11/06/2024 | #9364<br>#9365                                     |
| 186       | 319326   | 6400387  | Area of <i>E. crebra</i> to the NE on edge will require removal. <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive) within existing easement.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | 11/06/2024 | #9367<br>#9368                                     |
| 187       | 319350   | 6400514  | Smaller section of <i>E. crebra</i> to east. <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive) to west.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 | 11/06/2024 | #9371<br>#9372                                     |
| 188       | 319512   | 6400117  | Next to Creekline. PCT 4015. Two specimens of <i>E. crebra</i> (likely branch trimming). <i>Angophora floribunda</i> on creekline (require removal). Cut down leave stump to hold streambank #9381.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                | 11/06/2024 | #9379<br>#9380                                     |
| 189       | 319530   | 6401253  | <i>Angophora floribunda</i> trees on creekline (require removal or thinning). <i>Casuarina glauca</i> (Swamp Oak) on eastern side #9385. <i>Lomandra longifolia</i> (Spiny Mat Rush) along creekline. <i>Austrostipa verticillata</i> and <i>Cynodon dactylon</i> .                                                                                                                                                                                                                                                                                                                                                                                                                                                | 11/06/2024 | #9382<br>#9383                                     |
| 190       | 319676   | 6404754  | Glennies Creek Crossing Natives: <i>Casuarina cunninghamiana</i> (River Oak), <i>Melia azedarach</i> (White Cedar), <i>Austrostipa verticillata</i> (Slender Bamboo Grass), <i>Cynodon dactylon</i> (Couch). <i>Phragmites australis</i> along creek. Introduced species: <i>Galenia pubescens</i> (Galenia), <i>Cestrum parqui</i> (Green Cestrum), <i>Lycium ferocissimum</i> (African Boxthorn), <i>Galium aparine</i> (Cleavers), <i>Ricinus communis</i> (Castor Oil Plant), <i>Senecio madagascariensis</i> (Fireweed), <i>Gomphocarpus fruticosus</i> , <i>Bidens pilosa</i> (Cobblers Pegs). A small number of specimens of River Oak will require removal. Stumps are to be left behind on the creekbank. | 11/06/2024 | #9398<br>#9399                                     |
| 191       | 319561   | 6405081  | Mine Land Ashton Coal. Paddock grasses. Natives: <i>Sporobolus creber</i> . Weeds: <i>Hyparrhenia hirta</i> (Coolatai Grass), <i>Senecio madagascariensis</i> (Fireweed), <i>Lycium ferocissimum</i> (African Boxthorn), <i>Paspalum dilatatum</i> (Paspalum), <i>Plantago lanceolata</i> (Plantain).                                                                                                                                                                                                                                                                                                                                                                                                              | 11/06/2024 | #9407<br>#9408                                     |
| 192       | 319397   | 6405543  | PCT 3431<br>Area of dense <i>Allocasuarina luehmannii</i> (Bulloak), Smaller trees <i>E. crebra</i> , <i>E. tereticornis</i> and <i>E. moluccana</i> , <i>Aristida racemosa</i> , <i>Sporobolus creber</i> , <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Lomandra multiflora</i> , <i>Dianella revolutum</i> . Weeds: <i>Galenia pubescens</i> (Galenia).                                                                                                                                                                                                                                                                                                                                                  | 11/06/2024 | #9412<br>#9413<br>#9414<br>#9415                   |
| 193       | 318686   | 6406500  | Area of bare soil. Dense <i>Allocasuarina luehmannii</i> (Bulloak), on northern side. <i>Aristida racemosa</i> . Weeds: <i>Melinis repens</i> (Red Natel Grass).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   | 11/06/2024 | #9417<br>#9418                                     |
| 194       | 318566   | 6406632  | <u>Bowmans Creek Crossing near New England Highway</u><br><i>Casuarina cunninghamiana</i> (River Oak), <i>Lobelia purpurea</i> (Whiteroot), Weeds: <i>Ricinus communis</i> (Castor Oil Plant), <i>Solanum pseudocapsicum</i> (Jerusalem Cherry), <i>Megathyrsus maximus</i> (Guinea Grass), <i>Araujia sericifera</i> (Moth Vine), <i>Galium aparine</i> (Cleavers), <i>Bidens pilosa</i> (Cobblers Pegs).<br>Area of River Oak on northern side will require removal. (stumps to be left <i>in-situ</i> the ground).<br><i>Crinia signifera</i> (Common Eastern Froglet) heard calling from within the creek.                                                                                                     | 11/06/2024 | #9423<br>#9424<br>#9425<br>#9426<br>#9427<br>#9428 |
| 220       | 317870   | 6409290  | Next to cultural meeting Point<br>Small area of regenerating trees. Natives: <i>Brachychiton populneus</i> (Kurrajong), <i>Geijera parviflora</i> (Wilga), <i>Eucalyptus moluccana</i> (Grey Box), <i>Aristida racemosa</i> . Weeds: <i>Sida rhombifolia</i> (Paddys Lucerne), <i>Eragrostis curvula</i> (African Lovegrass), <i>Chloris gayana</i> (Rhodes Grass), <i>Acacia baileyana</i> (Cootamundra Wattle).                                                                                                                                                                                                                                                                                                  | 1/07/2024  | #0317                                              |
| 221       | 317860   | 6409410  | Natives: <i>Cynodon dactylon</i> (Couch), <i>Aristida racemosa</i> . Weeds: <i>Sida rhombifolia</i> (Paddys Lucerne), <i>Cenchrus clandestinus</i> (Kikuyu). Rabbit diggings.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      | 1/07/2024  | #0321<br>#0322                                     |
| 222       | 318324   | 6409410  | New England Highway, near Lemington Road. Weedy roadside.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          | 1/07/2024  | #0324<br>#0325                                     |
| 223       | 318448   | 6407119  | <u>Bowmans Creek (Southern Side)</u><br>Some casuarinas will require removal. Natives: <i>Casuarina cunninghamiana</i> (River Oak). Weeds: <i>Ricinus communis</i> (Castor Oil Plant), <i>Galenia pubescens</i> (Galenia), <i>Megathyrsus maximus</i> (Guinea Grass), <i>Bidens pilosa</i> (Cobblers Pegs). <i>Crinia signifera</i> calling from Bowmans Creek.                                                                                                                                                                                                                                                                                                                                                    | 1/07/2024  | #0326<br>#0327<br>#0328<br>#0329                   |
| 224       | 317507   | 6410192  | Grassy Paddock, mainly introduced species. Weeds: <i>Tagetes minuta</i> (Stinking Roger), Cattle grazing.<br>Black-faced Cuckoo Shrike.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | 1/07/2024  | #0337<br>#0338                                     |

| Point No. | Location |          | Point Description & Impact                                                                                                                                                                                                                                                                                                                                               | Date      | Photo and/or Photo No.           |
|-----------|----------|----------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------|----------------------------------|
|           | Easting  | Northing |                                                                                                                                                                                                                                                                                                                                                                          |           |                                  |
| 225       | 317413   | 6410316  | <u>Bowmans Creek</u><br><i>Casuarina cunninghamiana</i> (River Oak) – either side of the bank. <i>Cynodon dactylon</i> (Couch), <i>Austrostipa verticillata</i> (Slender Bamboo Grass). Weeds: <i>Juncus acutus</i> , <i>Tagetes minuta</i> (Stinking Roger). A small number of River Oaks will be required to be removed.                                               | 1/07/2024 | #0340<br>#0341<br>#0342<br>#0343 |
| 226       | 314327   | 6415962  | Looking back toward mine pits. Easement mainly <i>Acacia salicina</i> (Cooba) with weeds <i>Bidens pilosa</i> (Cobbers Pegs), <i>Galium aparine</i> (Cleavers), <i>Gomphocarpus fruticosus</i> , <i>Chloris gayana</i> (Rhodes Grass), <i>Galenia pubescens</i> (Galenia).                                                                                               | 1/07/2024 | #0349<br>#0350<br>#0351          |
| 227       | 314268   | 6416133  | <i>Acacia salicina</i> (Cooba). Weeds: <i>Gomphocarpus fruticosus</i> , <i>Chloris gayana</i> (Rhodes Grass), <i>Hyparrhenia hirta</i> (Coolatai Grass), <i>Cirsium vulgare</i> (Black Thistle), <i>Opuntia stricta</i> (Prickly Pear), <i>Senecio madagascariensis</i> (Fireweed).                                                                                      | 1/07/2024 | #0359<br>#0360                   |
| 228       | 314207   | 6416293  | <i>Acacia salicina</i> (Cooba). Weeds: <i>Hyparrhenia hirta</i> (Coolatai Grass), <i>Galenia pubescens</i> (Galenia), <i>Lysimachia arvensis</i> (Scarlet Pimpernel), <i>Megathyrsus maximus</i> (Guinea Grass).                                                                                                                                                         | 1/07/2024 | #0362<br>#0363                   |
| 229       | 314153   | 6416441  | <i>Acacia salicina</i> (Cooba). Weeds: <i>Gomphocarpus fruticosus</i> , <i>Chloris gayana</i> (Rhodes Grass), <i>Megathyrsus maximus</i> (Guinea Grass).                                                                                                                                                                                                                 | 1/07/2024 | #0364<br>#0365                   |
| 230       | 314128   | 6416494  | Near pit undergoing rehabilitation<br><i>Megathyrsus maximus</i> (Guinea Grass), <i>Galenia pubescens</i> (Galenia).                                                                                                                                                                                                                                                     | 1/07/2024 | #0366<br>#0367                   |
| 231       | 314742   | 6417030  | <i>Acacia salicina</i> (Cooba), <i>Acacia elongata</i> , <i>Lobelia purpurescens</i> (Whiteroot), <i>Dichondra repens</i> (Kidneyweed), <i>Einadia hastata</i> (Berry Saltbush). Weeds: <i>Gomphocarpus fruticosus</i> , <i>Chloris gayana</i> (Rhodes Grass), <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive), <i>Senecio madagascariensis</i> (Fireweed). | 1/07/2024 | #0368<br>#0369                   |
| 232       | 312862   | 6418187  | Point on Hebdon Road<br>Possible removal of two specimens of <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark). Weeds: <i>Chloris gayana</i> (Rhodes Grass), <i>Hyparrhenia hirta</i> (Coolatai Grass).                                                                                                                                                                  | 1/07/2024 | #0373                            |
| 233       | 312723   | 6418138  | Corner pole near Lake Liddell<br><i>Aristida racemosa</i> . Weeds: <i>Juncus acutus</i> (Spike Rush).                                                                                                                                                                                                                                                                    | 1/07/2024 | #0375<br>#0376                   |
| 234       | 312425   | 6418278  | Next to Lake Liddell <i>Juncus acutus</i> (Spike Rush), <b>White-bellied Sea Eagle observed flying over lake.</b>                                                                                                                                                                                                                                                        | 1/07/2024 | #0386<br>#0387                   |
| 236       | 312065   | 6418523  | Location of proposed Eastern Hub near the northern shore of Lake Liddell. Entire area of Eastern Hub composed of very low grasses such as <i>Cynodon dactylon</i> (Couch).                                                                                                                                                                                               | 1/07/2024 | #0401<br>#0402<br>#0403          |
| 237       | 311992   | 6418679  | Planted Native Vegetation <i>Eucalyptus moluccana</i> (Grey Box), <i>Eucalyptus blakelyi</i> (Blackley's Red Gum), <i>Eucalyptus punctata</i> (Grey Gum),                                                                                                                                                                                                                | 1/07/2024 | #0409<br>#0410<br>#0411<br>#0412 |
| 238       | 312002   | 6418702  | Planted Native Vegetation Next to Hebdon Road. Line connects to existing poles over Hebdon Road to the north.                                                                                                                                                                                                                                                            | 1/07/2024 | #0413<br>#0414                   |
| 239       | 303405   | 6430146  | Existing easement <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Sporobolus creber</i> (Slender Rats-tail Grass). <i>Eucalyptus crebra</i> to side of easement. Weeds: <i>Chloris gayana</i> (Rhodes Grass).                                                                                                                                                        | 4/06/2024 | #0090<br>#0091                   |
| 240       | 303464   | 6430384  | Narrow area of E. crebra to east, likely requires removal. <i>Myoporum montanum</i> , <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Austrostipa verticillata</i> (Slender Bamboo Grass). Weeds: <i>Senecio madagascariensis</i> (Fireweed).                                                                                                                        | 4/06/2024 | #0096<br>#0097                   |
| 241       | 303471   | 6430427  | Bend in easement <i>Cynodon dactylon</i> (Couch). Weeds: <i>Chloris gayana</i> (Rhodes Grass), <i>Nothoscordum gracile</i> (Onion Weed), <i>Plantago lanceolata</i> (Plantain).                                                                                                                                                                                          | 4/06/2024 | #0199<br>#0100                   |
| 242       | 303649   | 6430461  | Weeds: <i>Chloris gayana</i> (Rhodes Grass), <i>Nothoscordum gracile</i> (Onion Weed), <i>Plantago lanceolata</i> (Plantain). Yellow-faced Whip Snake.                                                                                                                                                                                                                   | 4/06/2024 | #0101<br>#0102                   |
| 243       | 303717   | 6430471  | Easement over a Dam<br>Clamorous reed warbler. Weeds: <i>Senecio madagascariensis</i> (Fireweed).                                                                                                                                                                                                                                                                        | 4/06/2024 | #0104<br>#0105                   |

| Point No. | Location |          | Point Description & Impact                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        | Date       | Photo and/or Photo No. |
|-----------|----------|----------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|------------------------|
|           | Easting  | Northing |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |            |                        |
|           |          |          |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |            | #0106                  |
| 244       | 303847   | 6430593  | Natives: <i>Microlaena stipoides</i> (Weeping Meadow Grass), <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Bulbine bulbosa</i> . Specimens of <i>E. crebra</i> either side of the easement.                                                                                                                                                                                                                                                                                                                 | 4/06/2024  | #0132<br>#0133         |
| 245       | 303937   | 6430679  | Low maintained easement. <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Sporobolus creber</i> (Slender Rats-tail Grass). <i>Arthropodium strictum</i> (Chocolate Lily).                                                                                                                                                                                                                                                                                                                                      | 4/06/2024  | #0173<br>#0174         |
| 246       | 304089   | 6430810  | <i>E. crebra</i> smaller trees will likely require removal. <i>Acacia paradoxa</i> (Kangaroo Wattle), <i>Cymbopogon refractus</i> (Barbed Wire Grass). <b>Brown Treecreeper present to north.</b>                                                                                                                                                                                                                                                                                                                 | 4/06/2024  | #0175                  |
| 247       | 304152   | 6430869  | <i>Eucalyptus blakelyi</i> (Blakely's Red Gum) side of easement. <i>Aristida racemosa</i> , <i>Cheilanthes sieberi</i> (Mulga Fern).                                                                                                                                                                                                                                                                                                                                                                              | 4/06/2024  | #0175                  |
| 248       | 304257   | 6430987  | Drainage line. <i>Eucalyptus blakelyi</i> (Blakely's Red Gum) to SSE. <i>Acacia paradoxa</i> (Kangaroo Wattle)., Red-necked Wallaby, <i>Crinia signifera</i> (Common Eastern Froglet), <i>Limnodynastes tasmaniensis</i> (Spotted Marsh Frog).                                                                                                                                                                                                                                                                    | 4/06/2024  | #0176<br>#0177         |
| 249       | 304415   | 6431134  | <i>Eucalyptus blakelyi</i> (Blakely's Red Gum) to SE.                                                                                                                                                                                                                                                                                                                                                                                                                                                             | 4/06/2024  | #0185<br>#0186         |
| 250       | 304728   | 6431439  | Derived Grassland. <i>Senecio madagascariensis</i> (Fireweed)-common.                                                                                                                                                                                                                                                                                                                                                                                                                                             | 4/06/2024  | #0189<br>#0190         |
| 251       | 304933   | 6431635  | <i>Sporobolus creber</i> (Slender Rats-tail Grass). <i>E. crebra</i> on easement edge.                                                                                                                                                                                                                                                                                                                                                                                                                            | 4/06/2024  | #0193<br>#0194         |
| 252       | 304975   | 6431668  | <i>E. crebra</i> on easement edge (possible branch trimming).                                                                                                                                                                                                                                                                                                                                                                                                                                                     | 4/06/2024  | #0195                  |
| 253       | 305032   | 6431720  | <i>E. crebra</i> may require trimming or removal.                                                                                                                                                                                                                                                                                                                                                                                                                                                                 | 4/06/2024  | #0196                  |
| 254       | 305138   | 6431825  | <i>E. crebra</i> may require trimming or removal.                                                                                                                                                                                                                                                                                                                                                                                                                                                                 | 4/06/2024  | #0197                  |
| 255       | 305498   | 6432141  | <i>E. crebra</i> on easement edge. Red-necked Wallaby.                                                                                                                                                                                                                                                                                                                                                                                                                                                            | 4/06/2024  | #0251<br>#0250         |
| 256       | 305692   | 6432267  | Disturbed paddock. Better quality derived grassland to east.                                                                                                                                                                                                                                                                                                                                                                                                                                                      | 4/06/2024  | #0252<br>#0253         |
| 257       | 306043   | 6432585  | End of the line. <i>Cymbopogon refractus</i> (Barbed Wire Grass),                                                                                                                                                                                                                                                                                                                                                                                                                                                 | 4/06/2024  | #0263<br>#0264         |
| 258       | 358914   | 6369420  | Southern side of Hunter Expressway, across from Kurri STS. <i>Eucalyptus parramattensis</i> subsp. <i>decadens</i> (Drooping Red Gum), <i>Eucalyptus fibrosa</i> (Broadleaved Ironbark), thick <i>Melaleuca nodosa</i> (Ball Honey myrtle).                                                                                                                                                                                                                                                                       | 4/04/2025  | #3084<br>#3085         |
| 259       | 326660   | 6387930  | Golden Highway (Mitchell Line of Road) – in drainage line. Canopy: <i>Angophora floribunda</i> (Rough-barked Apple), <i>Eucalyptus tereticornis</i> (Forest Red Gum), <i>Eucalyptus moluccana</i> (Grey Box). Shrub <i>Acacia implexa</i> (Hickory Wattle). Ground: <i>Juncus usitatus</i> . Weeds <i>Cirsium vulgare</i> (Black Thistle), <i>Rumex crispus</i> (Curled Dock).                                                                                                                                    | 12/11/2025 | #5375<br>#5376         |
| 260       | 326621   | 6387947  | Golden Highway (Mitchell Line of Road) –Canopy: <i>Eucalyptus tereticornis</i> (Forest Red Gum), <i>Eucalyptus moluccana</i> (Grey Box). Ground: <i>Juncus usitatus</i> . Weeds <i>Cirsium vulgare</i> (Black Thistle), <i>Rumex crispus</i> (Curled Dock). Fauna: Koel, Willy Wagtail, Masked Lapwing, Channell Bill Cuckoo.                                                                                                                                                                                     | 12/11/2025 | #5378<br>#5379         |
| 261       | 326454   | 6387996  | Golden Highway (Mitchell Line of Road) – Drainage Line to east. * <i>Acacia saligna</i> .                                                                                                                                                                                                                                                                                                                                                                                                                         | 12/11/2025 | #5384<br>#5385         |
| 262       | 326328   | 6388050  | Golden Highway (Mitchell Line of Road) – Open shrubby area. <i>Angophora floribunda</i> (Rough-barked Apple), <i>Acacia falcata</i> (Falcate Wattle), <i>Lomandra multiflora</i> , <i>Dichondra repens</i> (Kidney Weed), <i>Cheilanthes sieberi</i> (Mulga Fern). Weeds: <i>Cirsium vulgare</i> (Black Thistle), <i>Chloris gayana</i> (Rhodes Grass), <i>Hyparrhenia hirta</i> (Coolatai Grass). Fauna Pied Currawong, Magpie, Australian Raven.                                                                | 12/11/2025 | #5386<br>#5387         |
| 263       | 326252   | 6388073  | Golden Highway (Mitchell Line of Road) –Canopy: <i>Eucalyptus fibrosa</i> (Broad-leaved Ironbark), <i>Allocasuarina luehmannii</i> (Bullock). Shrub: <i>Daviesia ulicifolia</i> (Gorse Bitter Pea), <i>Cassinia quinquefaria</i> , <i>Ozothamnus diosmifolius</i> (White Dogwood), <i>Acacia cultriformis</i> (Knife Leaf Wattle). Ground <i>Dichondra repens</i> (Kidney Weed), <i>Hardenbergia violacea</i> , <i>Lomandra multiflora</i> (Many Flowered Mat Rush). Fauna: Grey Shrike Thrush, Australian Raven. | 12/11/2025 | #5393<br>#5394         |



| Point No. | Location |          | Point Description & Impact                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           | Date       | Photo and/or Photo No. |
|-----------|----------|----------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|------------------------|
|           | Easting  | Northing |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |            |                        |
| 264       | 326091   | 6388129  | Golden Highway (Mitchell Line of Road) – <i>Angophora floribunda</i> (Rough-barked Apple), <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Allocasuarina luehmannii</i> (Bullock). Shrub: <i>Breynia oblongifolia</i> (Breynia), <i>Olearia elliptica</i> (Sticky Daisy Bush), <i>Acacia parvippinula</i> (Silver-stemmed Wattle). Ground <i>Dichondra repens</i> (Kidney Weed), <i>Hardenbergia violacea</i> , <i>Lomandra multiflora</i> (Many Flowered Mat Rush).<br>Weed: <i>Sida rhombifolia</i> (Paddy's Lucerne), <i>Hyparrhenia hirta</i> (Coolatai Grass), <i>Conyza</i> sp., <i>Hypochaeris radicata</i> (Flatweed). | 12/11/2025 | #5401<br>#5402         |
| 265       | 325942   | 6388180  | Golden Highway (Mitchell Line of Road) – <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Allocasuarina luehmannii</i> (Bullock). Shrub: <i>Acacia parvippinula</i> (Silver-stemmed Wattle). Ground <i>Rytidosperma</i> sp. (Wallaby Grass), <i>Dichondra repens</i> (Kidney Weed), <i>Enchylaena tomentosa</i> (Ruby Saltbush), <i>Eremophila debilis</i> (Amulla).<br>Weeds: <i>Opuntia aurantiaca</i> (Tiger Pear), <i>Opuntia stricta</i> (Prickly Pear), <i>Verbena bonariensis</i> (Purple Top), <i>Hypochaeris radicata</i> (Flatweed), <i>Lantana camara</i> (Lantana).                                                 | 12/11/2025 | #5426<br>#5427         |
| 266       | 325864   | 6388214  | Golden Highway (Mitchell Line of Road) –Canopy: <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Eucalyptus tereticornis</i> (Forest Red Gum), <i>Allocasuarina luehmannii</i> (Bullock). Shrub: <i>Olearia elliptica</i> (Sticky Daisy Bush). Ground <i>Dichondra repens</i> (Kidney Weed), <i>Calotis lappulacea</i> (Yellow Burr-daisy).<br>Weeds: <i>Sida rhombifolia</i> (Paddy's Lucerne), <i>Cirsium vulgare</i> (Black Thistle), <i>Lantana camara</i> (Lantana), <i>Bidens pilosa</i> (Cobblers Pegs), <i>Verbena rigida</i> , <i>Plantago lanceolata</i> (Plantain).                                                  | 12/11/2025 | #5438<br>#5439         |
| 267       | 325753   | 6388264  | Golden Highway (Mitchell Line of Road) –Canopy: <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Eucalyptus moluccana</i> (Grey Box). Shrub: <i>Melia azedarach</i> (White Cedar). Ground <i>Austrostipa verticillata</i> (Speargrass), <i>Dichelachne micrantha</i> (Plumegrass).<br>Weeds: <i>Sida rhombifolia</i> (Paddy's Lucerne), <i>Hyparrhenia hirta</i> (Coolatai Grass), <i>Opuntia aurantiaca</i> (Tiger Pear), <i>Lantana camara</i> (Lantana), <i>Verbena rigida</i> , <i>Eragrostis curvula</i> (African Lovegrass), <i>Plantago lanceolata</i> (Plantain).<br>Fauna: Black-faced Cuckoo Shrike.                  | 12/11/2025 | #5455<br>#5456         |
| 268       | 325679   | 6388301  | Golden Highway (Mitchell Line of Road) –Canopy: <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Eucalyptus moluccana</i> (Grey Box). Shrub: <i>Olearia elliptica</i> (Sticky Daisy Bush). Ground <i>Calotis lappulacea</i> (Yellow Burr-daisy), <i>Asperula conferta</i> .<br>Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive), <i>Heliotropium amplexicaule</i> (Blue Heliotrope), <i>Opuntia aurantiaca</i> (Tiger Pear), <i>Hyparrhenia hirta</i> (Coolatai Grass), <i>Lantana camara</i> (Lantana), <i>Plantago lanceolata</i> (Plantain).                                                              | 12/11/2025 | #5469<br>#5470         |
| 269       | 325517   | 6388391  | Golden Highway (Mitchell Line of Road) –Canopy: <i>Eucalyptus moluccana</i> (Grey Box), <i>Allocasuarina luehmannii</i> (Bullock).. Shrub: <i>Olearia elliptica</i> (Sticky Daisy Bush), <i>Pimelea linifolia</i> (Rice Flower). Ground <i>Dichondra repens</i> (Kidney Weed), <i>Calotis lappulacea</i> (Yellow Burr-daisy), <i>Dichelachne micrantha</i> (Plumegrass), <i>Hardenbergia violacea</i> .<br>Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive). <i>Aloe vera</i> , <i>Opuntia aurantiaca</i> (Tiger Pear), <i>Plantago lanceolata</i> (Plantain).                                                    | 12/11/2025 | #5486<br>#5487         |
| 270       | 325428   | 6388439  | Golden Highway (Mitchell Line of Road) –Canopy: <i>Eucalyptus moluccana</i> (Grey Box), <i>Allocasuarina luehmannii</i> (Bullock).. Shrub: <i>Olearia elliptica</i> (Sticky Daisy Bush), <i>Pimelea linifolia</i> (Rice Flower). Ground <i>Dichondra repens</i> (Kidney Weed), <i>Calotis lappulacea</i> (Yellow Burr-daisy), <i>Dichelachne micrantha</i> (Plumegrass), <i>Hardenbergia violacea</i> .<br>Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive). <i>Opuntia aurantiaca</i> (Tiger Pear), <i>Plantago lanceolata</i> (Plantain).<br>Fauna: <i>Carlia tetradactyla</i> (Southern Rainbow Skink).        | 12/11/2025 | #5507<br>#5508         |
| 271       | 325338   | 6388487  | Golden Highway (Mitchell Line of Road) –Canopy: <i>Eucalyptus moluccana</i> (Grey Box), <i>Allocasuarina luehmannii</i> (Bullock).. Shrub: <i>Olearia elliptica</i> (Sticky Daisy Bush). Ground <i>Dichondra repens</i> (Kidney Weed), <i>Calotis lappulacea</i> (Yellow Burr-daisy), <i>Dichelachne micrantha</i> (Plumegrass), <i>Hardenbergia violacea</i> .<br>Weeds: <i>Eragrostis curvula</i> (African Lovegrass), <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive). <i>Cirsium vulgare</i> (Black Thistle), <i>Plantago lanceolata</i> (Plantain).                                                                | 12/11/2025 | #5511<br>#5512         |

| Point No. | Location |          | Point Description & Impact                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      | Date       | Photo and/or Photo No.  |
|-----------|----------|----------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|-------------------------|
|           | Eastings | Northing |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |            |                         |
| 272       | 325269   | 6388527  | Golden Highway (Mitchell Line of Road) – Weeds: <i>Chloris gayana</i> (Rhodes Grass). Fauna: Grey-crowned Babbler (BC Act).                                                                                                                                                                                                                                                                                                                                                                                     | 12/11/2025 | #5518<br>#5519          |
| 273       | 324985   | 6388684  | Near corner of Golden Highway (Mitchell Line of Road) and Putty Road – Highly disturbed area. Weeds: <i>Galenia pubescens</i> (Galenia), <i>Bidens pilosa</i> (Cobbers Pegs), <i>Conyza</i> sp., <i>Ricinus communis</i> (Castor Oil Plant), <i>Verbena bonariensis</i> (Purple Top), <i>Gomphocarpus fruticosus</i> (Swan Plant), <i>Chloris gayana</i> (Rhodes Grass).                                                                                                                                        | 12/11/2025 | #5520<br>#5521          |
| 274       | 324901   | 6388670  | East of Putty Road: Canopy: <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Allocasuarina luehmannii</i> (Bulloak).. Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive). <i>Plantago lanceolata</i> (Plantain), <i>Chloris gayana</i> (Rhodes Grass), <i>Pavonia hastata</i> , <i>Opuntia aurantiaca</i> (Tiger Pear), <i>Opuntia stricta</i> (Prickly Pear).                                                                                                                            | 12/11/2025 | #5535<br>#5536          |
| 275       | 324818   | 6388660  | East of Putty Road: Canopy: <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Allocasuarina luehmannii</i> (Bulloak). Ground <i>Dichondra repens</i> (Kidney Weed), <i>Eremophila debilis</i> (Amulla).. Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive). <i>Plantago lanceolata</i> (Plantain), <i>Chloris gayana</i> (Rhodes Grass), <i>Pavonia hastata</i> , <i>Verbena bonariensis</i> (Purple Top), <i>Opuntia aurantiaca</i> (Tiger Pear), <i>Opuntia stricta</i> (Prickly Pear). | 12/11/2025 | #5541<br>#5542          |
| 276       | 324693   | 6388658  | East of Putty Road: Canopy: <i>Eucalyptus moluccana</i> (Grey Box) to south. Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive). <i>Plantago lanceolata</i> (Plantain), <i>Schinus molle</i> var. <i>areira</i> (Pepper Tree), <i>Foeniculum vulgare</i> (Fennel), <i>Chloris gayana</i> (Rhodes Grass), <i>Verbena bonariensis</i> (Purple Top).                                                                                                                                              | 12/11/2025 | #5550<br>#5551          |
| 277       | 324573   | 6388664  | East of Putty Road: Start of native vegetation to the south. Canopy: <i>Eucalyptus moluccana</i> (Grey Box), <i>Allocasuarina luehmannii</i> (Bulloak). Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive). <i>Lantana camara</i> (Lantana), <i>Plantago lanceolata</i> (Plantain), <i>Chloris gayana</i> (Rhodes Grass).                                                                                                                                                                      | 12/11/2025 | #5555<br>#5556          |
| 278       | 324517   | 6388658  | East of Putty Road: Canopy: <i>Eucalyptus moluccana</i> (Grey Box), <i>Allocasuarina luehmannii</i> (Bulloak). Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive). <i>Lantana camara</i> (Lantana), <i>Chloris gayana</i> (Rhodes Grass).                                                                                                                                                                                                                                                      | 12/11/2025 | #5557<br>#5558          |
| 279       | 324404   | 6388687  | East of Putty Road: Canopy: <i>Allocasuarina luehmannii</i> (Bulloak)(Dominant), <i>Eucalyptus crebra</i> . Shrubs: <i>Notelaea microcarpa</i> (Native Olive). Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive). <i>Lantana camara</i> (Lantana), <i>Chloris gayana</i> (Rhodes Grass). Fauna: Striped Skink.                                                                                                                                                                                | 12/11/2025 | #5566<br>#5567          |
| 280       | 324259   | 6388725  | East of Putty Road: Canopy: <i>Eucalyptus crebra</i> , <i>Angophora floribunda</i> (Rough-barked Apple). Shrub <i>Acacia salicina</i> (Cooba). Ground <i>Dichondra repens</i> (Kidney Weed), <i>Dichelachne micrantha</i> (Plumegrass). Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive). <i>Lantana camara</i> (Lantana), <i>Chloris gayana</i> (Rhodes Grass). Fauna: Red-Necked Wallaby.                                                                                                  | 12/11/2025 | #5573<br>#5574          |
| 281       | 324147   | 6388809  | From intersection of Putty and Broke Roads. Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive). <i>Lantana camara</i> (Lantana), <i>Foeniculum vulgare</i> (Fennel), <i>Chloris gayana</i> (Rhodes Grass).                                                                                                                                                                                                                                                                                     | 12/11/2025 | #5582                   |
| 282       | 321940   | 6389648  | Mount Thorley STS: Canopy: <i>Eucalyptus moluccana</i> (Grey Box), Weeds: <i>Chloris gayana</i> (Rhodes Grass), <i>Galenia pubescens</i> (Galenia), <i>Bidens pilosa</i> (Cobbers Pegs), <i>Lycium ferocissimum</i> (African Boxthorn).                                                                                                                                                                                                                                                                         | 12/11/2025 | #5593<br>#5594          |
| 283       | 321856   | 6389675  | Mount Thorley STS: Canopy: <i>Eucalyptus moluccana</i> (Grey Box), <i>Brachychiton populneus</i> (Kurrajong). Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive). <i>Chloris gayana</i> (Rhodes Grass), <i>Galenia pubescens</i> (Galenia), <i>Bidens pilosa</i> (Cobbers Pegs).                                                                                                                                                                                                               | 12/11/2025 | #5598<br>#5599<br>#5600 |
| 284       | 321843   | 6389844  | Mount Thorley: Canopy: <i>Eucalyptus moluccana</i> (Grey Box), <i>Eucalyptus crebra</i> Shrub <i>Acacia salicina</i> (Cooba). Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive). <i>Schinus molle</i> var. <i>areira</i> (Pepper Tree), <i>Chloris gayana</i> (Rhodes Grass), <i>Galenia pubescens</i> (Galenia), <i>Bidens pilosa</i> (Cobbers Pegs).                                                                                                                                        | 12/11/2025 | #5595<br>#5596          |
| 285       | 320740   | 6392517  | Golden Highway: Area of the Endangered Population (NSW BC Act) <i>Acacia pendula</i> (Weeping Myall). Other natives <i>Marsilea drummondii</i> (Nardoo), <i>Lobelia concolor</i> (Poison Pratia).                                                                                                                                                                                                                                                                                                               | 12/11/2025 | #5620<br>#5621          |
| 286       | 320563   | 6392824  | Easement crossing over Golden Highway: <i>Allocasuarina luehmannii</i> (Bulloak), <i>Eucalyptus blakelyi</i> (Blakely's Red Gum)                                                                                                                                                                                                                                                                                                                                                                                | 12/11/2025 | #5613                   |

| Point No. | Location |          | Point Description & Impact                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | Date       | Photo and/or Photo No. |
|-----------|----------|----------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|------------------------|
|           | Easting  | Northing |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |            |                        |
| 287       | 312558   | 6418264  | Revised impact areas Eastern Hub (Lake Liddell, Muswellbrook), Open low grazed grassy area, west planted trees <i>Eucalyptus blakelyi</i> (Blakely's Red Gum), <i>Eucalyptus crebra</i> , <i>Melaleuca stypheloides</i> (Prickly-leaved Paperbark), <i>Casuarina glauca</i> (Swamp Oak). Weeds: <i>Phyla canescens</i> (Lippia).                                                                                                                                                                                                                      | 22/01/2026 | #9224<br>#9225         |
| 288       | 312558   | 6418264  | Revised impact areas Eastern Hub (Lake Liddell, Muswellbrook). Tree planting to east. <i>Eucalyptus moluccana</i> (Grey Box)(dbh 0.47m) 7m west may require removal.                                                                                                                                                                                                                                                                                                                                                                                  | 22/01/2026 | #9231<br>#9232         |
| 289       | 312457   | 6418298  | Revised impact areas Eastern Hub (Lake Liddell, Muswellbrook). <i>Casuarina glauca</i> (Swamp Oak) to west. Weeds: <i>Phyla canescens</i> (Lippia), <i>Juncus acutus</i> (Spiny Mat Rush).                                                                                                                                                                                                                                                                                                                                                            | 22/01/2026 | #9251<br>#9252         |
| 290       | 312397   | 6418344  | Revised impact areas Eastern Hub (Lake Liddell, Muswellbrook). West of point – Planted <i>Eucalyptus blakelyi</i> (Blakely's Red Gum), <i>Casuarina glauca</i> (Swamp Oak). Weeds: <i>Phyla canescens</i> (Lippia), <i>Juncus acutus</i> (Spiny Mat Rush).                                                                                                                                                                                                                                                                                            | 22/01/2026 | #9255<br>#9256         |
| 291       | 312350   | 6418371  | Revised impact areas Eastern Hub (Lake Liddell, Muswellbrook). Near drainage. Planted <i>Eucalyptus blakelyi</i> (Blakely's Red Gum), <i>Corymbia maculata</i> (Spotted Gum), <i>Callistemon</i> sp.                                                                                                                                                                                                                                                                                                                                                  | 22/01/2026 | #9259<br>#9260         |
| 292       | 312252   | 6418405  | Revised impact areas Eastern Hub (Lake Liddell, Muswellbrook). West other side of drainage line from 291. East planted <i>Corymbia maculata</i> (Spotted Gum) and <i>Eucalyptus melliodora</i> (Yellow Box).                                                                                                                                                                                                                                                                                                                                          | 22/01/2026 | #9264<br>#9265         |
| 293       | 312138   | 6418603  | Revised impact areas Eastern Hub (Lake Liddell, Muswellbrook). North planted trees: <i>Eucalyptus blakelyi</i> (Blakely's Red Gum), <i>Eucalyptus moluccana</i> (Grey Box), <i>Melaleuca linariifolia</i> (Snow in Summer), <i>Acacia pendula</i> (Weeping Myall).                                                                                                                                                                                                                                                                                    | 22/01/2026 | #9267<br>#9268         |
| 294       | 312151   | 6418659  | Revised impact areas Eastern Hub (Lake Liddell, Muswellbrook). North Hebdon Road, highly disturbed Weeds: <i>Chloris gayana</i> (Rhodes Grass).                                                                                                                                                                                                                                                                                                                                                                                                       | 22/01/2026 | #9271<br>#9272         |
| 295       | 312672   | 6418428  | Revised impact areas Eastern Hub (Lake Liddell, Muswellbrook). Highly disturbed Weeds: <i>Chloris gayana</i> (Rhodes Grass).                                                                                                                                                                                                                                                                                                                                                                                                                          | 22/01/2026 | #9273<br>#9274         |
| 296       | 332018   | 6387270  | <i>Casuarina glauca</i> (Swamp Oak).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  | 20/11/2025 | #6657<br>#6658         |
| 297       | 332348   | 6387213  | Shrub - <i>Acacia parvipinnula</i> (Silver-stemmed Wattle),                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           | 20/11/2025 | #6664<br>#6665         |
| 298       | 332167   | 6387196  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       | 20/11/2025 | #6670                  |
| 299       |          |          | Near Roadworks Office. <i>Eucalyptus moluccana</i> (Grey Box), <i>Acacia parvipinnula</i> (Silver-stemmed Wattle). <i>Allocasuarina luehmannii</i> (Bulloak). Weeds <i>Cirsium vulgare</i> (Black Thistle).                                                                                                                                                                                                                                                                                                                                           |            |                        |
| 300       | 332338   | 6387182  | Weeds: <i>Rumex crispus</i> (Curled Dock), <i>Cirsium vulgare</i> (Black Thistle).                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | 20/11/2025 | #6692<br>#6693         |
| 306       | 340746   | 6385431  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |            | #8549                  |
| 309       | 340745   | 6385446  | Old New England Highway.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |            | #8555<br>#8556         |
| 319       | 340869   | 6385325  | <b>Lower Belford</b> , Canopy: Mostly younger <i>Eucalyptus moluccana</i> (Grey Box), some <i>Eucalyptus tereticornis</i> (Forest Red Gum). Ground cover grazed low, natives present <i>Aristida vagans</i> (Three-awn Grass).                                                                                                                                                                                                                                                                                                                        | 10/03/2026 | #1403<br>#1405         |
| 320       | 340949   | 6385214  | <b>Lower Belford</b> , Ground cover grazed low, natives present <i>Microlaena stipoides</i> (Weeping Meadow Grass), <i>Cyanthillium cinereum</i> (Ironweed). Large specimen of <i>Eucalyptus fibrosa</i> (Broad-leaved Ironbark) nearby.                                                                                                                                                                                                                                                                                                              | 10/03/2026 | #1491<br>#1492         |
| 321       | 341056   | 6385052  | <b>Lower Belford, Within easement</b> Ground cover grazed low, Scattered <i>Eucalyptus tereticornis</i> (Forest Red Gum).                                                                                                                                                                                                                                                                                                                                                                                                                             | 10/03/2026 | #1436                  |
| 324       | 340791   | 6385383  | <b>Lower Belford</b> , Ground cover grazed low, Scattered <i>Eucalyptus tereticornis</i> (Forest Red Gum).                                                                                                                                                                                                                                                                                                                                                                                                                                            | 10/03/2026 | #1454<br>#1455         |
| 337       | 340734   | 6385469  | <b>Lower Belford</b> , Open Forest: Canopy- <i>Corymbia maculata</i> (Spotted Gum)(Dominant), <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), Shrub <i>Breynia oblongifolia</i> (Breynia), <i>Cassinia quinquefaria</i> , <i>Pultenaea spinosa</i> . Ground; <i>Lomandra longifolia</i> (Spiny Mat Rush), <i>Lomandra multiflora</i> (Many-flowered Mat Rush), <i>Dianella revoluta</i> , <i>Brachyscome multifida</i> . Climber – <i>Glycine clandestina</i> (Love Creeper). Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive). | 20/05/2026 | #4896<br>#4897         |



| Point No. | Location |          | Point Description & Impact                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          | Date       | Photo and/or Photo No. |
|-----------|----------|----------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|------------------------|
|           | Easting  | Northing |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |            |                        |
| 338       | 340721   | 6385487  | <b>Lower Belford</b> , Corner of easement next to the New England Highway. Open Forest: Canopy- <i>Corymbia maculata</i> (Spotted Gum)(Dominant). Shrubs: <i>Acacia falcata</i> (Falcate Wattle), <i>Breynia oblongifolia</i> (Breynia), <i>Pultenaea spinosa</i> . Ground; <i>Lomandra multiflora</i> (Many-flowered Mat Rush), <i>Dianella revoluta</i> , <i>Pomax umbelata</i> . Weeds: Closer to road - <i>Melinis repens</i> (Red Natel Grass), <i>Eragrostis curvula</i> (African Lovegrass).                                                                                                                                                                                                                                                                                                                                                                                                                                                 | 20/05/2026 | #4899<br>#4900         |
| 339       | 340618   | 6385482  | <b>Lower Belford</b> Along New England Highway. Canopy- <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), Shrub - <i>Jacksonia scoparia</i> (Dogwood), <i>Acacia falcata</i> (Falcate Wattle), <i>Leucopogon juniperinus</i> (Bearded Heath). Ground; <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Lomandra multiflora</i> (Many-flowered Mat Rush), <i>Dianella revoluta</i> , <i>Calotis cuneifolia</i> (Purple Burr-daisy), Climber – <i>Glycine clandestina</i> (Love Creeper). Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive). Closer to road - <i>Melinis repens</i> (Red Natel Grass), <i>Eragrostis curvula</i> (African Lovegrass).                                                                                                                                                                                                                                                                           | 20/05/2026 | #4907<br>#4908         |
| 340       | 340536   | 6385468  | <b>Lower Belford</b> Along New England Highway. Existing detention basin to west. <i>Typha orientalis</i> (Cumbungi). Open Forest: Canopy- <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus moluccana</i> (Grey Box). Shrub - <i>Acacia falcata</i> (Falcate Wattle), <i>Acacia parvipinnula</i> (Silver-stemmed Wattle). Ground; <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Lomandra multiflora</i> (Many-flowered Mat Rush), <i>Dianella revoluta</i> , Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive), <i>Megathyrsus maximus</i> (Guinea Grass), <i>Verbena rigida</i> . <b><i>Crinia signifera</i> (Common Eastern Froglet)</b> heard calling from detention basin.                                                                                                                                                                                                                                         | 20/05/2026 | #4916<br>#4917         |
| 341       | 340409   | 6385464  | <b>Lower Belford</b> Along New England Highway. Open Forest: Canopy- <i>Corymbia maculata</i> (Spotted Gum)(Dominant), <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Eucalyptus moluccana</i> (Grey Box). Shrub <i>Breynia oblongifolia</i> (Breynia), <i>Acacia falcata</i> (Falcate Wattle), <i>Bursaria spinosa</i> (Blackthorn). Ground; <i>Lomandra longifolia</i> (Spiny Mat Rush), <i>Lomandra multiflora</i> (Many-flowered Mat Rush). Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive). <i>Megathyrsus maximus</i> (Guinea Grass), <i>Eragrostis curvula</i> (African Lovegrass).<br><b>The majority of the trees appear to have been planted as part of works related to past roadworks for the section of New England Highway through Lower Belford which was completed in the early 1990's.</b>                                                                                                              | 20/05/2026 | #4918<br>#4919         |
| 342       | 340214   | 6385450  | <b>Lower Belford</b> Along New England Highway. Open Forest: Canopy- <i>Corymbia maculata</i> (Spotted Gum)(Dominant), Shrub <i>Breynia oblongifolia</i> (Breynia), <i>Pultenaea spinosa</i> . Ground; <i>Lomandra multiflora</i> (Many-flowered Mat Rush), <i>Calotis cuneifolia</i> (Purple Burr-daisy), <i>Hibbertia diffusa</i> . Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive). <i>Megathyrsus maximus</i> (Guinea Grass), <i>Eragrostis curvula</i> (African Lovegrass).<br><b>The majority of the trees appear to have been planted as part of works related to past roadworks for the section of New England Highway through Lower Belford which was completed in the early 1990's.</b>                                                                                                                                                                                                                               | 20/05/2026 | #4121<br>#4922         |
| 343       | 340125   | 6385449  | <b>Lower Belford</b> Along New England Highway. Open Forest: Canopy- <i>Corymbia maculata</i> (Spotted Gum)(Dominant), <i>Eucalyptus fibrosa</i> (Broad-leaved Ironbark), <i>Eucalyptus moluccana</i> (Grey Box), Shrub <i>Breynia oblongifolia</i> (Breynia), <i>Acacia falcata</i> (Falcate Wattle), <i>Olearia elliptica</i> (Sticky Daisy Bush), <i>Pultenaea spinosa</i> . Ground; <i>Lomandra multiflora</i> (Many-flowered Mat Rush), <i>Dianella revoluta</i> , <i>Cheilanthes sieberi</i> (Mulga Fern) Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive). <i>Megathyrsus maximus</i> (Guinea Grass), <i>Eragrostis curvula</i> (African Lovegrass), <i>Bryophyllum delagoense</i> (Mother of Millions).<br><b>The majority of the trees appear to have been planted as part of works related to past roadworks for the section of New England Highway through Lower Belford which was completed in the early 1990's.</b> | 20/05/2026 | #4925<br>#4926         |
| 344       | 340091   | 6385449  | <b>Lower Belford</b> Along New England Highway. Small gaps in open forest. Weeds: <i>Megathyrsus maximus</i> (Guinea Grass), <i>Eragrostis curvula</i> (African Lovegrass), <i>Bryophyllum delagoense</i> (Mother of Millions).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     | 20/05/2026 | #4927<br>#4928         |
| 345       | 340034   | 6385453  | <b>Lower Belford</b> Along New England Highway. Open Forest: Canopy- <i>Corymbia maculata</i> (Spotted Gum)(Dominant), <i>Eucalyptus fibrosa</i> (Broad-leaved Ironbark). Disturbed on the ground. Weeds: <i>Megathyrsus maximus</i> (Guinea Grass), <i>Chloris gayana</i> (Rhodes Grass), <i>Verbena bonariensis</i> (Purple Top). .                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               | 20/05/2026 | #4929<br>#4930         |

| Point No. | Location |          | Point Description & Impact                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | Date       | Photo and/or Photo No. |
|-----------|----------|----------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|------------------------|
|           | Easting  | Northing |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |            |                        |
| 346       | 339932   | 6385457  | <b>Lower Belford</b> Along New England Highway. Open disturbed area; Weeds: <i>Chloris gayana</i> (Rhodes Grass), <i>Hyparrhenia hirta</i> (Coolatai Grass), <i>Verbena bonariensis</i> (Purple Top), <i>Verbena rigida</i> , <i>Bidens pilosa</i> (Cobblers Pegs).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   | 20/05/2026 | #4932<br>#4933         |
| 347       | 339790   | 6385469  | <b>Lower Belford</b> Along New England Highway. Open Forest: Canopy- <i>Corymbia maculata</i> (Spotted Gum)(Dominant), <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Eucalyptus sideroxylon</i> (Mugga). Shrub <i>Breynia oblongifolia</i> (Breynia), <i>Acacia falcata</i> (Falcate Wattle), Ground; <i>Lomandra longifolia</i> (Spiny Mat Rush), <i>Lomandra multiflora</i> (Many-flowered Mat Rush).<br>Weeds: <i>Olea europaea</i> subsp. <i>cuspidata</i> (African Olive). <i>Pavonia hastata</i> .<br><b>The majority of the trees appear to have been planted as part of works related to past roadworks for the section of New England Highway through Lower Belford which was completed in the early 1990's.</b>                                                                                                                                                     | 20/05/2026 | #4943<br>#4944         |
| 348       | 339660   | 6385483  | <b>Lower Belford</b> Along New England Highway. Open Forest: Canopy- <i>Corymbia maculata</i> (Spotted Gum)(Dominant), Shrub <i>Breynia oblongifolia</i> (Breynia), <i>Jacksonia scoparia</i> (Dogwood), <i>Dillwynia retorta</i> , <i>Acacia falcata</i> (Falcate Wattle), <i>Acacia parvipinnula</i> (Silver-stemmed Wattle). Ground; <i>Lomandra multiflora</i> (Many-flowered Mat Rush), <i>Lepidosperma laterale</i> (Sword Grass). <i>Hardenbergia violacea</i> (False Sarsaparilla)<br>Weeds: <i>Megathyrsus maximus</i> (Guinea Grass), <i>Eragrostis curvula</i> (African Lovegrass).<br><b>The majority of the trees appear to have been planted as part of works related to past roadworks for the section of New England Highway through Lower Belford which was completed in the early 1990's.</b><br><b><i>Ctenotus robusta</i> (Striped Skink)</b> found under debris. | 20/05/2026 | #4947<br>#4948         |
| 349       | 339539   | 6385503  | <b>Lower Belford</b> Along New England Highway, just east of intersection with Pothana Road. Introduced maintained grasses on ground: <i>Chloris gayana</i> (Rhodes Grass), <i>Hyparrhenia hirta</i> (Coolatai Grass), <i>Melinis repens</i> (Red Natel Grass).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       | 20/05/2026 | #4956<br>#4957         |
| 350       | 339448   | 6385523  | <b>Lower Belford</b> Along New England Highway. Open Forest: Canopy- <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus moluccana</i> (Grey Box). Shrub - <i>Acacia falcata</i> (Falcate Wattle), <i>Breynia oblongifolia</i> (Breynia), <i>Acacia parvipinnula</i> (Silver-stemmed Wattle).<br>Disturbed open ground to west: <i>Chloris gayana</i> (Rhodes Grass), <i>Hyparrhenia hirta</i> (Coolatai Grass), <i>Acacia fimbriata</i> (Fimbriate Wattle).<br><b>The majority of the trees appear to have been planted as part of works related to past roadworks for the section of New England Highway through Lower Belford which was completed in the early 1990's.</b>                                                                                                                                                                                                       | 20/05/2026 | #4959<br>#4960         |
| 351       | 339330   | 6385560  | <b>Lower Belford</b> Along New England Highway. Steep Slope back from Highway. Open Forest: Canopy- <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Eucalyptus sideroxylon</i> (Mugga).<br><i>Megathyrsus maximus</i> (Guinea Grass), <i>Lantana camara</i> (Lantana).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  | 20/05/2026 | #4969<br>#4970         |
| 352       | 339154   | 6385610  | <b>Lower Belford</b> Along New England Highway. Steep Slope back from Highway. Canopy- <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus sideroxylon</i> (Mugga). Shrub: <i>Acacia fimbriata</i> (Fimbriate Wattle).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              | 20/05/2026 | #4974<br>#4975         |
| 353       | 338994   | 6385641  | <b>Lower Belford</b> Along New England Highway. Steep Slope back from Highway. Canopy- <i>Corymbia maculata</i> (Spotted Gum), Shrub: <i>Acacia fimbriata</i> (Fimbriate Wattle), <i>Breynia oblongifolia</i> (Breynia), <i>Acacia parvipinnula</i> (Silver-stemmed Wattle). <i>Dillwynia retorta</i> .<br>Weeds: <i>Lantana camara</i> (Lantana), <i>Eragrostis curvula</i> (African Lovegrass).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     | 20/05/2026 | #4979<br>#4980         |
| 354       | 338906   | 6385658  | <b>Lower Belford</b> Along New England Highway. Canopy- <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus moluccana</i> (Grey Box). Shrub - <i>Acacia fimbriata</i> (Fimbriate Wattle),<br>Weeds: <i>Melinis repens</i> (Red Natel Grass), <i>Verbena bonariensis</i> (Purple Top), <i>Verbena rigida</i> .                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       | 20/05/2026 | #4986<br>#4987         |
| 355       | 338829   | 6385676  | <b>Lower Belford</b> Along New England Highway. Canopy- <i>Eucalyptus tereticornis</i> (Forest Red Gum), <i>Eucalyptus sideroxylon</i> (Mugga).<br>Weeds: <i>Chloris gayana</i> (Rhodes Grass), <i>Hyparrhenia hirta</i> (Coolatai Grass).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | 20/05/2026 | #4989                  |

## **APPENDIX B**

### **DETAILED MAPS OF REZ**





|          |          |
|----------|----------|
| Job Ref  | 12875    |
| A4 Scale | 1:12,000 |

Disclaimer: While all reasonable care has been taken to ensure the information contained on this map is up to date and accurate, no guarantee is given that the information portrayed is free from error or omission. In addition the spatial accuracy of the map is wholly dependant on source data. Please verify the accuracy of all information prior to use. Development footprint areas should be used for indicative areas only.





Figure B2

Segment 1

25 June 2026

WILDTHING  
Environmental Consultants  
(a Division of Tattersall Lander Pty Ltd)  
ABN 41 003 509 215





|          |          |
|----------|----------|
| Job Ref  | 12875    |
| A4 Scale | 1:12,000 |

Disclaimer: While all reasonable care has been taken to ensure the information contained on this map is up to date and accurate, no guarantee is given that the information portrayed is free from error or omission. In addition the spatial accuracy of the map is wholly dependant on source data. Please verify the accuracy of all information prior to use. Development footprint areas should be used for indicative areas only.




Figure B3

Segment 2


25 June 2026

WILDTHING  
Environmental Consultants

(a Division of Tattersall Lander Pty Ltd)

ABN 41 003 509 215





|          |          |
|----------|----------|
| Job Ref  | 12875    |
| A4 Scale | 1:12,000 |

Disclaimer: While all reasonable care has been taken to ensure the information contained on this map is up to date and accurate, no guarantee is given that the information portrayed is free from error or omission. In addition the spatial accuracy of the map is wholly dependant on source data. Please verify the accuracy of all information prior to use. Development footprint areas should be used for indicative areas only.

0 250 500 m

Map Projection GDA2020 MGA Zone 56

Data Sources: LPI (2023),


Figure B4

**Segment 3**


25 June 2026

**WILDTHING**  
Environmental Consultants  
(a Division of Tattersall Lander Pty Ltd)  
ABN 41 003 509 215























|          |          |
|----------|----------|
| Job Ref  | 12875    |
| A4 Scale | 1:12,000 |

Disclaimer: While all reasonable care has been taken to ensure the information contained on this map is up to date and accurate, no guarantee is given that the information portrayed is free from error or omission. In addition the spatial accuracy of the map is wholly dependant on source data. Please verify the accuracy of all information prior to use. Development footprint areas should be used for indicative areas only.




Figure B9

Segment 8


25 June 2026

WILDTHING  
Environmental Consultants

(a Division of Tattersall Lander Pty Ltd)

ABN 41 003 509 215





|          |          |
|----------|----------|
| Job Ref  | 12875    |
| A4 Scale | 1:12,000 |

Disclaimer: While all reasonable care has been taken to ensure the information contained on this map is up to date and accurate, no guarantee is given that the information portrayed is free from error or omission. In addition the spatial accuracy of the map is wholly dependant on source data. Please verify the accuracy of all information prior to use. Development footprint areas should be used for indicative areas only.

Map Projection GDA2020 MGA Zone 56  
Data Sources: LPI (2023), NearMap (2024)


Figure B10

**Segment 9**

11 September 2026

**WILDTHING**  
Environmental Consultants  
(a Division of Tattersall Lander Pty Ltd)  
ABN 41 003 509 215





|          |          |
|----------|----------|
| Job Ref  | 12875    |
| A4 Scale | 1:12,000 |

Disclaimer: While all reasonable care has been taken to ensure the information contained on this map is up to date and accurate, no guarantee is given that the information portrayed is free from error or omission. In addition the spatial accuracy of the map is wholly dependant on source data. Please verify the accuracy of all information prior to use. Development footprint areas should be used for indicative areas only.

0 250 500 m

Map Projection GDA2020 MGA Zone 56

Data Sources: LPI (2023), NearMap (2024)


Figure B11

Segment 10


11 September 2026

WILDTHING  
Environmental Consultants  
(a Division of Tattersall Lander Pty Ltd)  
ABN 41 003 509 215









|          |         |
|----------|---------|
| Job Ref  | 12875   |
| A4 Scale | 1:4,000 |

Disclaimer: While all reasonable care has been taken to ensure the information contained on this map is up to date and accurate, no guarantee is given that the information portrayed is free from error or omission. In addition the spatial accuracy of the map is wholly dependant on source data. Please verify the accuracy of all information prior to use. Development footprint areas should be used for indicative areas only.




Figure B11.2

Segment 10b

11 September 2026

WILDTHING  
Environmental Consultants

(a Division of Tattersall Lander Pty Ltd)  
ABN 41 003 509 215





|          |          |
|----------|----------|
| Job Ref  | 12875    |
| A4 Scale | 1:12,000 |

Disclaimer: While all reasonable care has been taken to ensure the information contained on this map is up to date and accurate, no guarantee is given that the information portrayed is free from error or omission. In addition the spatial accuracy of the map is wholly dependant on source data. Please verify the accuracy of all information prior to use. Development footprint areas should be used for indicative areas only.

0 250 500 m

Map Projection GDA2020 MGA Zone 56

Data Sources: LPI (2023), NearMap (2024)


Figure B12

**Segment 11**

25 June 2026

**WILDTHING**  
Environmental Consultants  
(a Division of Tattersall Lander Pty Ltd)  
ABN 41 003 509 215





|          |          |
|----------|----------|
| Job Ref  | 12875    |
| A4 Scale | 1:12,000 |

Disclaimer: While all reasonable care has been taken to ensure the information contained on this map is up to date and accurate, no guarantee is given that the information portrayed is free from error or omission. In addition the spatial accuracy of the map is wholly dependant on source data. Please verify the accuracy of all information prior to use. Development footprint areas should be used for indicative areas only.




Figure B13

Segment 12


11 September 2026

WILDTHING  
Environmental Consultants

(a Division of Tattersall Lander Pty Ltd)

ABN 41 003 509 215





|          |          |
|----------|----------|
| Job Ref  | 12875    |
| A4 Scale | 1:12,000 |

Disclaimer: While all reasonable care has been taken to ensure the information contained on this map is up to date and accurate, no guarantee is given that the information portrayed is free from error or omission. In addition the spatial accuracy of the map is wholly dependant on source data. Please verify the accuracy of all information prior to use. Development footprint areas should be used for indicative areas only.




Figure B14


Segment 13

11 September 2026


WILDTHING  
Environmental Consultants

(a Division of Tattersall Lander Pty Ltd)  
ABN 41 003 509 215









|          |          |
|----------|----------|
| Job Ref  | 12875    |
| A4 Scale | 1:12,000 |

Disclaimer: While all reasonable care has been taken to ensure the information contained on this map is up to date and accurate, no guarantee is given that the information portrayed is free from error or omission. In addition the spatial accuracy of the map is wholly dependant on source data. Please verify the accuracy of all information prior to use. Development footprint areas should be used for indicative areas only.




Figure B16

Segment 15

11 September 2026

WILDTHING  
Environmental Consultants

(a Division of Tattersall Lander Pty Ltd)

ABN 41 003 509 215





|          |          |
|----------|----------|
| Job Ref  | 12875    |
| A4 Scale | 1:12,000 |

Disclaimer: While all reasonable care has been taken to ensure the information contained on this map is up to date and accurate, no guarantee is given that the information portrayed is free from error or omission. In addition the spatial accuracy of the map is wholly dependant on source data. Please verify the accuracy of all information prior to use. Development footprint areas should be used for indicative areas only.




Figure B17

Segment 16


11 September 2026

WILDTHING  
Environmental Consultants

(a Division of Tattersall Lander Pty Ltd)

ABN 41 003 509 215





|          |          |
|----------|----------|
| Job Ref  | 12875    |
| A4 Scale | 1:12,000 |

Disclaimer: While all reasonable care has been taken to ensure the information contained on this map is up to date and accurate, no guarantee is given that the information portrayed is free from error or omission. In addition the spatial accuracy of the map is wholly dependant on source data. Please verify the accuracy of all information prior to use. Development footprint areas should be used for indicative areas only.



Map Projection GDA2020 MGA Zone 56


Data Sources: LPI (2023), NearMap (2024)

Figure B18


## Segment 17

11 September 2026


























































|          |          |
|----------|----------|
| Job Ref  | 12875    |
| A4 Scale | 1:12,000 |

Disclaimer: While all reasonable care has been taken to ensure the information contained on this map is up to date and accurate, no guarantee is given that the information portrayed is free from error or omission. In addition the spatial accuracy of the map is wholly dependant on source data. Please verify the accuracy of all information prior to use. Development footprint areas should be used for indicative areas only.



Map Projection GDA2020 MGA Zone 56  
Data Sources: LPI (2024), Nearmap (2024)

Figure B30


Segment 29

11 September 2026

**WILDTHING**  
Environmental Consultants

(a Division of Tattersall Lander Pty Ltd)  
ABN 41 003 509 215





|          |          |
|----------|----------|
| Job Ref  | 12875    |
| A4 Scale | 1:12,000 |

Disclaimer: While all reasonable care has been taken to ensure the information contained on this map is up to date and accurate, no guarantee is given that the information portrayed is free from error or omission. In addition the spatial accuracy of the map is wholly dependant on source data. Please verify the accuracy of all information prior to use. Development footprint areas should be used for indicative areas only.





Figure B31

Segment 30

11 September 2026

WILDTHING  
Environmental Consultants  
(a Division of Tattersall Lander Pty Ltd)  
ABN 41 003 509 215





|          |         |
|----------|---------|
| Job Ref  | 12875   |
| A4 Scale | 1:2,400 |

Disclaimer: While all reasonable care has been taken to ensure the information contained on this map is up to date and accurate, no guarantee is given that the information portrayed is free from error or omission. In addition the spatial accuracy of the map is wholly dependant on source data. Please verify the accuracy of all information prior to use. Development footprint areas should be used for indicative areas only.





Figure B32  
**Liddell Switching Station**


11 September 2026

**WILDTHING**  
Environmental Consultants  
(a Division of Tattersall Lander Pty Ltd)  
ABN 41 003 509 215














## **APPENDIX C**

### **SECTION 7.3 OF THE BC ACT 2016 TEST OF SIGNIFICANCE**



## C1.0 SECTION 7.3 OF THE BC ACT 2016 TEST OF SIGNIFICANCE (THREATENED ECOLOGICAL COMMUNITIES)

### c1.1 KURRI SAND WOODLAND IN THE SYDNEY BASIN BIOREGION

#### Kurri Sand Swamp Woodland in the Sydney Basin Bioregion

|                                         |                                                                                                                                                                                                                                                                                                                              |
|-----------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Description</b>                      | Kurri Sand Swamp Woodland is a low woodland or heathland, generally with a low open canopy rarely exceeding 15 m in height and a shrubby understorey. The overstorey is usually dominated by <i>Eucalyptus parramattensis</i> subsp. <i>decadens</i> (Parramatta Red Gum) and <i>Angophora bakeri</i> (Narrow-leaved Apple). |
| <b>Habitat Requirements and Ecology</b> | Occurs on soils developed on poorly drained Tertiary sand deposits that blanket Permian sediments.                                                                                                                                                                                                                           |
| <b>Distribution</b>                     | Known to occur in the Kurri Kurri–Cessnock area of the Cessnock LGA in the lower Hunter Valley.                                                                                                                                                                                                                              |
| <b>BC Act 2016</b>                      | Endangered Ecological Community                                                                                                                                                                                                                                                                                              |
| <b>EPBC Act 1999</b>                    | Endangered Ecological Community                                                                                                                                                                                                                                                                                              |

#### ***Five Part Test of Significance***

***The objective of section 7.3 of the Biodiversity Conservation Act 2016 (BC Act), the test of significance, is to provide standardised and transparent consideration of threatened species and ecological communities, and their habitats, through the development assessment process. The following is to be taken into account for the purposes of determining whether a proposed development or activity is likely to significantly affect threatened species or ecological communities, or their habitats:***

- a. *have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction,*

NA

- b. *in the case of an endangered ecological community or critically endangered ecological community, whether the proposed development or activity:*
- is likely to have an adverse effect on the extent of the ecological community such that its local occurrence is likely to be placed at risk of extinction, or*
  - is likely to substantially and adversely modify the composition of the ecological community such that its local occurrence is likely to be placed at risk of extinction,*

Areas mapped as PCT 3631 Kurri Sand Clay Woodland occur west of the Kurri Kurri STS and west of Hart Road were found to be consistent with the Endangered Ecological Community; Kurri Sand Woodland in the Sydney Basin Bioregion. Between the Kurri STS and the Hunter Expressway and a section west of the Hunter Expressway across from the Kurri STS there was no existing easement. These areas had been subject to previous disturbance with areas highly disturbed to relatively intact. Between Hart Road and the Hunter Expressway this EEC occurs within an existing easement and is composed of low maintained grown covers with a moderate diversity. Better quality areas of Kurri Sand Heathy Woodland occur either side of the easement. The proposal will result in the removal/modification of a relatively small amount of Kurri Sand Swamp Woodland. Both areas of Kurri Sand Swamp Woodland are connected to much larger significant areas of this EEC. The proposal will result in an incremental reduction in Kurri Sand Swamp Woodland; however, it is considered unlikely to have a significant impact which might lead to extinction locally.

- c. *in relation to the habitat of a threatened species or ecological community:*
- the extent to which habitat is likely to be removed or modified as a result of the proposed development or activity, and*
  - whether an area of habitat is likely to become fragmented or isolated from other areas of habitat as a result of the proposed development or activity, and*
  - the importance of the habitat to be removed, modified, fragmented or isolated to the long-term survival of the species or ecological community in the locality,*

The REZ will impact 0.60ha Kurri Sand Swamp Woodland for approximately 150m of a new section of easement between the Kurri STS and Hunter Expressway (0.2a) and another section on the other side of

### Kurri Sand Swamp Woodland in the Sydney Basin Bioregion

the Hunter Expressway (0.40ha). The REZ is will only have a small impact on Kurri Sand Swamp Woodland within the existing easement west of Hart Road. Areas of impacted Kurri Sand Swamp Woodland were connected to much larger areas of this EEC. No areas are likely to become fragmented or isolated from other areas that are important to the long-term survival of Kurri Sand Swamp Woodland in the locality.

- d. *whether the proposed development or activity is likely to have an adverse effect on any declared area of outstanding biodiversity value (either directly or indirectly),*

No areas declared areas of outstanding biodiversity value were present within proximity to the REZ proposal.

- e. *whether the proposed development or activity is or is part of a key threatening process or is likely to increase the impact of a key threatening process.*

The 'Key Threatening Processes' currently listed under Schedule 4 of the BC Act 2016 that are relevant to the proposal have been addressed below:

#### **Clearing of Native Vegetation.**

The proposal will result in the removal of native vegetation and may be viewed as being part of this Key Threatening Process. The action is unlikely to be responsible for the significant loss of this EEC provided that recommendations for impact minimisation as listed within Section 6.4 are undertaken.

#### **Invasion of native plant communities by exotic perennial grasses.**

Specimens of exotic perennial grasses such as *Chloris gayana* (Rhodes Grass), *Eragrostis curvula* (African Love Grass), *Andropogon virginicus* (Whisky Grass) and *Hyparrhenia hirta* (Coolatai Grass) were present. These exotic grasses compete with species within this EEC for space, light, and other resources and may threaten its persistence at several sites. The action is unlikely to be responsible for the significant loss of this EEC provided recommendations for impact minimisation as listed within Section 6.4 are undertaken.

#### **High frequency fire resulting in the disruption of life cycle processes in plants and animals and loss of vegetation structure and composition**

It is unknown what impact fire has had within areas of this EEC. The proposal is unlikely to increase

#### **Competition and grazing by the feral European rabbit, *Oryctolagus cuniculus***

*Observations and secondary indications of the European Rabbit such as scats and diggings were noted along parts of the REZ. The proposal is not likely to result in an increase in feral numbers of this introduced species.*

#### **Herbivory and environmental degradation caused by feral deer**

Tracks of Feral Deer were recorded within several sections of the REZ. Feral Deer would be considered to have an impact on native flora in the local area. The proposal is not likely to result in an increase in feral numbers of this introduced species.

#### **Infection of native plants by *Phytophthora cinnamomi***

No evidence of *Phytophthora cinnamomi* was observed during site visits. The proposal is unlikely to cause the spread of this fungus.

#### **Invasion, establishment and spread of Lantana (*Lantana camara*)**

Lantana was recorded within parts of the REZ proposal. The proposal is unlikely to cause the spread of this weed species.

#### **Introduction and establishment of Exotic Rust Fungi of the order Pucciniales pathogenic on plants of the family Myrtaceae.**

No evidence of the fungi was observed during site visits. The proposal is unlikely to cause the spread of this fungus.

#### **Invasion of native plant communities by African Olive *Olea europaea* subsp. *cuspidata***

The African Olive was recorded along parts of the REZ route. The proposal is unlikely to cause further spread of this introduced tree.



## Kurri Sand Swamp Woodland in the Sydney Basin Bioregion

### Removal of dead wood and dead trees

Dead wood and trees were recorded along small sections of the REZ route. Dead wood and trees such be avoided wherever possible.

### Predation, habitat degradation, competition and disease transmission by *Sus scrofa* (Feral Pigs)

Signs of Feral Pigs were noted along parts of the REZ route. The proposal is not likely to result in an increase in feral numbers of this introduced species.

### Conclusion

The proposal will result in an incremental reduction in Kurri Sand Swamp Woodland; however, it is considered unlikely to have a significant impact on this EEC.

### Bibliography:

Bell, S. and Driscoll, C. (2007) Vegetation of the Cessnock-Kurri region, Cessnock LGA, New South Wales: Survey, Classification & Mapping. Report to Department of Environment & Climate Change, Newcastle.

NSW Scientific Committee (2001) Kurri sand swamp woodland in the Sydney Basin Bioregion- Endangered ecological community determination - final. DEC (NSW), Sydney.

## c1.2 LOWER HUNTER SPOTTED GUM-IRONBARK FOREST

| Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW North Coast Bioregions |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|---------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Description</b>                                                                          | This community is dominated by <i>Corymbia maculata</i> (Spotted Gum) and <i>Eucalyptus fibrosa</i> (Broad-leaved Ironbark) while <i>E. punctata</i> (Grey Gum) and <i>E. crebra</i> (Narrow-leaved Ironbark) occur occasionally. The understorey is marked by the tall shrub, <i>Acacia parvipinnula</i> , and by the prickly shrubs, <i>Daviesia ulicifolia</i> , <i>Bursaria spinosa</i> , <i>Melaleuca nodosa</i> and <i>Lissanthe strigosa</i> . The ground layer is diverse; frequent species include <i>Cheilanthes sieberi</i> , <i>Cymbopogon refractus</i> , <i>Dianella revoluta</i> , <i>Entolasia stricta</i> , <i>Glycine clandestina</i> , <i>Lepidosperma laterale</i> , <i>Lomandra multiflora</i> , <i>Microlaena stipoides</i> , <i>Pomax umbellata</i> , <i>Pratia purpurascens</i> , <i>Themeda australis</i> and <i>Phyllanthus hirtellus</i> . |
| <b>Habitat Requirements and Ecology</b>                                                     | Occurs principally on Permian geology in the central to lower Hunter Valley. The Permian substrates most commonly supporting the community belong to the Dalwood Group, the Maitland Group and the Greta and Tomago Coal Measures, although smaller areas of the community may also occur on the Permian Singleton and Newcastle Coal Measures and the Triassic Narrabeen Group.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| <b>Distribution</b>                                                                         | Restricted to a range of approximately 65 km by 35 km centred on the Cessnock - Beresfield area in the Central and Lower Hunter Valley.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| <b>BC Act 2016</b>                                                                          | Endangered Ecological Community                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| <b>EPBC Act 1999</b>                                                                        | Not listed                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |

**Five-Part Test of Significance**

***The objective of section 7.3 of the Biodiversity Conservation Act 2016 (BC Act), the test of significance, is to provide standardised and transparent consideration of threatened species and ecological communities, and their habitats, through the development assessment process. The following is to be taken into account for the purposes of determining whether a proposed development or activity is likely to significantly affect threatened species or ecological communities, or their habitats:***

- a. *have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction,*

NA

- b. *in the case of an endangered ecological community or critically endangered ecological community, whether the proposed development or activity:*
- is likely to have an adverse effect on the extent of the ecological community such that its local occurrence is likely to be placed at risk of extinction, or*
  - is likely to substantially and adversely modify the composition of the ecological community such that its local occurrence is likely to be placed at risk of extinction,*

Areas mapped as PCT 3433 Hunter Coast Foothills Spotted Gum-Ironbark Forest and PCT 3444 Lower Hunter Spotted Gum-Ironbark Forest were found to be consistent with the EEC; Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW North Coast Bioregions. Sections of this EEC were present along the Hunter Expressway near Kurri Kurri, NW of Harts Road and East of Old Maitland Road. This EEC largely occurred within areas of existing easement, composed of low maintained grown covers with a varying diversity. Better quality areas of Lower Hunter Spotted Gum Ironbark Forest were generally present on the edges of the existing easement. The proposal will result in the removal/modification of a relatively small amount (0.53ha) of generally disturbed Lower Hunter Spotted Gum Ironbark Forest resulting in an incremental reduction in the local area, however, it is considered unlikely to have a significant impact which might lead to extinction locally.

- c. *in relation to the habitat of a threatened species or ecological community:*
- the extent to which habitat is likely to be removed or modified as a result of the proposed development or activity, and*
  - whether an area of habitat is likely to become fragmented or isolated from other areas of habitat as a result of the proposed development or activity, and*
  - the importance of the habitat to be removed, modified, fragmented or isolated to the long-term survival of the species or ecological community in the locality,*

The REZ is will only have a relatively small impact on Lower Hunter Spotted Gum Ironbark Forest within areas of existing easement and parts of the easement edge. No areas are likely to become fragmented or



### Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW North Coast Bioregions

isolated from other areas that are important to the long-term survival of Lower Hunter Spotted Gum Ironbark Forest in the locality.

- d. *whether the proposed development or activity is likely to have an adverse effect on any declared area of outstanding biodiversity value (either directly or indirectly),*

No areas declared areas of outstanding biodiversity value were present within proximity to the REZ proposal.

- e. *whether the proposed development or activity is or is part of a key threatening process or is likely to increase the impact of a key threatening process.*

The 'Key Threatening Processes' currently listed under Schedule 4 of the BC Act 2016 that are relevant to the proposal have been addressed below:

#### **Clearing of Native Vegetation.**

The proposal will result in the removal of native vegetation and may be viewed as being part of this Key Threatening Process. The action is unlikely to be responsible for the significant loss of this EEC provided that recommendations for impact minimisation as listed within Section 6.4 are undertaken.

#### **Invasion of native plant communities by exotic perennial grasses.**

Specimens of exotic perennial grasses such as *Chloris gayana* (Rhodes Grass), *Eragrostis curvula* (African Love Grass), *Andropogon virginicus* (Whisky Grass) and *Hyparrhenia hirta* (Coolatai Grass) were present. These exotic grasses compete with species within this EEC for space, light, and other resources and may threaten its persistence at several sites. The action is unlikely to be responsible for the significant loss of this EEC provided recommendations for impact minimisation as listed within Section 6.4 are undertaken.

#### **High frequency fire resulting in the disruption of life cycle processes in plants and animals and loss of vegetation structure and composition**

It is unknown what impact fire has had within areas of this EEC. The proposal is unlikely to increase

#### **Competition and grazing by the feral European rabbit, *Oryctolagus cuniculus***

Observations and secondary indications of the European Rabbit such as scats and diggings were noted along parts of the REZ. The proposal is not likely to result in an increase in feral numbers of this introduced species.

#### **Herbivory and environmental degradation caused by feral deer**

Tracks of Feral Deer were recorded within several sections of the REZ. Feral Deer would be considered to have an impact on native flora in the local area. The proposal is not likely to result in an increase in feral numbers of this introduced species.

#### **Infection of native plants by *Phytophthora cinnamomi***

No evidence of *Phytophthora cinnamomi* was observed during site visits. The proposal is unlikely to cause the spread of this fungus.

#### **Invasion, establishment and spread of Lantana (*Lantana camara*)**

Lantana was recorded within parts of the REZ proposal. The proposal is unlikely to cause the spread of this weed species.

#### **Introduction and establishment of Exotic Rust Fungi of the order Pucciniales pathogenic on plants of the family Myrtaceae.**

No evidence of the fungi was observed during site visits. The proposal is unlikely to cause the spread of this fungus.

#### **Invasion of native plant communities by African Olive *Olea europaea* subsp. *cuspidata***

The African Olive was recorded along parts of the REZ route. The proposal is unlikely to cause further spread of this introduced tree.

## Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW North Coast Bioregions

### Removal of dead wood and dead trees

Dead wood and trees were recorded along small sections of the REZ route. Dead wood and trees such be avoided wherever possible.

### Predation, habitat degradation, competition and disease transmission by *Sus scrofa* (Feral Pigs)

Signs of Feral Pigs were noted along parts of the REZ route. The proposal is not likely to result in an increase in feral numbers of this introduced species.

### Conclusion

The proposal will result in a small incremental reduction in Lower Hunter Spotted Gum Ironbark Forest; however, it is considered unlikely to have a significant impact on this EEC.

### Bibliography:

Bell, S. and Driscoll, C. (2007) Vegetation of the Cessnock-Kurri region, Cessnock LGA, New South Wales: Survey, Classification & Mapping. Report to Department of Environment & Climate Change, Newcastle.

NSW Scientific Committee (2005) Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin Bioregion - Endangered ecological community determination - final. DEC (NSW), Sydney.



### c1.3 HUNTER LOWLAND REDGUM FOREST IN THE SYDNEY BASIN AND NSW NORTH COAST BIOREGIONS

#### Hunter Lowland Redgum Forest in the Sydney Basin and New South Wales North Coast Bioregions

|                                         |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
|-----------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Description</b>                      | Hunter Lowland Redgum Forest is an open forest where the most common canopy tree species are <i>Eucalyptus tereticornis</i> (Forest Red Gum) and <i>E. punctata</i> (Grey Gum). Other frequently occurring canopy species are <i>Angophora floribunda</i> (Rough-barked Apple), <i>E. crebra</i> (Narrow-leaved Ironbark), <i>E. moluccana</i> (Grey Box) and <i>Corymbia maculata</i> (Spotted Gum). The shrub layer is open and common shrub species include <i>Breynia oblongifolia</i> (Coffee Bush), <i>Leucopogon juniperinus</i> (Prickly Beard-heath) and <i>Daviesia ulicifolia</i> (Gorse Bitter Pea). The ground cover typically comprises grasses and herbs with common species being <i>Microlaena stipoides</i> var. <i>stipoides</i> (Weeping Meadow Grass), <i>Pratia purpurascens</i> (Whiteroot), <i>Lomandra multiflora</i> (Many-flowered Mat-rush), <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Cheilanthes sieberi</i> (Poison Rock Fern) and <i>Dichondra repens</i> (Kidney Weed). |
| <b>Habitat Requirements and Ecology</b> | Permian sediments of the Hunter Valley floor on gentle slopes of depressions and drainage flats.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| <b>Distribution</b>                     | Occurs between Muswellbrook, Beresfield, Mulbring and Cessnock in the Lower Hunter in the Sydney Basin and North Coast bioregions. It has been recorded from the Maitland, Cessnock, Port Stephens, Muswellbrook and Singleton LGAs.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| <b>BC Act 2016</b>                      | Endangered Ecological Community                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <b>EPBC Act 1999</b>                    | Not listed                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |

#### Five-Part Test of Significance

**The objective of section 7.3 of the Biodiversity Conservation Act 2016 (BC Act), the test of significance, is to provide standardised and transparent consideration of threatened species and ecological communities, and their habitats, through the development assessment process. The following is to be taken into account for the purposes of determining whether a proposed development or activity is likely to significantly affect threatened species or ecological communities, or their habitats:**

- a. *have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction,*

NA

- b. *in the case of an endangered ecological community or critically endangered ecological community, whether the proposed development or activity:*
- is likely to have an adverse effect on the extent of the ecological community such that its local occurrence is likely to be placed at risk of extinction, or*
  - is likely to substantially and adversely modify the composition of the ecological community such that its local occurrence is likely to be placed at risk of extinction,*

Areas mapped as PCT 3446 Lower North Foothills Ironbark-Box-Gum Grassy Forest and PCT 3634 Quorrobolong Sand Flats Forest were found to be consistent with the EEC; Hunter Lowland Redgum Forest in the Sydney Basin and New South Wales North Coast Bioregions. Sections of this EEC were present along present along Scales Road at Loxford and Majors Lane Sawyers Gully. This EEC largely occurred within areas of existing easement, composed of low maintained grown covers with a varying diversity. Better quality areas of Hunter Lowland Redgum Forest were generally present on parts of the Scales Road reserve. The proposal will result in the removal/modification of a relatively small amount of generally disturbed Lower Hunter Spotted Gum Ironbark Forest resulting in an incremental reduction in the local area, however, it is considered unlikely to have a significant impact which might lead to extinction locally.

- c. *in relation to the habitat of a threatened species or ecological community:*
- the extent to which habitat is likely to be removed or modified as a result of the proposed development or activity, and*
  - whether an area of habitat is likely to become fragmented or isolated from other areas of habitat as a result of the proposed development or activity, and*
  - the importance of the habitat to be removed, modified, fragmented or isolated to the long-term survival of the species or ecological community in the locality,*

**Hunter Lowland Redgum Forest in the Sydney Basin and New South Wales North Coast Bioregions**

The REZ is will only have a relatively small impact on Hunter Lowland Redgum Forest within areas of existing easement and parts of the easement edge. No areas are likely to become fragmented or isolated from other areas that are important to the long-term survival of Hunter Lowland Redgum Forest in the local area.

- d. *whether the proposed development or activity is likely to have an adverse effect on any declared area of outstanding biodiversity value (either directly or indirectly),*

No areas declared areas of outstanding biodiversity value were present within proximity to the REZ proposal.

- e. *whether the proposed development or activity is or is part of a key threatening process or is likely to increase the impact of a key threatening process.*

The 'Key Threatening Processes' currently listed under Schedule 4 of the BC Act 2016 that are relevant to the proposal have been addressed below:

**Clearing of Native Vegetation.**

The proposal will result in the removal of native vegetation and may be viewed as being part of this Key Threatening Process. The action is unlikely to be responsible for the significant loss of this EEC provided that recommendations for impact minimisation as listed within Section 6.4 are undertaken.

**Invasion of native plant communities by exotic perennial grasses.**

Specimens of exotic perennial grasses such as *Chloris gayana* (Rhodes Grass), *Eragrostis curvula* (African Love Grass), *Andropogon virginicus* (Whisky Grass) and *Hyparrhenia hirta* (Coolatai Grass) were present. These exotic grasses compete with species within this EEC for space, light, and other resources and may threaten its persistence at several sites. The action is unlikely to be responsible for the significant loss of this EEC provided recommendations for impact minimisation as listed within Section 6.4 are undertaken.

**High frequency fire resulting in the disruption of life cycle processes in plants and animals and loss of vegetation structure and composition**

It is unknown what impact fire has had within areas of this EEC. The proposal is unlikely to increase

**Competition and grazing by the feral European rabbit, *Oryctolagus cuniculus***

Observations and secondary indications of the European Rabbit such as scats and diggings were noted along parts of the REZ. The proposal is not likely to result in an increase in feral numbers of this introduced species.

**Herbivory and environmental degradation caused by feral deer**

Tracks of Feral Deer were recorded within several sections of the REZ. Feral Deer would be considered to have an impact on native flora in the local area. The proposal is not likely to result in an increase in feral numbers of this introduced species.

**Infection of native plants by *Phytophthora cinnamomi***

No evidence of *Phytophthora cinnamomi* was observed during site visits. The proposal is unlikely to cause the spread of this fungus.

**Invasion, establishment and spread of Lantana (*Lantana camara*)**

Lantana was recorded within parts of the REZ proposal. The proposal is unlikely to cause the spread of this weed species.

**Introduction and establishment of Exotic Rust Fungi of the order Pucciniales pathogenic on plants of the family Myrtaceae.**

No evidence of the fungi was observed during site visits. The proposal is unlikely to cause the spread of this fungus.

**Invasion of native plant communities by African Olive *Olea europaea* subsp. *cuspidata***

The African Olive was recorded along parts of the REZ route. The proposal is unlikely to cause further spread of this introduced tree.



## Hunter Lowland Redgum Forest in the Sydney Basin and New South Wales North Coast Bioregions

### **Removal of dead wood and dead trees**

Dead wood and trees were recorded along small sections of the REZ route. Dead wood and trees such be avoided wherever possible.

### **Predation, habitat degradation, competition and disease transmission by *Sus scrofa* (Feral Pigs)**

Signs of Feral Pigs were noted along parts of the REZ route. The proposal is not likely to result in an increase in feral numbers of this introduced species.

### **Conclusion**

The proposal will result in a small incremental reduction in Hunter Lowland Redgum Forest; however, it is considered unlikely to have a significant impact on this EEC.

### **Bibliography:**

Bell, S. and Driscoll, C. (2007) Vegetation of the Cessnock-Kurri region, Cessnock LGA, New South Wales: Survey, Classification & Mapping. Report to Department of Environment & Climate Change, Newcastle.

**c1.4 CENTRAL HUNTER-IRONBARK-SPOTTED GUM GREY BOX FOREST IN THE NSW NORTH COAST AND SYDNEY BASIN BIOREGIONS**

**Central Hunter Ironbark-Spotted Gum-Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions**

|                                         |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|-----------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Description</b>                      | Typically forms an open forest or woodland dominated by <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Corymbia maculata</i> (Spotted Gum), <i>Eucalyptus moluccana</i> (Grey Box). Other tree species such as <i>Eucalyptus fibrosa</i> (Red Ironbark) and <i>Eucalyptus tereticornis</i> (Forest Red Gum) may be present, and occasionally dominate or co-dominate. A sparse layer of small trees including <i>Allocasuarina luehmannii</i> (Bulloak) or <i>Acacia parvipinnula</i> (Silver-stemmed Wattle) may be present in some areas. The shrub layer varies from sparse to moderately dense and includes shrub species include <i>Daviesia ulicifolia</i> subsp. <i>ulicifolia</i> (Gorse Bitter Pea), <i>Pultenaea spinosa</i> (Grey Bush-pea), <i>Breynia oblongifolia</i> (Breynia) and <i>Bursaria spinosa</i> subsp. <i>spinosa</i> (Blackthorn). Common species include <i>Cheilanthes sieberi</i> subsp. <i>sieberi</i> (Mulga Fern), <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Pratia purpurascens</i> (Whiteroot), <i>Lomandra multiflora</i> subsp. <i>multiflora</i> (Many-flowered Mat-rush), <i>Pomax umbellata</i> , <i>Glycine tabacina</i> , <i>Dianella revoluta</i> (Blue Flax-lily), <i>Laxmannia gracilis</i> (Slender Wire Lily), <i>Vernonia cinerea</i> var. <i>cinerea</i> , <i>Desmodium varians</i> (Slender Tick-trefoil) and <i>Dichondra repens</i> (Kidney Weed). |
| <b>Habitat Requirements and Ecology</b> | Occupies undulating country including low rises and slopes, occurring on all aspects. It may also occur on alluvial and colluvial soils in valleys. It mostly occurs on clayey soils found on Permian sediments.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| <b>Distribution</b>                     | Occurs in the central Hunter Valley mainly between Maitland and Muswellbrook. It has been recorded from Singleton, Cessnock and Muswellbrook LGAs.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| <b>BC Act 2016</b>                      | Endangered Ecological Community                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| <b>EPBC Act 1999</b>                    | Not listed                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |

**Five-Part Test of Significance**

**The objective of section 7.3 of the Biodiversity Conservation Act 2016 (BC Act), the test of significance, is to provide standardised and transparent consideration of threatened species and ecological communities, and their habitats, through the development assessment process. The following is to be taken into account for the purposes of determining whether a proposed development or activity is likely to significantly affect threatened species or ecological communities, or their habitats:**

- a. have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction,

NA

- b. in the case of an endangered ecological community or critically endangered ecological community, whether the proposed development or activity:
- is likely to have an adverse effect on the extent of the ecological community such that its local occurrence is likely to be placed at risk of extinction, or
  - is likely to substantially and adversely modify the composition of the ecological community such that its local occurrence is likely to be placed at risk of extinction,

Areas mapped as PCT 3315 Central Hunter Ironbark-Spotted Gum Forest were found to be consistent with the EEC; Central Hunter Ironbark-Spotted Gum-Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions. This EEC was present along narrow sections of easement near Branxton, Belford and Mount Thorley. The proposal will result in the removal/modification of narrow sections (totaling 13.27ha) of Central Hunter Ironbark-Spotted Gum-Grey Box Forest resulting in an incremental reduction in the local area, however, it is considered unlikely to have a significant impact which might lead to extinction locally.

- c. in relation to the habitat of a threatened species or ecological community:
- the extent to which habitat is likely to be removed or modified as a result of the proposed development or activity, and
  - whether an area of habitat is likely to become fragmented or isolated from other areas of habitat as a result of the proposed development or activity, and
  - the importance of the habitat to be removed, modified, fragmented or isolated to the long-term survival of the species or ecological community in the locality,



**Central Hunter Ironbark-Spotted Gum-Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions**

The REZ is will impact 13.24ha of Central Hunter Ironbark-Spotted Gum-Grey Box Forest occurring largely within new sections of easement and parts of the easement edge. No areas are likely to become fragmented or isolated from other areas that are important to the long-term survival of Central Hunter Ironbark-Spotted Gum-Grey Box Forest in the local area.

- d. *whether the proposed development or activity is likely to have an adverse effect on any declared area of outstanding biodiversity value (either directly or indirectly),*

No areas declared areas of outstanding biodiversity value were present within proximity to the REZ proposal.

- e. *whether the proposed development or activity is or is part of a key threatening process or is likely to increase the impact of a key threatening process.*

The 'Key Threatening Processes' currently listed under Schedule 4 of the BC Act 2016 that are relevant to the proposal have been addressed below:

**Clearing of Native Vegetation.**

The proposal will result in the removal of native vegetation and may be viewed as being part of this Key Threatening Process. The action is unlikely to be responsible for the significant loss of this EEC provided that recommendations for impact minimisation as listed within Section 6.4 are undertaken.

**Invasion of native plant communities by exotic perennial grasses.**

Specimens of exotic perennial grasses such as *Chloris gayana* (Rhodes Grass), *Eragrostis curvula* (African Love Grass), *Andropogon virginicus* (Whisky Grass) and *Hyparrhenia hirta* (Coolatai Grass) were present. These exotic grasses compete with species within this EEC for space, light, and other resources and may threaten its persistence at several sites. The action is unlikely to be responsible for the significant loss of this EEC provided recommendations for impact minimisation as listed within Section 6.4 are undertaken.

**High frequency fire resulting in the disruption of life cycle processes in plants and animals and loss of vegetation structure and composition**

It is unknown what impact fire has had within areas of this EEC. The proposal is unlikely to increase

**Competition and grazing by the feral European rabbit, *Oryctolagus cuniculus***

Observations and secondary indications of the European Rabbit such as scats and diggings were noted along parts of the REZ. The proposal is not likely to result in an increase in feral numbers of this introduced species.

**Herbivory and environmental degradation caused by feral deer**

Tracks of Feral Deer were recorded within several sections of the REZ. Feral Deer would be considered to have an impact on native flora in the local area. The proposal is not likely to result in an increase in feral numbers of this introduced species.

**Infection of native plants by *Phytophthora cinnamomi***

No evidence of *Phytophthora cinnamomi* was observed during site visits. The proposal is unlikely to cause the spread of this fungus.

**Invasion, establishment and spread of Lantana (*Lantana camara*)**

Lantana was recorded within parts of the REZ proposal. The proposal is unlikely to cause the spread of this weed species.

**Introduction and establishment of Exotic Rust Fungi of the order Pucciniales pathogenic on plants of the family Myrtaceae.**

No evidence of the fungi was observed during site visits. The proposal is unlikely to cause the spread of this fungus.

**Invasion of native plant communities by African Olive *Olea europaea* subsp. *cuspidata***

## Central Hunter Ironbark-Spotted Gum-Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions

The African Olive was recorded along parts of the REZ route. The proposal is unlikely to cause further spread of this introduced tree.

### Removal of dead wood and dead trees

Dead wood and trees were recorded along small sections of the REZ route. Dead wood and trees such be avoided wherever possible.

### Predation, habitat degradation, competition and disease transmission by *Sus scrofa* (Feral Pigs)

Signs of Feral Pigs were noted along parts of the REZ route. The proposal is not likely to result in an increase in feral numbers of this introduced species.

### Conclusion

The proposal will result in an incremental reduction in Central Hunter Ironbark-Spotted Gum-Grey Box Forest; however, it is considered unlikely to have a significant impact on this EEC.

### Bibliography:

Bell, S. and Driscoll, C. (2007) Vegetation of the Cessnock-Kurri region, Cessnock LGA, New South Wales: Survey, Classification & Mapping. Report to Department of Environment & Climate Change, Newcastle.



## c1.5 SWAMP OAK FLOODPLAIN FOREST OF THE NSW NORTH COAST, SYDNEY BASIN AND SOUTH EAST CORNER BIOREGIONS

### Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions

|                                         |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|-----------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Description</b>                      | Swamp Oak Floodplain Forest is a community of plants that is generally dominated by the tree/s <i>Casuarina glauca</i> (Swamp Oak) and/or <i>Melaleuca ericifolia</i> (Swamp Paperbark). The understorey of this community is characterised by frequent occurrences of vines such as: <i>Parsonsia straminea</i> (Common Silkpod), <i>Geitonoplesium cymosum</i> (Scrambling Lily) and <i>Stephania japonica</i> (Snake Vine). There may be a sparse layer of shrubs and a number of small herbs such as <i>Centella asiatica</i> (Indian Pennywort), <i>Commelina cyanea</i> (Commelina), <i>Persicaria decipiens</i> (Slender Knotweed) and <i>Viola</i> spp.. |
| <b>Habitat Requirements and Ecology</b> | Swamp Oak Floodplain Forest is associated with humic clay and sandy loam soils on waterlogged or periodically flooded areas. These soils are generally deposited during flood events and occur on the flats and drainage lines of the Coastal Floodplain. The community is usually found below 20m in elevation although sometimes up to 50 m elevation on small floodplains or where the larger floodplains adjoin lithic (rocky) substrates or coastal sand plains.                                                                                                                                                                                            |
| <b>Distribution</b>                     | Found in the NSW North Coast, Sydney Basin and South East Corner bioregions.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| <b>BC Act 2016</b>                      | Endangered Ecological Community                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| <b>EPBC Act 1999</b>                    | Not listed                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |

#### Five-Part Test of Significance

**The objective of section 7.3 of the Biodiversity Conservation Act 2016 (BC Act), the test of significance, is to provide standardised and transparent consideration of threatened species and ecological communities, and their habitats, through the development assessment process. The following is to be taken into account for the purposes of determining whether a proposed development or activity is likely to significantly affect threatened species or ecological communities, or their habitats:**

- a. have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction,

NA

- b. in the case of an endangered ecological community or critically endangered ecological community, whether the proposed development or activity:
  - i. is likely to have an adverse effect on the extent of the ecological community such that its local occurrence is likely to be placed at risk of extinction, or
  - ii. is likely to substantially and adversely modify the composition of the ecological community such that its local occurrence is likely to be placed at risk of extinction,

Areas mapped as PCT 4015 Central Hunter Riparian Forest and PCT 4023 Coastal Valley Riparian Forest were found to be consistent with the EEC; Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions. This EEC was present alongside drainage crossings such as Black Creek at Belford and Muds Creek Crossing at Whittingham along the Golden Highway. Swamp Oak Floodplain Forest largely occurred within and either side of areas of existing easement, composed of low maintained grown covers with a varying diversity. Some smaller specimens of trees such as *Casuarina glauca* will likely require removal. The proposal will result in the removal/modification of a relatively small amount of generally disturbed Swamp Oak Floodplain Forest resulting in an incremental reduction in the local area, however, it is considered unlikely to have a significant impact which might lead to extinction locally.

- c. in relation to the habitat of a threatened species or ecological community:
  - i. the extent to which habitat is likely to be removed or modified as a result of the proposed development or activity, and
  - ii. whether an area of habitat is likely to become fragmented or isolated from other areas of habitat as a result of the proposed development or activity, and
  - iii. the importance of the habitat to be removed, modified, fragmented or isolated to the long-term survival of the species or ecological community in the locality,

**Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions**

The REZ is will only have a relatively small impact on Swamp Oak Floodplain Forest occurring largely within areas of existing easement and parts of the easement edge. No areas are likely to become fragmented or isolated from other areas that are important to the long-term survival of Swamp Oak Floodplain Forest in the local area.

- d. whether the proposed development or activity is likely to have an adverse effect on any declared area of outstanding biodiversity value (either directly or indirectly),*

No areas declared areas of outstanding biodiversity value were present within proximity to the REZ proposal.

- e. whether the proposed development or activity is or is part of a key threatening process or is likely to increase the impact of a key threatening process.*

The 'Key Threatening Processes' currently listed under Schedule 4 of the BC Act 2016 that are relevant to the proposal have been addressed below:

***Clearing of Native Vegetation.***

The proposal will result in the removal of native vegetation and may be viewed as being part of this Key Threatening Process. The action is unlikely to be responsible for the significant loss of this EEC provided that recommendations for impact minimisation as listed within Section 6.4 are undertaken.

***Invasion of native plant communities by exotic perennial grasses.***

Specimens of exotic perennial grasses such as *Chloris gayana* (Rhodes Grass), *Eragrostis curvula* (African Love Grass), *Andropogon virginicus* (Whisky Grass) and *Hyparrhenia hirta* (Coolatai Grass) were present. These exotic grasses compete with species within this EEC for space, light, and other resources and may threaten its persistence at several sites. The action is unlikely to be responsible for the significant loss of this EEC provided recommendations for impact minimisation as listed within Section 6.4 are undertaken.

***High frequency fire resulting in the disruption of life cycle processes in plants and animals and loss of vegetation structure and composition***

It is unknown what impact fire has had within areas of this EEC. The proposal is unlikely to increase

***Competition and grazing by the feral European rabbit, *Oryctolagus cuniculus****

Observations and secondary indications of the European Rabbit such as scats and diggings were noted along parts of the REZ. The proposal is not likely to result in an increase in feral numbers of this introduced species.

***Herbivory and environmental degradation caused by feral deer***

Tracks of Feral Deer were recorded within several sections of the REZ. Feral Deer would be considered to have an impact on native flora in the local area. The proposal is not likely to result in an increase in feral numbers of this introduced species.

***Infection of native plants by *Phytophthora cinnamomi****

No evidence of *Phytophthora cinnamomi* was observed during site visits. The proposal is unlikely to cause the spread of this fungus.

***Invasion, establishment and spread of Lantana (*Lantana camara*)***

Lantana was recorded within parts of the REZ proposal. The proposal is unlikely to cause the spread of this weed species.

***Introduction and establishment of Exotic Rust Fungi of the order Pucciniales pathogenic on plants of the family Myrtaceae.***

No evidence of the fungi was observed during site visits. The proposal is unlikely to cause the spread of this fungus.

***Invasion of native plant communities by African Olive *Olea europaea* subsp. *cuspidata****



## Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions

The African Olive was recorded along parts of the REZ route. The proposal is unlikely to cause further spread of this introduced tree.

### Removal of dead wood and dead trees

Dead wood and trees were recorded along small sections of the REZ route. Dead wood and trees such be avoided wherever possible.

### Predation, habitat degradation, competition and disease transmission by *Sus scrofa* (Feral Pigs)

Signs of Feral Pigs were noted along parts of the REZ route. The proposal is not likely to result in an increase in feral numbers of this introduced species.

### Conclusion

The proposal will result in a small incremental reduction in Swamp Oak Floodplain Forest; however, it is considered unlikely to have a significant impact on this EEC.

### Bibliography:

Bell, S. and Driscoll, C. (2007) Vegetation of the Cessnock-Kurri region, Cessnock LGA, New South Wales: Survey, Classification & Mapping. Report to Department of Environment & Climate Change, Newcastle.

**c1.6 CENTRAL HUNTER GREY BOX-IRONBARK WOODLAND IN THE NSW NORTH COAST AND SYDNEY BASIN BIOREGIONS**

**Central Hunter Grey Box-Ironbark Woodland in the New South Wales North Coast and Sydney Basin Bioregions**

|                                         |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|-----------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Description</b>                      | Typically forms a woodland dominated by <i>Eucalyptus crebra</i> (Narrow-leaved Ironbark), <i>Brachychiton populneus</i> subsp. <i>populneus</i> (Kurrajong) and <i>Eucalyptus moluccana</i> (Grey Box). Other tree species such as <i>Angophora floribunda</i> (Rough-barked Apple) and <i>Callitris endlicheri</i> (Black Cypress Pine) may be present and occasionally dominate or co-dominate. A shrub layer is often present and common shrub species include <i>Notelaea microcarpa</i> var. <i>microcarpa</i> (Mock Olive), <i>Breynia oblongifolia</i> (Breynia), <i>Bursaria spinosa</i> subsp. <i>spinosa</i> (Blackthorn), <i>Cassinia quinquefaria</i> and <i>Dodonaea viscosa</i> (Sticky Hopbush). Subshrubs may also be common and include <i>Solanum cinereum</i> (Narrawa Burr), <i>Phyllanthus virgatus</i> and <i>Maireana microphylla</i> (Small-leaf Bluebush). Ground cover can be moderately dense to dense and consist of numerous forbs and grass species as well as a small number of ferns, sedges and twiners. The more common species include <i>Cymbopogon refractus</i> (Barbed Wire Grass), <i>Aristida ramosa</i> , <i>Dichondra repens</i> (Kidney Weed), <i>Cheilanthes sieberi</i> subsp. <i>sieberi</i> (Mulga Fern), <i>Cheilanthes distans</i> (Bristly Cloak Fern), <i>Chloris ventricosa</i> (Tall Windmill Grass), <i>Desmodium varians</i> (Slender Tick-trefoil), <i>Calotis lappulacea</i> (Yellow Burr-daisy), <i>Lomandra multiflora</i> subsp. <i>multiflora</i> (Many-flowered Mat-rush), <i>Brunoniella australis</i> (Blue Trumpet) and <i>Glycine tabacina</i> .. |
| <b>Habitat Requirements and Ecology</b> | Occurs in areas of relatively low rainfall and high temperatures. It is associated mostly with Permian lithology, and is situated on gently undulating hills, slopes and valleys, or occasionally on rocky knolls.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <b>Distribution</b>                     | Occurs in the Central Hunter Valley between about Singleton and Muswellbrook. It is known to occur in the Cessnock, Singleton and Muswellbrook LGAs.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| <b>BC Act 2016</b>                      | Endangered Ecological Community                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| <b>EPBC Act 1999</b>                    | Not listed                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |

**Five-Part Test of Significance**

***The objective of section 7.3 of the Biodiversity Conservation Act 2016 (BC Act), the test of significance, is to provide standardised and transparent consideration of threatened species and ecological communities, and their habitats, through the development assessment process. The following is to be taken into account for the purposes of determining whether a proposed development or activity is likely to significantly affect threatened species or ecological communities, or their habitats:***

- a. *have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction,*

NA

- b. *in the case of an endangered ecological community or critically endangered ecological community, whether the proposed development or activity:*
- is likely to have an adverse effect on the extent of the ecological community such that its local occurrence is likely to be placed at risk of extinction, or*
  - is likely to substantially and adversely modify the composition of the ecological community such that its local occurrence is likely to be placed at risk of extinction,*

Areas mapped as PCT 3431 Central Hunter Ironbark Grassy Woodland were found to be consistent with the EEC; Central Hunter Grey Box-Ironbark Woodland in the New South Wales North Coast and Sydney Basin Bioregions. This EEC was present within and around the Muswellbrook STS where it consisted of areas of woodland and derived grassland. The proposal will result in the removal/modification of Central Hunter Grey Box-Ironbark Woodland resulting in an incremental reduction in the local area; however, it is considered unlikely to have a significant impact which might lead to extinction locally.

- c. *in relation to the habitat of a threatened species or ecological community:*
- the extent to which habitat is likely to be removed or modified as a result of the proposed development or activity, and*
  - whether an area of habitat is likely to become fragmented or isolated from other areas of habitat as a result of the proposed development or activity, and*
  - the importance of the habitat to be removed, modified, fragmented or isolated to the long-term survival of the species or ecological community in the locality,*



**Central Hunter Grey Box-Ironbark Woodland in the New South Wales North Coast and Sydney Basin Bioregions**

The REZ will impact 4.69ha of Central Hunter Grey Box-Ironbark Woodland occurring within the proposed Muswellbrook STS and required associated easements was composed of woodland and derived grassland. Will require the direct removal of 2.99ha (PCT 3431 1.27ha & PCT 3431-derived grassland 1.72ha) for the Muswellbrook STS and 1.70ha (PCT 3431 0.66ha & PCT 3431-derived grassland 1.04ha) for the associated easements. No areas are likely to become fragmented or isolated from other areas that are important to the long-term survival of Central Hunter Grey Box-Ironbark Woodland in the local area.

- d. *whether the proposed development or activity is likely to have an adverse effect on any declared area of outstanding biodiversity value (either directly or indirectly),*

No areas declared areas of outstanding biodiversity value were present within proximity to the REZ proposal.

- e. *whether the proposed development or activity is or is part of a key threatening process or is likely to increase the impact of a key threatening process.*

The 'Key Threatening Processes' currently listed under Schedule 4 of the BC Act 2016 that are relevant to the proposal have been addressed below:

**Clearing of Native Vegetation.**

The proposal will result in the removal of native vegetation and may be viewed as being part of this Key Threatening Process. The action is unlikely to be responsible for the significant loss of this EEC provided that recommendations for impact minimisation as listed within Section 6.4 are undertaken.

**Invasion of native plant communities by exotic perennial grasses.**

Specimens of exotic perennial grasses such as *Chloris gayana* (Rhodes Grass), *Eragrostis curvula* (African Love Grass), *Andropogon virginicus* (Whisky Grass) and *Hyparrhenia hirta* (Coolatai Grass) were present. These exotic grasses compete with species within this CEEC for space, light, and other resources and may threaten its persistence at several sites. The action is unlikely to be responsible for the significant loss of this EEC provided recommendations for impact minimisation as listed within Section 6.4 are undertaken.

**High frequency fire resulting in the disruption of life cycle processes in plants and animals and loss of vegetation structure and composition**

It is unknown what impact fire has had within areas of this EEC. The proposal is unlikely to increase

**Competition and grazing by the feral European rabbit, *Oryctolagus cuniculus***

Observations and secondary indications of the European Rabbit such as scats and diggings were noted along parts of the REZ. The proposal is not likely to result in an increase in feral numbers of this introduced species.

**Herbivory and environmental degradation caused by feral deer**

Tracks of Feral Deer were recorded within several sections of the REZ. Feral Deer would be considered to have an impact on native flora in the local area. The proposal is not likely to result in an increase in feral numbers of this introduced species.

**Infection of native plants by *Phytophthora cinnamomi***

No evidence of *Phytophthora cinnamomi* was observed during site visits. The proposal is unlikely to cause the spread of this fungus.

**Invasion, establishment and spread of Lantana (*Lantana camara*)**

Lantana was recorded within parts of the REZ proposal. The proposal is unlikely to cause the spread of this weed species.

**Introduction and establishment of Exotic Rust Fungi of the order Pucciniales pathogenic on plants of the family Myrtaceae.**

No evidence of the fungi was observed during site visits. The proposal is unlikely to cause the spread of this fungus.

## Central Hunter Grey Box-Ironbark Woodland in the New South Wales North Coast and Sydney Basin Bioregions

### **Invasion of native plant communities by African Olive *Olea europaea* subsp. *cuspidata***

The African Olive was recorded along parts of the REZ route. The proposal is unlikely to cause further spread of this introduced tree.

### **Removal of dead wood and dead trees**

Dead wood and trees were recorded along small sections of the REZ route. Dead wood and trees such be avoided wherever possible.

### **Predation, habitat degradation, competition and disease transmission by *Sus scrofa* (Feral Pigs)**

Signs of Feral Pigs were noted along parts of the REZ route. The proposal is not likely to result in an increase in feral numbers of this introduced species.

### **Conclusion**

The proposal will result in a small incremental reduction in Central Hunter Grey Box-Ironbark Woodland, it is considered unlikely to have a significant impact on this EEC.

### **Bibliography:**

Bell, S. and Driscoll, C. (2007) Vegetation of the Cessnock-Kurri region, Cessnock LGA, New South Wales: Survey, Classification & Mapping. Report to Department of Environment & Climate Change, Newcastle.



## c1.7 WHITE BOX-YELLOW BOX-BLAKELY'S RED GUM GRASSY WOODLAND AND DERIVED NATIVE GRASSLAND

### White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grassland

|                                         |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|-----------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Description</b>                      | Characterised by widely-spaced trees and projected foliage cover generally less than 30%. Conversely, the canopy may be completely absent in areas of derived native grassland where tree removal has occurred, and in such areas higher abundance of groundcover species may be present. Dominated by one or more of the species <i>Eucalyptus albens</i> (White Box), <i>E. melliodora</i> (Yellow Box) and <i>E. blakelyi</i> (Blakely's Red Gum). <i>Eucalyptus moluccana</i> (Grey Box) may be codominant in the Nandewar Bioregion and in the north-western corner of the Sydney Basin Bioregion in the upper Hunter Valley. Understorey shrubs are typically sparse or absent. The groundcover is dominated by perennial tussock grasses interspersed with a diverse range of forb species with the families Asteraceae and Fabaceae, and the orders Liliales and Asparagales well represented. |
| <b>Habitat Requirements and Ecology</b> | Occurs on hilly to undulating landscapes in areas with soils of moderate fertility derived from a range of lithologies, including alkaline and acid volcanics, granites, sediments, serpentinites and metamorphics.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <b>Distribution</b>                     | Occurs in an arc along the western slopes and tablelands of the Great Dividing Range from Southern Queensland through NSW to central Victoria. It occurs in the Brigalow Belt South, Nandewar, New England Tableland, South Eastern Queensland, Sydney Basin, NSW North Coast, South Eastern Highlands, South East Corner, NSW South Western Slopes, Victorian Midlands and Riverina Bioregions.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| <b>BC Act 2016</b>                      | Critically Endangered Ecological Community                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| <b>EPBC Act 1999</b>                    | Critically Endangered Ecological Community                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |

#### Five-Part Test of Significance

*The objective of section 7.3 of the Biodiversity Conservation Act 2016 (BC Act), the test of significance, is to provide standardised and transparent consideration of threatened species and ecological communities, and their habitats, through the development assessment process. The following is to be taken into account for the purposes of determining whether a proposed development or activity is likely to significantly affect threatened species or ecological communities, or their habitats:*

- a. *have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction,*

NA

- b. *in the case of an endangered ecological community or critically endangered ecological community, whether the proposed development or activity:*
- is likely to have an adverse effect on the extent of the ecological community such that its local occurrence is likely to be placed at risk of extinction, or*
  - is likely to substantially and adversely modify the composition of the ecological community such that its local occurrence is likely to be placed at risk of extinction,*

Areas mapped as PCT 3525 Upper Hunter Box-Blakelys Red Gum Grassy Forest were found to be consistent with the CEEC; White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grassland. This CEEC was present along a section of existing easement to the north-east of the Muswellbrook STS. This CEEC largely occurred within areas of existing easement, composed of low maintained grown covers with a varying diversity (derived grassland). Better quality areas of White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grassland were generally present on the edges of the existing easement. The proposal will result in the removal/modification of a relatively small amount of generally disturbed White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grassland resulting in an incremental reduction in the local area; however, it is considered unlikely to have a significant impact which might lead to extinction locally.

- c. *in relation to the habitat of a threatened species or ecological community:*
- the extent to which habitat is likely to be removed or modified as a result of the proposed development or activity, and*
  - whether an area of habitat is likely to become fragmented or isolated from other areas of habitat as a result of the proposed development or activity, and*

**White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grassland**

- iii. *the importance of the habitat to be removed, modified, fragmented or isolated to the long-term survival of the species or ecological community in the locality,*

The REZ is will only have a relatively small impact on White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grassland occurring largely within areas of existing easement and parts of the easement edge. No areas are likely to become further fragmented or isolated from other areas that are important to the long-term survival of White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the local area.

- d. *whether the proposed development or activity is likely to have an adverse effect on any declared area of outstanding biodiversity value (either directly or indirectly),*

No areas declared areas of outstanding biodiversity value were present within proximity to the REZ proposal.

- e. *whether the proposed development or activity is or is part of a key threatening process or is likely to increase the impact of a key threatening process.*

The 'Key Threatening Processes' currently listed under Schedule 4 of the BC Act 2016 that are relevant to the proposal have been addressed below:

***Clearing of Native Vegetation.***

The proposal will result in the removal of native vegetation and may be viewed as being part of this Key Threatening Process. The action is unlikely to be responsible for the significant loss of this EEC provided that recommendations for impact minimisation as listed within Section 6.4 are undertaken.

***Invasion of native plant communities by exotic perennial grasses.***

Specimens of exotic perennial grasses such as *Chloris gayana* (Rhodes Grass), *Eragrostis curvula* (African Love Grass), *Andropogon virginicus* (Whisky Grass) and *Hyparrhenia hirta* (Coolatai Grass) were present. These exotic grasses compete with species within this CEEC for space, light, and other resources and may threaten its persistence at several sites. The action is unlikely to be responsible for the significant loss of this EEC provided recommendations for impact minimisation as listed within Section 6.4 are undertaken.

***High frequency fire resulting in the disruption of life cycle processes in plants and animals and loss of vegetation structure and composition***

It is unknown what impact fire has had within areas of this EEC. The proposal is unlikely to increase

***Competition and grazing by the feral European rabbit, *Oryctolagus cuniculus****

Observations and secondary indications of the European Rabbit such as scats and diggings were noted along parts of the REZ. The proposal is not likely to result in an increase in feral numbers of this introduced species.

***Herbivory and environmental degradation caused by feral deer***

Tracks of Feral Deer were recorded within several sections of the REZ. Feral Deer would be considered to have an impact on native flora in the local area. The proposal is not likely to result in an increase in feral numbers of this introduced species.

***Infection of native plants by *Phytophthora cinnamomi****

No evidence of *Phytophthora cinnamomi* was observed during site visits. The proposal is unlikely to cause the spread of this fungus.

***Invasion, establishment and spread of Lantana (*Lantana camara*)***

Lantana was recorded within parts of the REZ proposal. The proposal is unlikely to cause the spread of this weed species.

***Introduction and establishment of Exotic Rust Fungi of the order Pucciniales pathogenic on plants of the family Myrtaceae.***

No evidence of the fungi was observed during site visits. The proposal is unlikely to cause the spread of this fungus.



**White Box – Yellow Box – Blakely’s Red Gum Grassy Woodland and Derived Native Grassland**

**Invasion of native plant communities by African Olive *Olea europaea* subsp. *cuspidata***

The African Olive was recorded along parts of the REZ route. The proposal is unlikely to cause further spread of this introduced tree.

**Removal of dead wood and dead trees**

Dead wood and trees were recorded along small sections of the REZ route. Dead wood and trees such be avoided wherever possible.

**Predation, habitat degradation, competition and disease transmission by *Sus scrofa* (Feral Pigs)**

Signs of Feral Pigs were noted along parts of the REZ route. The proposal is not likely to result in an increase in feral numbers of this introduced species.

**Conclusion**

The proposal will result in a small incremental reduction in White Box – Yellow Box – Blakely’s Red Gum Grassy Woodland and Derived Native Grassland; however, it is considered unlikely to have a significant impact on this EEC.

**Bibliography:**

Bell, S. and Driscoll, C. (2007) Vegetation of the Cessnock-Kurri region, Cessnock LGA, New South Wales: Survey, Classification & Mapping. Report to Department of Environment & Climate Change, Newcastle.

## C2.0 SECTION 7.3 OF THE BC ACT 2016 TEST OF SIGNIFICANCE (THREATENED SPECIES)

### c2.1 *DIURIS TRICOLOR* PINE DONKEY ORCHID

| Diuris tricolor Pine Donkey Orchid                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Description</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         | A tuberous, deciduous terrestrial orchid that has between one and three leaves, to 30 centimetres long and 4mm wide. The flower stalk is between 20-40cm high and has 2-6 flowers, which are bright yellow to orange, speckled with red to purple and white markings. The sepals (the down-pointing slender green segments) are very long and often crossed. The species is a tuberous, deciduous terrestrial orchid and the flowers have a pleasant, light sweet scent.                                                                                                                                                                                                         |
| <b>Habitat Requirements and Ecology</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | Grows in sclerophyll forest among grass, often with native Cypress Pine ( <i>Callitris</i> spp.). It is found in sandy soils, either on flats or small rises. Also recorded from a red earth soil in a Bimble Box community in western NSW. Usually recorded as common and locally frequent in populations, however only one or two plants have also been observed at sites. The species has also been noted as growing in large colonies. Usually, flowers between early September to late October.                                                                                                                                                                             |
| <b>Distribution</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        | Sporadically distributed on the western slopes of NSW, extending from south of Narrandera all the way to the north of NSW. Localities in the south include several sites west of Wagga Wagga, Dubbo in the Central West and Pilliga National in the north. The population in the Muswellbrook LGA is at the eastern limit of the geographic range of the species. All other populations of the species are located west of the Great Dividing Range. The distance of the Muswellbrook LGA population to the nearest population of the species to the west is c. 100 km. Therefore, the population in the Muswellbrook LGA is disjunct and at the limits of its geographic range. |
| <b>BC Act 2016</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         | Vulnerable                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| <b>BC Act 2016</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         | Endangered population listing - <i>Diuris tricolor</i> , the Pine Donkey Orchid population in the Muswellbrook local government area -                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| <b>EPBC Act 1999</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       | Not Listed                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| <p><b><u>Five-Part Test of Significance</u></b></p> <p><b><i>The objective of section 7.3 of the Biodiversity Conservation Act 2016 (BC Act), the test of significance, is to provide standardised and transparent consideration of threatened species and ecological communities, and their habitats, through the development assessment process. The following is to be taken into account for the purposes of determining whether a proposed development or activity is likely to significantly affect threatened species or ecological communities, or their habitats:</i></b></p> <p><i>a. have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction,</i></p> <p>Approximately four specimens of <i>Diuris tricolor</i> (Pine Donkey Orchid) were recorded within the existing easement to the NE of the Muswellbrook STS during fieldwork. As these specimens of <i>D. tricolor</i> occur within the existing easement, they are not required to be impacted. It is recommended that these orchids be avoided during line works along the easement. Given the recommendation the proposal is unlikely to result in the extinction of any local population of this species.</p> <p><i>b. in the case of an endangered ecological community or critically endangered ecological community, whether the proposed development or activity:</i></p> <p><i>i. is likely to have an adverse effect on the extent of the ecological community such that its local occurrence is likely to be placed at risk of extinction, or</i></p> <p><i>ii. is likely to substantially and adversely modify the composition of the ecological community such that its local occurrence is likely to be placed at risk of extinction,</i></p> <p>NA</p> <p><i>c. in relation to the habitat of a threatened species or ecological community:</i></p> <p><i>i. the extent to which habitat is likely to be removed or modified as a result of the proposed development or activity, and</i></p> <p><i>ii. whether an area of habitat is likely to become fragmented or isolated from other areas of habitat as a result of the proposed development or activity, and</i></p> |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |



**Diuris tricolor Pine Donkey Orchid**

- iii. *the importance of the habitat to be removed, modified, fragmented or isolated to the long-term survival of the species or ecological community in the locality,*

Given specimens of *D. tricolor* were found in an existing easement no areas are likely to become further fragmented or isolated from other areas that are important to the long-term survival of this threatened orchid species in the local area.

- d. *whether the proposed development or activity is likely to have an adverse effect on any declared area of outstanding biodiversity value (either directly or indirectly),*

No areas declared areas of outstanding biodiversity value were present within proximity to the REZ proposal.

- e. *whether the proposed development or activity is or is part of a key threatening process or is likely to increase the impact of a key threatening process.*

The 'Key Threatening Processes' currently listed under Schedule 4 of the BC Act 2016 that are relevant to the proposal have been addressed below:

**Clearing of Native Vegetation.**

The proposal will result in the removal of native vegetation and may be viewed as being part of this Key Threatening Process. The action is unlikely to be responsible for the significant loss of this EEC provided that recommendations for impact minimisation as listed within Section 6.4 are undertaken.

**Invasion of native plant communities by exotic perennial grasses.**

Specimens of exotic perennial grasses such as *Chloris gayana* (Rhodes Grass), *Eragrostis curvula* (African Love Grass), *Andropogon virginicus* (Whisky Grass) and *Hyparrhenia hirta* (Coolatai Grass) were present. These exotic grasses compete with species within this CEEC for space, light, and other resources and may threaten its persistence at several sites. The action is unlikely to be responsible for the significant loss of this EEC provided recommendations for impact minimisation as listed within Section 6.4 are undertaken.

**High frequency fire resulting in the disruption of life cycle processes in plants and animals and loss of vegetation structure and composition**

It is unknown what impact fire has had within areas of this EEC. The proposal is unlikely to increase

**Competition and grazing by the feral European rabbit, *Oryctolagus cuniculus***

Observations and secondary indications of the European Rabbit such as scats and diggings were noted along parts of the REZ. The proposal is not likely to result in an increase in feral numbers of this introduced species.

**Herbivory and environmental degradation caused by feral deer**

Tracks of Feral Deer were recorded within several sections of the REZ. Feral Deer would be considered to have an impact on native flora in the local area. The proposal is not likely to result in an increase in feral numbers of this introduced species.

**Infection of native plants by *Phytophthora cinnamomi***

No evidence of *Phytophthora cinnamomi* was observed during site visits. The proposal is unlikely to cause the spread of this fungus.

**Invasion, establishment and spread of Lantana (*Lantana camara*)**

Lantana was recorded within parts of the REZ proposal. The proposal is unlikely to cause the spread of this weed species.

**Introduction and establishment of Exotic Rust Fungi of the order Pucciniales pathogenic on plants of the family Myrtaceae.**

No evidence of the fungi was observed during site visits. The proposal is unlikely to cause the spread of this fungus.

**Invasion of native plant communities by African Olive *Olea europaea* subsp. *cuspidata***

#### Diuris tricolor Pine Donkey Orchid

The African Olive was recorded along parts of the REZ route. The proposal is unlikely to cause further spread of this introduced tree.

#### Removal of dead wood and dead trees

Dead wood and trees were recorded along small sections of the REZ route. Dead wood and trees such be avoided wherever possible.

#### Predation, habitat degradation, competition and disease transmission by *Sus scrofa* (Feral Pigs)

Signs of Feral Pigs were noted along parts of the REZ route. The proposal is not likely to result in an increase in feral numbers of this introduced species.

#### Conclusion

The proposal will result in a small incremental modification of suitable habitat for *D. tricolor* however, it is considered unlikely to have a significant impact on this threatened orchid species.

#### Bibliography:

Jones, D.,J. (2006). *A Complete Guide to Native Orchids of Australia including the Island Territories*. New Holland Publishers Australia Pty Ltd.

### c2.2 GREVILLEA PARVIFLORA SUBSP. PARVIFLORA SMALL-FLOWERED GREVILLEA

#### *Grevillea parviflora* subsp. *parviflora* (Small-flowered Grevillea)

|                                         |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
|-----------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Description</b>                      | Is a low spreading to erect shrub usually less than a metre high. Its erect narrow leaves are 2-3.5 mm long and less than 1.3mm wide with silky hairs on the underside and a short-pointed tip. Leaf margins are curved back, or even rolled completely under. The small flowers are spider-like and clustered in groups of 6-12. Flowers are small and white or pinkish, with rusty-brown hairs that can become red with age. The fruiting capsule is 8-10 mm long and has 1-2 seeds (NSW NPWS 2002h, NSW OEH 2013o).                                                                                                                     |
| <b>Habitat Requirements and Ecology</b> | Grows in sandy or light clay soils usually over thin shales and occurs in a range of vegetation types from heath and shrubby woodland to open forest. It is found over a range of altitudes from flat, low-lying areas to upper slopes and ridge crests, and commonly occurs in open, slightly disturbed sites such as along tracks. Plants are capable of suckering from a rootstock and most populations demonstrate a degree of vegetative spread, particularly after disturbance such as fire. Flowering has been recorded between July to December as well as April-May. Flowers are insect-pollinated and seed dispersal is limited. |
| <b>Distribution</b>                     | Sporadically distributed throughout the Sydney Basin with the main occurrence centred around Picton, Appin and Bargo (and possibly further south to the Moss Vale area). Separate populations are also known further north from Putty to Wyong and Lake Macquarie on the Central Coast and Cessnock and Kurri Kurri in the Lower Hunter.                                                                                                                                                                                                                                                                                                   |
| <b>BC Act 2016</b>                      | Vulnerable                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| <b>EPBC Act 1999</b>                    | Vulnerable                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |

#### Five-Part Test of Significance

**The objective of section 7.3 of the Biodiversity Conservation Act 2016 (BC Act), the test of significance, is to provide standardised and transparent consideration of threatened species and ecological communities, and their habitats, through the development assessment process. The following is to be taken into account for the purposes of determining whether a proposed development or activity is likely to significantly affect threatened species or ecological communities, or their habitats:**

- have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction,



***Grevillea parviflora* subsp. *parviflora* (Small-flowered Grevillea)**

A small number of specimens of *Grevillea parviflora* subsp. *parviflora* were present within sections of existing easement west of the Kurri STS to Graham Road, Sawyers Gully. As these specimens of *G. parviflora* subsp. *parviflora* occur within the existing sections of easement, they are not required to be impacted. It is recommended that these grevilleas be avoided during upgrading works along the easement. Given the recommendation the proposal is unlikely to result in the extinction of any local population of this species.

- b. *in the case of an endangered ecological community or critically endangered ecological community, whether the proposed development or activity:*
- i. *is likely to have an adverse effect on the extent of the ecological community such that its local occurrence is likely to be placed at risk of extinction, or*
  - ii. *is likely to substantially and adversely modify the composition of the ecological community such that its local occurrence is likely to be placed at risk of extinction,*

NA

- c. *in relation to the habitat of a threatened species or ecological community:*
- i. *the extent to which habitat is likely to be removed or modified as a result of the proposed development or activity, and*
  - ii. *whether an area of habitat is likely to become fragmented or isolated from other areas of habitat as a result of the proposed development or activity, and*
  - iii. *the importance of the habitat to be removed, modified, fragmented or isolated to the long-term survival of the species or ecological community in the locality,*

Given specimens of *G. parviflora* subsp. *parviflora* were found in an existing easement no areas are likely to become further fragmented or isolated from other areas that are important to the long-term survival of this threatened grevillea species in the local area.

- d. *whether the proposed development or activity is likely to have an adverse effect on any declared area of outstanding biodiversity value (either directly or indirectly),*

No areas declared areas of outstanding biodiversity value were present within proximity to the REZ proposal.

- e. *whether the proposed development or activity is or is part of a key threatening process or is likely to increase the impact of a key threatening process.*

The 'Key Threatening Processes' currently listed under Schedule 4 of the BC Act 2016 that are relevant to the proposal have been addressed below:

***Clearing of Native Vegetation.***

The proposal will result in the removal of native vegetation and may be viewed as being part of this Key Threatening Process. The action is unlikely to be responsible for the significant loss of this EEC provided that recommendations for impact minimisation as listed within Section 6.4 are undertaken.

***Invasion of native plant communities by exotic perennial grasses.***

Specimens of exotic perennial grasses such as *Chloris gayana* (Rhodes Grass), *Eragrostis curvula* (African Love Grass), *Andropogon virginicus* (Whisky Grass) and *Hyparrhenia hirta* (Coolatai Grass) were present. These exotic grasses compete with species within this CEEC for space, light, and other resources and may threaten its persistence at several sites. The action is unlikely to be responsible for the significant loss of this EEC provided recommendations for impact minimisation as listed within Section 6.4 are undertaken.

***High frequency fire resulting in the disruption of life cycle processes in plants and animals and loss of vegetation structure and composition***

It is unknown what impact fire has had within areas of this EEC. The proposal is unlikely to increase

***Competition and grazing by the feral European rabbit, *Oryctolagus cuniculus****

Observations and secondary indications of the European Rabbit such as scats and diggings were noted along parts of the REZ. The proposal is not likely to result in an increase in feral numbers of this introduced species.

***Grevillea parviflora* subsp. *parviflora* (Small-flowered Grevillea)**

***Herbivory and environmental degradation caused by feral deer***

Tracks of Feral Deer were recorded within several sections of the REZ. Feral Deer would be considered to have an impact on native flora in the local area. The proposal is not likely to result in an increase in feral numbers of this introduced species.

***Infection of native plants by *Phytophthora cinnamomi****

No evidence of *Phytophthora cinnamomi* was observed during site visits. The proposal is unlikely to cause the spread of this fungus.

***Invasion, establishment and spread of Lantana (*Lantana camara*)***

Lantana was recorded within parts of the REZ proposal. The proposal is unlikely to cause the spread of this weed species.

***Introduction and establishment of Exotic Rust Fungi of the order Pucciniales pathogenic on plants of the family Myrtaceae.***

No evidence of the fungi was observed during site visits. The proposal is unlikely to cause the spread of this fungus.

***Invasion of native plant communities by African Olive *Olea europaea* subsp. *cuspidata****

The African Olive was recorded along parts of the REZ route. The proposal is unlikely to cause further spread of this introduced tree.

***Removal of dead wood and dead trees***

Dead wood and trees were recorded along small sections of the REZ route. Dead wood and trees such be avoided wherever possible.

***Predation, habitat degradation, competition and disease transmission by *Sus scrofa* (Feral Pigs)***

Signs of Feral Pigs were noted along parts of the REZ route. The proposal is not likely to result in an increase in feral numbers of this introduced species.

**Conclusion**

The proposal will result in a small incremental modification of suitable habitat for *G. parviflora* subsp. *parviflora* however, it is considered unlikely to have a significant impact on this threatened grevillea species.

**Bibliography:**

Bell, S.A.J. (2004). Vegetation of Werakata National Park, Hunter Valley, New South Wales. *Cunninghamia*. 8(3):331-347.

Driscoll, C. (2013). The ecology and biology of *Grevillea parviflora* subsp. *parviflora*: A review. Report prepared for Lake Macquarie City Council.

Harden, G.J. (ed.) (2002) *Flora of New South Wales Volume 2 Revised Edition*. UNSW, Sydney.

Lake Macquarie City Council (2013). Interim Lake Macquarie *Grevillea parviflora* subsp. *parviflora* Planning and Management Guidelines. Lake Macquarie City Council June 2013.

Mamott, I. (2013). *Notes on a Threatened (Vulnerable) flora species on the NSW Mid North Coast: Small flower Grevillea (*Grevillea parviflora* subsp. *parviflora*)*. Consulting Ecology V30:37-39.

NSW Scientific Committee (1998) *Grevillea parviflora* subsp. *parviflora* (a shrub) - Vulnerable species determination - final. DEC (NSW), Sydney.



## c2.3 EUCALYPTUS GLAUCINA (SLATY RED GUM)

| <i>Eucalyptus glaucina</i> (Slaty Red Gum) |                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|--------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Description</b>                         | A medium-sized tree to 30 m tall. The bark is smooth and mottled white to slaty grey. The juvenile leaves are oval in shape and blue-green with a whitish bloom, and the buds and fruit are similarly coloured. The flowers are white, or occasionally pink, and are produced between August and December. The fruits are oval-shaped and 7–10 mm long. The three to five raised valves are surrounded by a domed disk raised above the fruit. |
| <b>Habitat Requirements and Ecology</b>    | Grows in grassy woodland and dry eucalypt forest on deep, moderately fertile and well-watered soils.                                                                                                                                                                                                                                                                                                                                           |
| <b>Distribution</b>                        | Found in separate districts along the eastern seaboard of NSW, from near Casino, to Taree, south to Broke, and recently discovered on the eastern side of the Blue Mountains National Park near Warragamba Dam.                                                                                                                                                                                                                                |
| <b>BC Act 2016</b>                         | Vulnerable                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| <b>EPBC Act 1999</b>                       | Vulnerable                                                                                                                                                                                                                                                                                                                                                                                                                                     |

**Five-Part Test of Significance**

*The objective of section 7.3 of the Biodiversity Conservation Act 2016 (BC Act), the test of significance, is to provide standardised and transparent consideration of threatened species and ecological communities, and their habitats, through the development assessment process. The following is to be taken into account for the purposes of determining whether a proposed development or activity is likely to significantly affect threatened species or ecological communities, or their habitats:*

- a. *have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction,*

A small number of specimens of *Eucalyptus glaucina* (Slaty Red Gum) were present within close proximity of the proposed and existing easement around Belford and Whittingham. The majority of these specimens will not be required to be impacted. It is recommended that specimens of *E. glaucina* be avoided wherever possible within the scope of the proposal. Given the recommendation the proposal is unlikely to result in the extinction of any local population of this eucalypt species.

- b. *in the case of an endangered ecological community or critically endangered ecological community, whether the proposed development or activity:*
- is likely to have an adverse effect on the extent of the ecological community such that its local occurrence is likely to be placed at risk of extinction, or*
  - is likely to substantially and adversely modify the composition of the ecological community such that its local occurrence is likely to be placed at risk of extinction,*

NA

- c. *in relation to the habitat of a threatened species or ecological community:*
- the extent to which habitat is likely to be removed or modified as a result of the proposed development or activity, and*
  - whether an area of habitat is likely to become fragmented or isolated from other areas of habitat as a result of the proposed development or activity, and*
  - the importance of the habitat to be removed, modified, fragmented or isolated to the long-term survival of the species or ecological community in the locality,*

Specimens of *E. glaucina* are unlikely to become further fragmented or isolated from other areas that are important to the long-term survival of this threatened eucalypt species in the local area.

- d. *whether the proposed development or activity is likely to have an adverse effect on any declared area of outstanding biodiversity value (either directly or indirectly),*

No areas declared areas of outstanding biodiversity value were present within proximity to the proposal.

- e. *whether the proposed development or activity is or is part of a key threatening process or is likely to increase the impact of a key threatening process.*

***Eucalyptus glaucina* (Slaty Red Gum)**

The 'Key Threatening Processes' currently listed under Schedule 4 of the BC Act 2016 that are relevant to the proposal have been addressed below:

***Clearing of Native Vegetation.***

The proposal will result in the removal of native vegetation and may be viewed as being part of this Key Threatening Process. The action is unlikely to be responsible for the significant loss of this EEC provided that recommendations for impact minimisation as listed within Section 6.4 are undertaken.

***Invasion of native plant communities by exotic perennial grasses.***

Specimens of exotic perennial grasses such as *Chloris gayana* (Rhodes Grass), *Eragrostis curvula* (African Love Grass), *Andropogon virginicus* (Whisky Grass) and *Hyparrhenia hirta* (Coolatai Grass) were present. These exotic grasses compete with species within this CEEC for space, light, and other resources and may threaten its persistence at several sites. The action is unlikely to be responsible for the significant loss of this EEC provided recommendations for impact minimisation as listed within Section 6.4 are undertaken.

***High frequency fire resulting in the disruption of life cycle processes in plants and animals and loss of vegetation structure and composition***

It is unknown what impact fire has had within areas of this EEC. The proposal is unlikely to increase

***Competition and grazing by the feral European rabbit, *Oryctolagus cuniculus****

Observations and secondary indications of the European Rabbit such as scats and diggings were noted along parts of the REZ. The proposal is not likely to result in an increase in feral numbers of this introduced species.

***Herbivory and environmental degradation caused by feral deer***

Tracks of Feral Deer were recorded within several sections of the REZ. Feral Deer would be considered to have an impact on native flora in the local area. The proposal is not likely to result in an increase in feral numbers of this introduced species.

***Infection of native plants by *Phytophthora cinnamomi****

No evidence of *Phytophthora cinnamomi* was observed during site visits. The proposal is unlikely to cause the spread of this fungus.

***Invasion, establishment and spread of Lantana (*Lantana camara*)***

Lantana was recorded within parts of the REZ proposal. The proposal is unlikely to cause the spread of this weed species.

***Introduction and establishment of Exotic Rust Fungi of the order Pucciniales pathogenic on plants of the family Myrtaceae.***

No evidence of the fungi was observed during site visits. The proposal is unlikely to cause the spread of this fungus.

***Invasion of native plant communities by African Olive *Olea europaea* subsp. *cuspidata****

The African Olive was recorded along parts of the REZ route. The proposal is unlikely to cause further spread of this introduced tree.

***Removal of dead wood and dead trees***

Dead wood and trees were recorded along small sections of the REZ route. Dead wood and trees such be avoided wherever possible.

***Predation, habitat degradation, competition and disease transmission by *Sus scrofa* (Feral Pigs)***

Signs of Feral Pigs were noted along parts of the REZ route. The proposal is not likely to result in an increase in feral numbers of this introduced species.

**Conclusion**

The proposal will result in a small incremental modification of suitable habitat for *E. glaucina* however, it is considered unlikely to have a significant impact on this threatened eucalypt species.



***Eucalyptus glaucina* (Slaty Red Gum)**
**Bibliography:**

Harden, G.J. (ed.) (2002) Flora of New South Wales. Volume 2, Revised Edition. UNSW, Sydney.

**c2.4 EUCALYPTUS PARRAMATTENSIS SUBSP. DECADENS (DROOPING RED GUM)**
***Eucalyptus parramattensis* subsp. *decadens* (Drooping Red Gum)**

|                                         |                                                                                                                                                                                                                                                                                               |
|-----------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Description</b>                      | A woodland tree, up to 15 m, but usually smaller. height. Buds are ovoid 4 – 10mm long, 4 – 6 mm in diameter with a scar present. Fruit is hemispherical or globose 4 – 9 mm long, 5 – 9 mm in diameter, with the disc flat or slightly raised, usually with four exerted valves.             |
| <b>Habitat Requirements and Ecology</b> | Generally, occupies deep, low-nutrient sands, often those subject to periodic inundation or where water tables are relatively high.                                                                                                                                                           |
| <b>Distribution</b>                     | There are two separate meta-populations; the Kurri Kurri meta-population is bordered by Cessnock—Kurri Kurri in the north and Mulbring—Abedare in the south. The Tomago Sandbeds meta-population is bounded by Salt Ash and Tanilba Bay in the north and Williamtown and Tomago in the south. |
| <b>BC Act 2016</b>                      | Vulnerable                                                                                                                                                                                                                                                                                    |
| <b>BC Act 2016</b>                      | Vulnerable                                                                                                                                                                                                                                                                                    |
| <b>EPBC Act 1999</b>                    | Not Listed                                                                                                                                                                                                                                                                                    |

**Five-Part Test of Significance**

*The objective of section 7.3 of the Biodiversity Conservation Act 2016 (BC Act), the test of significance, is to provide standardised and transparent consideration of threatened species and ecological communities, and their habitats, through the development assessment process. The following is to be taken into account for the purposes of determining whether a proposed development or activity is likely to significantly affect threatened species or ecological communities, or their habitats:*

- a. *have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction,*

A number of specimens of *Eucalyptus parramattensis* subsp. *decadens* were present within two sections of proposed easement composed of Kurri Sand Woodland west of the Kurri STS to the Hunter Expressway and west over the expressway from the Kurri STS. The proposal will result in the removal of up to 32 specimens of *E. parramattensis* subsp. *decadens*. It is recommended that specimens of *E. parramattensis* subsp. *decadens* be avoided wherever possible. Given the presence of large numbers of *E. parramattensis* subsp. *decadens* within the local area, and given the recommendations, the proposal is unlikely to result in the extinction of any local population of this species.

- b. *in the case of an endangered ecological community or critically endangered ecological community, whether the proposed development or activity:*
  - i. *is likely to have an adverse effect on the extent of the ecological community such that its local occurrence is likely to be placed at risk of extinction, or*
  - ii. *is likely to substantially and adversely modify the composition of the ecological community such that its local occurrence is likely to be placed at risk of extinction,*

NA

- c. *in relation to the habitat of a threatened species or ecological community:*
  - i. *the extent to which habitat is likely to be removed or modified as a result of the proposed development or activity, and*

***Eucalyptus parramattensis* subsp. *decadens* (Drooping Red Gum)**

- ii. *whether an area of habitat is likely to become fragmented or isolated from other areas of habitat as a result of the proposed development or activity, and*
- iii. *the importance of the habitat to be removed, modified, fragmented or isolated to the long-term survival of the species or ecological community in the locality,*

The proposal will result in the removal of up to 32 specimens of *E. parramattensis* subsp. *decadens* within two sections of proposed REZ easement composed of Kurri Sand Woodland, west of the Kurri STS to the Hunter Expressway and west over the expressway from the Kurri STS. No areas are likely to become further fragmented or isolated from other areas that are important to the long-term survival of this threatened eucalypt species in the local area.

- d. *whether the proposed development or activity is likely to have an adverse effect on any declared area of outstanding biodiversity value (either directly or indirectly),*

No areas declared areas of outstanding biodiversity value were present within proximity to the REZ proposal.

- e. *whether the proposed development or activity is or is part of a key threatening process or is likely to increase the impact of a key threatening process.*

The 'Key Threatening Processes' currently listed under Schedule 4 of the BC Act 2016 that are relevant to the proposal have been addressed below:

***Clearing of Native Vegetation.***

The proposal will result in the removal of native vegetation and may be viewed as being part of this Key Threatening Process. The action is unlikely to be responsible for the significant loss of this EEC provided that recommendations for impact minimisation as listed within Section 6.4 are undertaken.

***Invasion of native plant communities by exotic perennial grasses.***

Specimens of exotic perennial grasses such as *Chloris gayana* (Rhodes Grass), *Eragrostis curvula* (African Love Grass), *Andropogon virginicus* (Whisky Grass) and *Hyparrhenia hirta* (Coolatai Grass) were present. These exotic grasses compete with species within this CEEC for space, light, and other resources and may threaten its persistence at several sites. The action is unlikely to be responsible for the significant loss of this EEC provided recommendations for impact minimisation as listed within Section 6.4 are undertaken.

***High frequency fire resulting in the disruption of life cycle processes in plants and animals and loss of vegetation structure and composition***

It is unknown what impact fire has had within areas of this EEC. The proposal is unlikely to increase

***Competition and grazing by the feral European rabbit, *Oryctolagus cuniculus****

Observations and secondary indications of the European Rabbit such as scats and diggings were noted along parts of the REZ. The proposal is not likely to result in an increase in feral numbers of this introduced species.

***Herbivory and environmental degradation caused by feral deer***

Tracks of Feral Deer were recorded within several sections of the REZ. Feral Deer would be considered to have an impact on native flora in the local area. The proposal is not likely to result in an increase in feral numbers of this introduced species.

***Infection of native plants by *Phytophthora cinnamomi****

No evidence of *Phytophthora cinnamomi* was observed during site visits. The proposal is unlikely to cause the spread of this fungus.

***Invasion, establishment and spread of Lantana (*Lantana camara*)***

Lantana was recorded within parts of the REZ proposal. The proposal is unlikely to cause the spread of this weed species.

***Introduction and establishment of Exotic Rust Fungi of the order Pucciniales pathogenic on plants of the family Myrtaceae.***



***Eucalyptus parramattensis* subsp. *decadens* (Drooping Red Gum)**

No evidence of the fungi was observed during site visits. The proposal is unlikely to cause the spread of this fungus.

**Invasion of native plant communities by African Olive *Olea europaea* subsp. *cuspidata***

The African Olive was recorded along parts of the REZ route. The proposal is unlikely to cause further spread of this introduced tree.

**Removal of dead wood and dead trees**

Dead wood and trees were recorded along small sections of the REZ route. Dead wood and trees such be avoided wherever possible.

**Predation, habitat degradation, competition and disease transmission by *Sus scrofa* (Feral Pigs)**

Signs of Feral Pigs were noted along parts of the REZ route. The proposal is not likely to result in an increase in feral numbers of this introduced species.

**Conclusion**

The proposal will result in a small incremental modification of suitable habitat for *E. parramattensis* subsp. *decadens* however, it is considered unlikely to have a significant impact on this threatened eucalypt species.

**Bibliography:**

Harden, G.J. (ed.) (2002) Flora of New South Wales. Volume 2, Revised Edition. UNSW, Sydney.

c2.5 DELMA VESCOLINEATA HUNTER VALLEY DELMA

| <b>Delma vescolineata Hunter Valley Delma</b>                                           |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|-----------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Description</b>                                                                      | <i>Delma vescolineata</i> (Hunter Valley Delma) is a legless or flap-footed lizard that grows to approximately 25 cm in total length, with a maximum snout-vent length (SVL) of 10 cm (Mahony et al. 2022). Lacks forelimbs and has very reduced hind limbs which are apparent as small flaps on either side of the vent. The top of the head is dark grey to light brown and uniform or with darker streaks or spots. There are scattered dark bars, spots or smudges on the scales bordering the mouth, the ear and sides toward the front of the body.                              |
| <b>Habitat Requirements and Ecology</b>                                                 | Has been recorded in secondary native grassland remaining after the removal or dieback of previous woody canopy vegetation (Benson 1996) in association with sparse box-gum or ironbark woodland (Mahony et al. 2022). The primary canopy species consists of <i>Eucalyptus melliodora</i> and <i>E. crebra</i> , with a diverse ground cover layer containing multiple grasses including <i>Austrostipa</i> spp., <i>Bothriochloa</i> spp., and <i>Chloris</i> spp. (Mahony et al. 2022).                                                                                             |
| <b>Distribution</b>                                                                     | Known almost entirely from a 25 km wide corridor in the Hunter Valley, between Maitland and Muswellbrook, New South Wales (Mahony et al. 2022). A further specimen has been identified from just north of Parraweena on the Liverpool Plains, approximately 80 km north-west of Muswellbrook (Mahony et al. 2022). The primary canopy species consists of <i>Eucalyptus melliodora</i> and <i>E. crebra</i> , with a diverse ground cover layer containing multiple grasses including <i>Austrostipa</i> spp., <i>Bothriochloa</i> spp., and <i>Chloris</i> spp. (Mahony et al. 2022). |
| <b>Photo</b><br>Muswellbrook<br>25 March 2024<br>(just east of the<br>Muswellbrook STS) |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| <b>BC Act 2016</b>                                                                      | Endangered                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| <b>EPBC Act 1999</b>                                                                    | Endangered                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |

**Five-Part Test of Significance**

*The objective of section 7.3 of the Biodiversity Conservation Act 2016 (BC Act), the test of significance, is to provide standardised and transparent consideration of threatened species and ecological communities, and their habitats, through the development assessment process. The following is to be taken into account for the purposes of determining whether a proposed development or activity is likely to significantly affect threatened species or ecological communities, or their habitats:*

- a. *have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction,*

*Delma vescolineata* (Hunter Valley Delma) has been recorded in 2024 within the neighbouring property to the west of the Muswellbrook proposed STS within similar habitat. According to the BioNet Atlas records (DPE, 2024) *D. vescolineata* has also been found within close proximity to the Muswellbrook STS. Taking this into consideration it is highly likely that *D. vescolineata* would utilise habitat within the impact area of the Muswellbrook STS. The proposal will likely result in a small incremental reduction of suitable habitat for *D. vescolineata*. Given the presence of large areas of similar adjoining woodland and derived grassland habitat the proposal is unlikely to result in the extinction of any local population of this reptile species.



**Delma vescolineata Hunter Valley Delma**

- b. *in the case of an endangered ecological community or critically endangered ecological community, whether the proposed development or activity:*
- is likely to have an adverse effect on the extent of the ecological community such that its local occurrence is likely to be placed at risk of extinction, or*
  - is likely to substantially and adversely modify the composition of the ecological community such that its local occurrence is likely to be placed at risk of extinction,*

NA

- c. *in relation to the habitat of a threatened species or ecological community:*
- the extent to which habitat is likely to be removed or modified as a result of the proposed development or activity, and*
  - whether an area of habitat is likely to become fragmented or isolated from other areas of habitat as a result of the proposed development or activity, and*
  - the importance of the habitat to be removed, modified, fragmented or isolated to the long-term survival of the species or ecological community in the locality,*

Given the presence of large areas of similar adjoining woodland and derived grassland habitat no areas are likely to become further fragmented or isolated from other areas that are important to the long-term survival of this threatened eucalypt species in the local area.

- d. *whether the proposed development or activity is likely to have an adverse effect on any declared area of outstanding biodiversity value (either directly or indirectly),*

No areas declared areas of outstanding biodiversity value were present within proximity to the REZ proposal.

- e. *whether the proposed development or activity is or is part of a key threatening process or is likely to increase the impact of a key threatening process.*

The 'Key Threatening Processes' currently listed under Schedule 4 of the Biodiversity Conservation Act 2016 Act that are relevant to the site have been listed in bold below followed by an assessment of the applicability of the threatening process in regards to the proposal and the species considered.

The 'Key Threatening Processes' currently listed under Schedule 4 of the BC Act 2016 that are relevant to the proposal have been addressed below:

**Clearing of Native Vegetation.**

The proposal will result in the removal of native vegetation and may be viewed as being part of this Key Threatening Process. The action is unlikely to be responsible for the significant loss of this EEC provided that recommendations for impact minimisation as listed within Section 6.4 are undertaken.

**Invasion of native plant communities by exotic perennial grasses.**

Specimens of exotic perennial grasses such as *Chloris gayana* (Rhodes Grass), *Eragrostis curvula* (African Love Grass), *Andropogon virginicus* (Whisky Grass) and *Hyparrhenia hirta* (Coolatai Grass) were present. These exotic grasses compete with species within this CEEC for space, light, and other resources and may threaten its persistence at several sites. The action is unlikely to be responsible for the significant loss of this EEC provided recommendations for impact minimisation as listed within Section 6.4 are undertaken.

**High frequency fire resulting in the disruption of life cycle processes in plants and animals and loss of vegetation structure and composition**

It is unknown what impact fire has had within areas of this EEC. The proposal is unlikely to increase

**Competition and grazing by the feral European rabbit, *Oryctolagus cuniculus***

Observations and secondary indications of the European Rabbit such as scats and diggings were noted along parts of the REZ. The proposal is not likely to result in an increase in feral numbers of this introduced species.

***Delma vescolineata* Hunter Valley Delma**
***Herbivory and environmental degradation caused by feral deer***

Tracks of Feral Deer were recorded within several sections of the REZ. Feral Deer would be considered to have an impact on native flora in the local area. The proposal is not likely to result in an increase in feral numbers of this introduced species.

***Invasion, establishment and spread of Lantana (*Lantana camara*)***

Lantana was recorded within parts of the REZ proposal. The proposal is unlikely to cause the spread of this weed species.

***Invasion of native plant communities by African Olive *Olea europaea* subsp. *cuspidata****

The African Olive was recorded along parts of the REZ route. The proposal is unlikely to cause further spread of this introduced tree.

***Removal of dead wood and dead trees***

Dead wood and trees were recorded along small sections of the REZ route. Dead wood and trees such be avoided wherever possible.

***Predation, habitat degradation, competition and disease transmission by *Sus scrofa* (Feral Pigs)***

Signs of Feral Pigs were noted along parts of the REZ route. The proposal is not likely to result in an increase in feral numbers of this introduced species.

***Predation by the *Vulpes vulpes* (Red Fox)***

The Red Fox was recorded along the REZ route a number of times and would be considered to have an impact on native fauna in the local area. The proposal is not likely to result in an increase in numbers of this introduced species.

***Predation by the *Felis catus* (Feral Cat)***

The Feral Cat was not recorded along the REZ route during fieldwork; however, this species would be considered to have an impact on native fauna in the local area. The proposal is not likely to result in an increase in feral numbers of this introduced species.

**Conclusion**

The proposal will result in a small incremental reduction of suitable habitat for *D. vescolineata* however, it is considered unlikely to have a significant impact on this threatened reptile species.

**Bibliography:**

Cogger, H.G. (2014). Reptiles and Amphibians of Australia (7th edn.). CSIRO Publishing.

Hunter Eco (2019). Biodiversity Assessment Report to develop an underground coal mining operation, (Maxwell Project) Muswellbrook NSW. Malabar Coal Limited.

Mahony SV, Cutajar T, Rowley JJJ (2022) A new species of *Delma* Gray 1831 (Squamata: Pygopodidae) from the Hunter Valley and Liverpool Plains of New South Wales. Zootaxa 5162: 541-556.



## c2.6 CLIMACTERIS PICUMNUS VICTORIAE

## BROWN TREECREEPER

| <i>Climacteris picumnus victoriae</i> Brown Treecreeper                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Description</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       | The Brown Treecreeper, Australia's largest treecreeper, is a grey-brown bird with black streaking on the lower breast and belly and black bars on the undertail.                                                                                                                                                                                                                                                                                                                              |
| <b>Habitat Requirements and Ecology</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  | This species is a medium sized insectivorous bird that occupies Eucalypt woodlands, particularly open woodlands lacking a dense understorey, River Red Gums on watercourses and around lake shores. It is sedentary and nests in tree hollows within permanent territories. They forage on tree trunks and on the ground amongst leaf litter and on fallen logs for ants, beetles and larvae. Breeding occurs from May to December.                                                           |
| <b>Distribution</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      | Distributed along the east coast of Australia from Spencer Gulf in South Australia, north to Townsville in Queensland and west to Channel Country. The eastern subspecies, <i>Climacteris picumnus victoriae</i> , is distributed through central NSW on the western side of the Great Dividing range and sparsely scattered to the east of the divide in drier areas such as the Cumberland Plain of Western Sydney, and in parts of the Hunter, Clarence, Richmond and Snowy River Valleys. |
| <b>photo</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| <b>BC Act 2016</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       | Vulnerable                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <b>EPBC Act 1999</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     | Vulnerable                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <p><b>Five-Part Test of Significance</b><br/> <i>The objective of section 7.3 of the Biodiversity Conservation Act 2016 (BC Act), the test of significance, is to provide standardised and transparent consideration of threatened species and ecological communities, and their habitats, through the development assessment process. The following is to be taken into account for the purposes of determining whether a proposed development or activity is likely to significantly affect threatened species or ecological communities, or their habitats:</i></p> <p>a. <i>have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction,</i></p> <p><i>Climacteris picumnus victoriae</i> (Brown Treecreeper) was observed just outside the existing easement within the REZ route NW of the Muswellbrook STS. As areas of suitable habitat for the Brown Treecreeper occur primarily outside the existing easement areas along the REZ route and only smaller areas of suitable habitat will be impacted the proposal is unlikely to result in the extinction of any local population of this species.</p> <p>b. <i>in the case of an endangered ecological community or critically endangered ecological community, whether the proposed development or activity:</i></p> <ol style="list-style-type: none"> <li><i>is likely to have an adverse effect on the extent of the ecological community such that its local occurrence is likely to be placed at risk of extinction, or</i></li> <li><i>is likely to substantially and adversely modify the composition of the ecological community such that its local occurrence is likely to be placed at risk of extinction,</i></li> </ol> <p>NA</p> <p>c. <i>in relation to the habitat of a threatened species or ecological community:</i></p> <ol style="list-style-type: none"> <li><i>the extent to which habitat is likely to be removed or modified as a result of the proposed development or activity, and</i></li> <li><i>whether an area of habitat is likely to become fragmented or isolated from other areas of habitat as a result of the proposed development or activity, and</i></li> <li><i>the importance of the habitat to be removed, modified, fragmented or isolated to the long-term survival of the species or ecological community in the locality,</i></li> </ol> <p>Given the presence of large areas of similar adjoining woodland and open forest habitat, no areas are likely to become further fragmented or isolated from other areas for the mobile Brown Treecreeper that are important to the long-term survival of this threatened bird species in the local area.</p> <p>d. <i>whether the proposed development or activity is likely to have an adverse effect on any declared area of outstanding biodiversity value (either directly or indirectly),</i></p> |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |

***Climacteris picumnus victoriae* Brown Treecreeper**

No areas declared areas of outstanding biodiversity value were present within proximity to the REZ proposal.

- e. *whether the proposed development or activity is or is part of a key threatening process or is likely to increase the impact of a key threatening process.*

*The 'Key Threatening Processes' currently listed under Schedule 4 of the Biodiversity Conservation Act 2016 Act that are relevant to the site have been listed in bold below followed by an assessment of the applicability of the threatening process in regards to the proposal and the species considered.*

The 'Key Threatening Processes' currently listed under Schedule 4 of the BC Act 2016 that are relevant to the proposal have been addressed below:

***Clearing of Native Vegetation.***

The proposal will result in the removal of native vegetation and may be viewed as being part of this Key Threatening Process. The action is unlikely to be responsible for the significant loss of this EEC provided that recommendations for impact minimisation as listed within Section 6.4 are undertaken.

***Invasion of native plant communities by exotic perennial grasses.***

Specimens of exotic perennial grasses such as *Chloris gayana* (Rhodes Grass), *Eragrostis curvula* (African Love Grass), *Andropogon virginicus* (Whisky Grass) and *Hyparrhenia hirta* (Coolatai Grass) were present. These exotic grasses compete with species within this CEEC for space, light, and other resources and may threaten its persistence at several sites. The action is unlikely to be responsible for the significant loss of this EEC provided recommendations for impact minimisation as listed within Section 6.4 are undertaken.

***High frequency fire resulting in the disruption of life cycle processes in plants and animals and loss of vegetation structure and composition***

It is unknown what impact fire has had within areas of this EEC. The proposal is unlikely to increase

***Herbivory and environmental degradation caused by feral deer***

Tracks of Feral Deer were recorded within several sections of the REZ. Feral Deer would be considered to have an impact on native flora in the local area. The proposal is not likely to result in an increase in feral numbers of this introduced species.

***Invasion, establishment and spread of Lantana (*Lantana camara*)***

Lantana was recorded within parts of the REZ proposal. The proposal is unlikely to cause the spread of this weed species.

***Invasion of native plant communities by African Olive *Olea europaea* subsp. *cuspidata****

The African Olive was recorded along parts of the REZ route. The proposal is unlikely to cause further spread of this introduced tree.

***Removal of dead wood and dead trees***

Dead wood and trees were recorded along small sections of the REZ route. Dead wood and trees such be avoided wherever possible.

***Predation by the *Vulpes vulpes* (Red Fox)***

The Red Fox was recorded along the REZ route a number of times and would be considered to have an impact on native fauna in the local area. The proposal is not likely to result in an increase in numbers of this introduced species.

***Predation by the *Felis catus* (Feral Cat)***

The Feral Cat was not recorded along the REZ route during fieldwork; however, this species would be considered to have an impact on native fauna in the local area. The proposal is not likely to result in an increase in feral numbers of this introduced species.



***Climacteris picumnus victoriae* Brown Treecreeper**

**Conclusion**

The proposal will result in a small incremental reduction of suitable habitat for the Brown Treecreeper however, it is considered unlikely to have a significant impact on this threatened bird species.

**Bibliography:**

Gardner J., L. (2002). Breeding Biology of the Speckled Warbler *Chthonicola sagittata*. Australian Journal of Biology, 2002, 50, 169-181.

Gardner J., L., Marsack, P. R & Blackmore, C. J (2008). Timing and sequence of moult, and the trade-off with breeding, in the Speckled Warbler (*Chthonicola sagittata*). Emu, 2008, 108, 90–96. CSIRO Publishing.

Pizzey, G. & Knight, F. (2012). The Field Guide to the Birds of Australia. 9th edn, Harper Collins Publishers Pty Limited.

**c2.7 POMATOSTOMUS TEMPORALIS TEMPORALIS GREY-CROWNED BABBLER**

***Pomatostomus temporalis temporalis* Grey-crowned Babbler**

|                                         |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
|-----------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Description</b>                      | Reaching to 30 cm long. It has a curved beak with a broad white eyebrow and a pale grey crown-stripe. A dark band passes from the bill through the eye, separating the pale throat and brow to giving a 'masked' look. It has dark greyish-brown upperparts and is paler brown on the underparts, grading to a whitish throat. It is distinctive in flight, showing white tips to the tail feathers, and orange-buff patches in the broad, rounded wings. Young birds have dark brown eyes, with the iris becoming paler with age, reaching a yellow colour by about three years.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| <b>Habitat Requirements and Ecology</b> | <p>Ranges through open forest, woodland, scrubland, farmland, and outer suburbs. Prefers woodlands with regenerating trees, tall shrubs, and an intact ground cover of grass and forbs. Members of this species are rare in settled areas and are threatened by vegetation clearance, habitat fragmentation and degradation, and competition from introduced species.</p> <p>Grey-crowned Babblers occur in sedentary family units of 2-13 birds. Babbler groups' home ranges vary from 2-53 ha (Blakers et al. 1984) Groups normally consist of a primary breeding pair along with several non-breeding birds (sometimes groups may contain two breeding pairs or two females that both breed). Most members of the group help to build nests, with the primary female contributing the most effort.</p> <p>Two types of nests are built: roost-nests (usually larger and used by the whole group) and brood-nests (for the breeding females), and often old nest sites are renovated and re-used from year to year. The large domed nests are placed in a tree fork usually 4-7 m high and are made of thick sticks with projections that make a hood and landing platform for the entrance tunnel. The nest chamber is lined with soft grass, bark, wool, and feathers. The brooding female (sometimes more than one) is fed by the other group members and all help to feed the nestlings. Larger groups tend to raise a greater number of young and two broods are usually raised per season.</p> <p>The breeding season is between July and February, however, within NSW, almost all observations of breeding activity have been from July to December. The species produces 2-6 buff, purple-brown or dark grey eggs with dark brown scribbles.</p> |
| <b>Distribution</b>                     | The Grey-crowned Babbler occurs in NSW. on the western slopes and plains, with isolated populations in coastal woodlands on the North Coast, Hunter Valley and South Coast near Nowra.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |

***Pomatostomus temporalis temporalis* Grey-crowned Babbler**



|                      |            |
|----------------------|------------|
| <b>BC Act 2016</b>   | Vulnerable |
| <b>EPBC Act 1999</b> | Not Listed |

**Five-Part Test of Significance**

**The objective of section 7.3 of the Biodiversity Conservation Act 2016 (BC Act), the test of significance, is to provide standardised and transparent consideration of threatened species and ecological communities, and their habitats, through the development assessment process. The following is to be taken into account for the purposes of determining whether a proposed development or activity is likely to significantly affect threatened species or ecological communities, or their habitats:**

- a. have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction,

*Pomatostomus temporalis* subsp. *temporalis* (Grey-crowned Babbler) was observed along the REZ route at several locations during fieldwork in areas such as the Singleton Army Base. A number of Nest/Roosts were also noted along the edge of the existing easement. Suitable foraging nesting/roosting habitat was present along large sections of the REZ route for the Grey-crowned Babbler. Taking into consideration the of preclearance searches for nests before any tree removal, and relocation of deadwood outside impact areas the proposal is unlikely to have a significant impact on the Grey-crowned Babbler such that a local population would be placed at risk of extinction.

- b. in the case of an endangered ecological community or critically endangered ecological community, whether the proposed development or activity:
  - i. is likely to have an adverse effect on the extent of the ecological community such that its local occurrence is likely to be placed at risk of extinction, or
  - ii. is likely to substantially and adversely modify the composition of the ecological community such that its local occurrence is likely to be placed at risk of extinction,

NA

- c. in relation to the habitat of a threatened species or ecological community:
  - i. the extent to which habitat is likely to be removed or modified as a result of the proposed development or activity, and
  - ii. whether an area of habitat is likely to become fragmented or isolated from other areas of habitat as a result of the proposed development or activity, and
  - iii. the importance of the habitat to be removed, modified, fragmented or isolated to the long-term survival of the species or ecological community in the locality,



**Pomatostomus temporalis temporalis      Grey-crowned Babbler**

Given the presence of large areas of similar adjoining woodland and open forest habitat, no areas are likely to become further fragmented or isolated from other areas for the Grey-crowned Babbler that are important to the long-term survival of this threatened bird species in the local area.

- d. *whether the proposed development or activity is likely to have an adverse effect on any declared area of outstanding biodiversity value (either directly or indirectly),*

No areas declared areas of outstanding biodiversity value were present within proximity to the REZ proposal.

- e. *whether the proposed development or activity is or is part of a key threatening process or is likely to increase the impact of a key threatening process.*

*The 'Key Threatening Processes' currently listed under Schedule 4 of the Biodiversity Conservation Act 2016 Act that are relevant to the site have been listed in bold below followed by an assessment of the applicability of the threatening process in regards to the proposal and the species considered.*

The 'Key Threatening Processes' currently listed under Schedule 4 of the BC Act 2016 that are relevant to the proposal have been addressed below:

**Clearing of Native Vegetation.**

The proposal will result in the removal of native vegetation and may be viewed as being part of this Key Threatening Process. The action is unlikely to be responsible for the significant loss of this EEC provided that recommendations for impact minimisation as listed within Section 6.4 are undertaken.

**Invasion of native plant communities by exotic perennial grasses.**

Specimens of exotic perennial grasses such as *Chloris gayana* (Rhodes Grass), *Eragrostis curvula* (African Love Grass), *Andropogon virginicus* (Whisky Grass) and *Hyparrhenia hirta* (Coolatai Grass) were present. These exotic grasses compete with species within this CEEC for space, light, and other resources and may threaten its persistence at several sites. The action is unlikely to be responsible for the significant loss of this EEC provided recommendations for impact minimisation as listed within Section 6.4 are undertaken.

**High frequency fire resulting in the disruption of life cycle processes in plants and animals and loss of vegetation structure and composition**

It is unknown what impact fire has had within areas of this EEC. The proposal is unlikely to increase

**Herbivory and environmental degradation caused by feral deer**

Tracks of Feral Deer were recorded within several sections of the REZ. Feral Deer would be considered to have an impact on native flora in the local area. The proposal is not likely to result in an increase in feral numbers of this introduced species.

**Invasion, establishment and spread of Lantana (*Lantana camara*)**

Lantana was recorded within parts of the REZ proposal. The proposal is unlikely to cause the spread of this weed species.

**Invasion of native plant communities by African Olive *Olea europaea* subsp. *cuspidata***

The African Olive was recorded along parts of the REZ route. The proposal is unlikely to cause further spread of this introduced tree.

**Removal of dead wood and dead trees**

Dead wood and trees were recorded along small sections of the REZ route. Dead wood and trees such be avoided wherever possible.

**Predation by the *Vulpes vulpes* (Red Fox)**

The Red Fox was recorded along the REZ route a number of times and would be considered to have an impact on native fauna in the local area. The proposal is not likely to result in an increase in numbers of this introduced species.

***Pomatostomus temporalis temporalis* Grey-crowned Babbler**

**Predation by the *Felis catus* (Feral Cat)**

The Feral Cat was not recorded along the REZ route during fieldwork; however, this species would be considered to have an impact on native fauna in the local area. The proposal is not likely to result in an increase in feral numbers of this introduced species.

**Conclusion**

The proposal will result in a small incremental reduction of suitable habitat for the Grey-crowned Babbler however, it is considered unlikely to have a significant impact on this threatened bird species.

**Bibliography:**

Blackmore, C. J., and Heinsohn, R. (2007). *Reproductive success and helper effects in the cooperatively breeding Grey-crowned Babbler Pomatostomus temporalis*. Journal of Zoology (London, England) 273, 326–332.

Blakers, M., Davies, S.J.J.F., & Reilly, P.N. (1984) *The Atlas of Australian Birds*. Melbourne University Press, Melbourne.

Davidson, I. and Robinson, D. (1992). 'Grey-crowned Babbler, *Pomatostomus temporalis*: Action Statement No. 34.' Department of Conservation and environment: Victoria.

Dow, D. D., and King, B. R. (1984). *Communal building of brood and roost nests by the Grey-crowned Babbler Pomatostomus temporalis*. Emu 84, 193–199.

Fischer, S. E. (2011). The effect of fire on the breeding ecology of the Grey-crowned Babbler (*Pomatostomus temporalis*) in the Australian monsoon tropics. Student thesis: Masters by Research - Charles Darwin University.

King, B. R. (1980). *Social organisation and behaviour of the Grey-crowned Babbler Pomatostomus temporalis* Emu 80: 59-76.

Murray, M., Bell, S., Hoyer, G. (2002). *Local Biodiversity Assessment Guidelines: Lower Hunter Central Coast Region 2002*. Lower Hunter & Central Coast Regional Environmental Management Strategy, Callaghan.

NSW National Parks and Wildlife Service. (2001). *Threatened Species Conservation Act 1995 NSW Scientific Committee Final Determination, Grey-crowned Babbler (eastern subspecies) Pomatostomus temporalis*.

Pizzey, G. & Knight, F. (2012). *The Field Guide to the Birds of Australia*. 9<sup>th</sup> edn, Harper Collins Publishers Pty Limited.

Slater, P., Slater, P., & Slater, R., (1997). *The Slater Field Guide to Australian Birds*, Lansdowne Publishing, Sydney.

Kate P. Stevens, K. P., Holland, G. J., Clarke, R. H., Cooke, R., Bennett, A F (2015). *What Determines Habitat Quality for a Declining Woodland Bird in a Fragmented Environment: The Grey-Crowned Babbler Pomatostomus temporalis in South-Eastern Australia?*. PLOS ONE DOI:10.1371/journal.pone.0130738 June 22, 2015.



## c2.8 HALIAEETUS LEUCOGASTER WHITE-BELLIED SEA-EAGLE

| <i>Haliaeetus leucogaster</i> (White-bellied Sea-Eagle)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Description</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      | The White-bellied Sea-Eagle has long broad wings and a short, wedge-shaped tail. It measures 75–85 cm in length, and has a wingspan of 180–220 cm. Adults are predominantly white and grey. The head, breast and belly, and the feathering on the legs, are white. The tail is grey at the base and has a white tip. The large, hooked bill is grey with a darker tip, and the eye is dark brown. The legs and feet are cream-white, with long black talons. Both sexes are similar in appearance, but females are larger than the males. Juveniles are brown with lighter markings.                               |
| <b>Habitat Requirements and Ecology</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 | Habitats are characterised by the presence of large areas of open water including larger rivers, swamps, lakes, and the sea. Breeding habitat consists of mature tall open forest, open forest, tall woodland, and swamp sclerophyll forest close to foraging habitat. Nest trees are typically large emergent eucalypts and often have emergent dead branches or large dead trees nearby which are used as 'guard roosts'. Nests are large structures built from sticks and lined with leaves or grass. Typically lays two eggs between June and September with young birds remaining in the nest for 65-70 days. |
| <b>Distribution</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     | Found around the Australian coastline, including Tasmania, and well inland along rivers and wetlands of the Murray Darling Basin. In New South Wales it is widespread along the east coast, and along all major inland rivers and waterways.                                                                                                                                                                                                                                                                                                                                                                       |
| <b>Nest in Aberglasslyn 2010.</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| <b>BC Act 2016</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      | Vulnerable                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| <b>EPBC Act 1999</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | Listed marine migratory species.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| <b>Five-Part Test of Significance</b><br><b>The objective of section 7.3 of the Biodiversity Conservation Act 2016 (BC Act), the test of significance, is to provide standardised and transparent consideration of threatened species and ecological communities, and their habitats, through the development assessment process. The following is to be taken into account for the purposes of determining whether a proposed development or activity is likely to significantly affect threatened species or ecological communities, or their habitats:</b>                                                                                                           |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <p>a. have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction,</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <p>One specimen of <i>Haliaeetus leucogaster</i> (White-bellied Sea-Eagle) was observed flying over the proposed site for the Eastern Hub near the northern shore of Lake Liddell during fieldwork. Suitable hunting habitat was present within the waters of Lake Liddell; however, no hunting habitat was present within the impact area for the Eastern Hub. No likely nesting habitat would also be impacted is part of the proposal. Taking into consideration the lack of habitat within the impact area the proposal is not likely to have a significant impact on <i>H. leucogaster</i> such that a local population would be placed at risk of extinction.</p> |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |

***Haliaeetus leucogaster (White-bellied Sea-Eagle)***

- b. *in the case of an endangered ecological community or critically endangered ecological community, whether the proposed development or activity:*
  - i. *is likely to have an adverse effect on the extent of the ecological community such that its local occurrence is likely to be placed at risk of extinction, or*
  - ii. *is likely to substantially and adversely modify the composition of the ecological community such that its local occurrence is likely to be placed at risk of extinction,*
- c. *in relation to the habitat of a threatened species or ecological community:*
  - i. *the extent to which habitat is likely to be removed or modified as a result of the proposed development or activity, and*
  - ii. *whether an area of habitat is likely to become fragmented or isolated from other areas of habitat as a result of the proposed development or activity, and*
  - iii. *the importance of the habitat to be removed, modified, fragmented or isolated to the long-term survival of the species or ecological community in the locality,*
- d. *whether the proposed development or activity is likely to have an adverse effect on any declared area of outstanding biodiversity value (either directly or indirectly),*
- e. *whether the proposed development or activity is or is part of a key threatening process or is likely to increase the impact of a key threatening process.*

*The 'Key Threatening Processes' currently listed under Schedule 4 of the Biodiversity Conservation Act 2016 Act that are relevant to the site have been listed in bold below followed by an assessment of the applicability of the threatening process in regards to the proposal and the species considered.*

The 'Key Threatening Processes' currently listed under Schedule 4 of the BC Act 2016 that are relevant to the proposal have been addressed below:

***Clearing of Native Vegetation.***

The proposal will result in the removal of native vegetation and may be viewed as being part of this Key Threatening Process. The action is unlikely to be responsible for the significant loss of habitat for this fauna species provided that recommendations for impact minimisation as listed within Section 6.4 are undertaken.

***High frequency fire resulting in the disruption of life cycle processes in plants and animals and loss of vegetation structure and composition***

It is unknown what impact fire has had within areas of habitat.

***Invasion, establishment and spread of Lantana (*Lantana camara*)***

Lantana was recorded within parts of the REZ proposal. The proposal is unlikely to cause the spread of this weed species.

***Removal of dead wood and dead trees***

Dead wood and trees were recorded along small sections of the REZ route. Dead wood and trees such be avoided wherever possible.

***Predation by the *Vulpes vulpes* (Red Fox)***

The Red Fox was recorded along the REZ route a number of times and would be considered to have an impact on native fauna in the local area. The proposal is not likely to result in an increase in numbers of this introduced species.

***Predation by the *Felis catus* (Feral Cat)***

The Feral Cat was not recorded along the REZ route during fieldwork; however, this species would be considered to have an impact on native fauna in the local area. The proposal is not likely to result in an increase in feral numbers of this introduced species.



***Haliaeetus leucogaster* (White-bellied Sea-Eagle)**

**Conclusion**

The proposal will not likely impact any areas of hunting or potential nesting habitat for *H. leucogaster*, therefore the proposal it is considered not likely to have a significant impact on this threatened bird species.

**Bibliography:**

Pizzey, G. & Knight, F. (2012). The Field Guide to the Birds of Australia. 9th edn, Harper Collins Publishers Pty Limited.

Pryor, K.J. (2022) Effects of human disturbance during residential developments on the productivity of White-bellied Sea-Eagle in the Hunter Region, New South Wales. *The Whistler* **16 (2022)**: 66-73.

Shephard, J.M., C.P. Catterall & J.M. Hughes (2005). Long-term variation in the distribution of the White-bellied Sea-Eagle (*Haliaeetus leucogaster*) across Australia. *Austral Ecology*. 30:131-145.

## **APPENDIX D**

### **TOTAL FLORA**



Introduced species are indicated by an asterisk (“\*”).

The following standard abbreviations are used to indicate subspecific taxa:

- subsp.** subspecies
- var.-** variety
- x -** hybrid between the two indicated species

## Threatened Species - NSW Biodiversity Conservation Act 2016 (BC Act)

- V** Vulnerable
- E1** Endangered
- E2** Endangered Population
- E4A** Critically Endangered Population

## Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act)

- V** Vulnerable
- E** Endangered
- CE** Critically Endangered

## Serious and Irreversible Impact SAIL

## Regional Significance (Hunter Rare Plants Database – Version 1 2003)

- L** endemic to Hunter Region
- DA** disjunct in the Hunter Region, rare or localized (aggregated)
- DB** disjunct in the Hunter Region, widespread and uncommon (broad)
- R** rare but extends beyond the Hunter Region
- U** everywhere uncommon
- N** at northern distributional limit in the Hunter
- E** at eastern distributional limit in the Hunter
- S** at southern distributional limited in the Hunter
- W** at western distributional limited in the Hunter
- T** may be threatened in the Hunter Region
- S** Probably secure in the Hunter Region

## Weeds

### Priorities under the Biosecurity Act 2015

- G** General Biosecurity Duty - any person dealing with plant matter must take measures to prevent, minimise or eliminate the biosecurity risk (as far as is reasonably practicable).
- P** Prohibition on dealings - Must not be imported into the State or sold.
- R** Regional Recommended Measure - Land managers mitigate the risk of the plant being introduced to their land. Land managers reduce impacts from the plant on priority assets. Land managers prevent spread from their land where feasible. The plant or parts of the plant are not traded, carried, grown or released into the environment.

### NSW BC Act 2016

- T** Listed as a Threatening Process under the NSW BC Act 2016.

### National

- N** Weed of National Significance (WoNS)

Table D1 Total Flora List

| SCIENTIFIC NAME                                              | COMMON NAME            | BC ACT | EPBC ACT | SAI | REGIONALLY SIGNIFICANT | BIOSECURITY ACT 2015 | FLOWERING PERIOD |
|--------------------------------------------------------------|------------------------|--------|----------|-----|------------------------|----------------------|------------------|
| <b>CLASS FILICOPSIDA (Ferns)</b>                             |                        |        |          |     |                        |                      |                  |
| <b>Azollaceae</b>                                            |                        |        |          |     |                        |                      |                  |
| <i>Azolla pinnata</i>                                        | Ferny Azolla           |        |          |     |                        |                      |                  |
|                                                              |                        |        |          |     |                        |                      |                  |
| <b>Dennstaedtiaceae</b>                                      |                        |        |          |     |                        |                      |                  |
| <i>Hypolepis muelleri</i>                                    | Harsh Ground Fern      |        |          |     |                        |                      |                  |
| <i>Pteridium esculentum</i>                                  | Bracken                |        |          |     |                        |                      |                  |
|                                                              |                        |        |          |     |                        |                      |                  |
| <b>Pteridaceae</b>                                           |                        |        |          |     |                        |                      |                  |
| <i>Adiantum aethiopicum</i>                                  | Common Maidenhair Fern |        |          |     |                        |                      |                  |
| <i>Cheilanthes distans</i>                                   | Bristly Cloak Fern     |        |          |     |                        |                      |                  |
| <i>Cheilanthes sieberi</i> ssp. <i>sieberi</i>               | Mulga Fern             |        |          |     |                        |                      |                  |
|                                                              |                        |        |          |     |                        |                      |                  |
| <b>CYCADOPSIDA (Cycads)</b>                                  |                        |        |          |     |                        |                      |                  |
| <b>Zamiaceae</b>                                             |                        |        |          |     |                        |                      |                  |
| <i>Macrozamia flexuosa</i>                                   |                        |        |          |     | L N S W                |                      |                  |
|                                                              |                        |        |          |     |                        |                      |                  |
| <b>MAGNOLIOPSIDA: Magnoliidae</b>                            |                        |        |          |     |                        |                      |                  |
| <b>LILOPSIDA: (Monocotyledons)</b>                           |                        |        |          |     |                        |                      |                  |
|                                                              |                        |        |          |     |                        |                      |                  |
| <b>Amaryllidaceae</b>                                        |                        |        |          |     |                        |                      |                  |
| <i>*Nothoscordum gracile</i>                                 | Onion Weed             |        |          |     |                        |                      |                  |
|                                                              |                        |        |          |     |                        |                      |                  |
| <b>Asparagaceae</b>                                          |                        |        |          |     |                        |                      |                  |
| <i>*Asparagus asparagoides</i>                               | Bridal Creeper         |        |          |     |                        |                      |                  |
| <i>*Asparagus aethiopicus</i>                                | Asparagus Fern         |        |          |     |                        |                      |                  |
| <i>*Asparagus officinalis</i>                                | Asparagus              |        |          |     |                        |                      |                  |
| <i>Arthropodium minus</i>                                    | Small Vanilla Lily     |        |          |     |                        |                      |                  |
| <i>Arthropodium strictum</i> syn. <i>Dichopogon strictus</i> | Chocolate Lily         |        |          |     |                        |                      |                  |
| <i>Eustrephus latifolius</i>                                 | Wombat Berry           |        |          |     |                        |                      |                  |
| <i>Laxmannia gracilis</i>                                    | Slender Wire Lily      |        |          |     |                        |                      |                  |
| <i>Lomandra confertifolia</i>                                |                        |        |          |     |                        |                      |                  |
| <i>Lomandra filiformis</i>                                   |                        |        |          |     |                        |                      |                  |
| <i>Lomandra gracilis</i>                                     |                        |        |          |     |                        |                      |                  |
| <i>Lomandra longifolia</i>                                   | Spiny Mat Rush         |        |          |     |                        |                      | Aug, Sept        |
| <i>Lomandra multiflora</i> subsp. <i>multiflora</i>          | Many-flowered Mat-rush |        |          |     |                        |                      | Sept             |

| SCIENTIFIC NAME                                 | COMMON NAME         | BC ACT | EPBC ACT | SAII | REGIONALLY SIGNIFICANT | BIOSECURITY ACT 2015 | FLOWERING PERIOD |
|-------------------------------------------------|---------------------|--------|----------|------|------------------------|----------------------|------------------|
| <b>Asphodelaceae</b>                            |                     |        |          |      |                        |                      |                  |
| <i>Caesia parviflora</i>                        | Pale Grass-lily     |        |          |      |                        |                      |                  |
| <i>Dianella caerulea</i> var. <i>caerulea</i>   | Blue Flax-lily      |        |          |      |                        |                      |                  |
| <i>Dianella caerulea</i> var. <i>producta</i>   | Blue Flax-lily      |        |          |      |                        |                      |                  |
| <i>Dianella longifolia</i>                      | Smooth Flax Lily    |        |          |      |                        |                      |                  |
| <i>Dianella revoluta</i>                        | Blue Flax-lily      |        |          |      |                        |                      |                  |
| <i>Geitonoplesium cymosum</i>                   | Scrambling Lily     |        |          |      |                        |                      |                  |
| <i>Stypandra glauca</i>                         | Nodding Blue Lily   |        |          |      |                        |                      |                  |
| <i>Tricoryne elatior</i>                        | Yellow Rush-lily    |        |          |      |                        |                      |                  |
| <i>Xanthorrhoea glauca</i> subsp. <i>glauca</i> | Giant Grass Tree    |        |          |      |                        |                      |                  |
|                                                 |                     |        |          |      |                        |                      |                  |
| <b>Colchicaceae syn Uvulariaceae</b>            |                     |        |          |      |                        |                      |                  |
| <i>Wurmbea dioica</i>                           | Early Nancy         |        |          |      |                        |                      |                  |
|                                                 |                     |        |          |      |                        |                      |                  |
| <b>Commelinaceae</b>                            |                     |        |          |      |                        |                      |                  |
| <i>Commelina cyanea</i>                         | Scurvy Weed         |        |          |      |                        |                      |                  |
| * <i>Tradescantia fluminensis</i>               | Wandering Jew       |        |          |      |                        |                      |                  |
|                                                 |                     |        |          |      |                        |                      |                  |
| <b>Cyperaceae</b>                               |                     |        |          |      |                        |                      |                  |
| <i>Carex appressa</i>                           | Saw Sedge           |        |          |      |                        |                      |                  |
| * <i>Cyperus brevifolius</i>                    | Mullumbimby Couch   |        |          |      |                        |                      |                  |
| <i>Cyperus difformis</i>                        | Dirty Dora          |        |          |      |                        |                      |                  |
| * <i>Cyperus eragrostis</i>                     | Umbrella Sedge      |        |          |      |                        |                      |                  |
| <i>Cyperus gracilis</i>                         | Slender Flat-sedge  |        |          |      |                        |                      |                  |
| <i>Eleocharis sphacelata</i>                    | Tall Spike-rush     |        |          |      |                        |                      |                  |
| <i>Fimbristylis dichotoma</i>                   | Common Fringe Sedge |        |          |      |                        |                      |                  |
| <i>Gahnia aspera</i>                            | Rough Saw Sedge     |        |          |      |                        |                      |                  |
| <i>Lepidosperma laterale</i>                    | Sword Sedge         |        |          |      |                        |                      |                  |
| <i>Ptilothrix deusta</i>                        |                     |        |          |      |                        |                      |                  |
| <i>Schoenoplectus mucronatus</i>                |                     |        |          |      |                        |                      |                  |
| <i>Schoenus apogon</i>                          | Common Bog-rush     |        |          |      |                        |                      |                  |
|                                                 |                     |        |          |      |                        |                      |                  |
| <b>Iridaceae</b>                                |                     |        |          |      |                        |                      |                  |
| * <i>Romulea rosea</i> var. <i>australis</i>    | Onion Grass         |        |          |      |                        |                      |                  |
|                                                 |                     |        |          |      |                        |                      |                  |
| <b>Juncaceae</b>                                |                     |        |          |      |                        |                      |                  |
| * <i>Juncus acutus</i>                          | Spiny Rush          |        |          |      |                        |                      |                  |
| * <i>Juncus cognatus</i>                        |                     |        |          |      |                        |                      |                  |



| SCIENTIFIC NAME                                                   | COMMON NAME                | BC ACT | EPBC ACT | SAII | REGIONALLY SIGNIFICANT | BIOSECURITY ACT 2015 | FLOWERING PERIOD |
|-------------------------------------------------------------------|----------------------------|--------|----------|------|------------------------|----------------------|------------------|
| <i>Juncus planifolius</i>                                         |                            |        |          |      |                        |                      |                  |
| <i>Juncus prismatocarpus</i>                                      |                            |        |          |      |                        |                      |                  |
| <i>Juncus subsecundus</i>                                         | Finger Rush                |        |          |      |                        |                      |                  |
| <i>Juncus usitatus</i>                                            | Common Rush                |        |          |      |                        |                      |                  |
|                                                                   |                            |        |          |      |                        |                      |                  |
| <b>Orchidaceae</b>                                                |                            |        |          |      |                        |                      |                  |
| <i>Diuris tricolor</i>                                            | Pine Donkey Orchid         | V      | V        |      | E U R*                 |                      | Sept. Oct, Nov   |
|                                                                   |                            |        |          |      |                        |                      |                  |
| <b>Philydraceae</b>                                               |                            |        |          |      |                        |                      |                  |
| <i>Philydrum lanuginosum</i>                                      | Woolly Frogsmouth          |        |          |      |                        |                      |                  |
|                                                                   |                            |        |          |      |                        |                      |                  |
| <b>Poaceae</b>                                                    |                            |        |          |      |                        |                      |                  |
| * <i>Andropogon virginicus</i>                                    | Whisky Grass               |        |          |      |                        |                      |                  |
| * <i>Anthoxanthum odoratum</i>                                    | Sweet Vernal Grass         |        |          |      |                        |                      |                  |
| <i>Aristida ramosa</i> var. <i>ramosa</i>                         | Three-awn Speargrass       |        |          |      | W?                     |                      |                  |
| <i>Aristida vagans</i>                                            | Three-awn Speargrass       |        |          |      |                        |                      |                  |
| * <i>Arundo donax</i>                                             | Giant Reed                 |        |          |      |                        |                      |                  |
| <i>Austrostipa verticillata</i>                                   | Slender Bamboo Grass       |        |          |      |                        |                      |                  |
| * <i>Avena fatua</i>                                              | Wild Oats                  |        |          |      |                        |                      |                  |
| * <i>Axonopus fissifolius</i>                                     | Narrow-leaved Carpet Grass |        |          |      |                        |                      |                  |
| <i>Bothriochloa decipens</i>                                      | Red grass                  |        |          |      |                        |                      |                  |
| * <i>Briza maxima</i>                                             | Quaking Grass              |        |          |      |                        |                      |                  |
| * <i>Briza minor</i>                                              | Shivery Grass              |        |          |      |                        |                      |                  |
| * <i>Bromus catharticus</i>                                       | Prairie Grass              |        |          |      |                        |                      |                  |
| <i>Capillipedium spicigerum</i>                                   | Scented Top                |        |          |      |                        |                      |                  |
| * <i>Cenchrus clandestinus</i> syn <i>Pennisetum clandestinum</i> | Kikuyu                     |        |          |      |                        |                      |                  |
| * <i>Chloris gayana</i>                                           | Rhodes Grass               |        |          |      |                        |                      |                  |
| <i>Chloris ventricosa</i>                                         | Tall Windmill Grass        |        |          |      |                        |                      |                  |
| <i>Cynodon dactylon</i>                                           | Common Couch               |        |          |      |                        |                      |                  |
| <i>Cymbopogon refractus</i>                                       | Barbed Wire Grass          |        |          |      |                        |                      |                  |
| <i>Dichelachne micrantha</i>                                      | Plume Grass                |        |          |      |                        |                      |                  |
| <i>Digitaria parviflora</i>                                       | Smallflower Fingergrass    |        |          |      |                        |                      |                  |
| * <i>Ehrharta erecta</i>                                          | Panic Veldt Grass          |        |          |      |                        |                      |                  |
| * <i>Elusine indica</i>                                           | Crowsfoot Grass            |        |          |      |                        |                      |                  |
| <i>Enteropogon acicularis</i>                                     |                            |        |          |      |                        |                      |                  |
| <i>Entolasia marginata</i>                                        | Bordered Panic             |        |          |      |                        |                      |                  |
| <i>Entolasia stricta</i>                                          | Wiry Panic                 |        |          |      |                        |                      |                  |

| SCIENTIFIC NAME                                           | COMMON NAME                | BC ACT | EPBC ACT | SAII | REGIONALLY SIGNIFICANT | BIOSECURITY ACT 2015 | FLOWERING PERIOD |
|-----------------------------------------------------------|----------------------------|--------|----------|------|------------------------|----------------------|------------------|
| <i>Eragrostis brownii</i>                                 | Browns Love Grass          |        |          |      |                        |                      |                  |
| * <i>Eragrostis cilianensis</i>                           | Stinkgrass                 |        |          |      |                        |                      |                  |
| * <i>Eragrostis curvula</i>                               | African Lovegrass          |        |          |      |                        |                      |                  |
| <i>Eragrostis leptostachya</i>                            | Paddock Lovegrass          |        |          |      |                        |                      |                  |
| * <i>Eragrostis tenuifolia</i>                            | Elastic Grass              |        |          |      |                        |                      |                  |
| * <i>Hyparrhenia hirta</i>                                | Coolatai Grass             |        |          |      |                        |                      | Sept             |
| <i>Imperata cylindrica</i> var. <i>major</i>              | Blady Grass                |        |          |      |                        |                      |                  |
| * <i>Lolium perenne</i>                                   | Perennial Ryegrass         |        |          |      |                        |                      |                  |
| * <i>Megathyrsus maximus</i> syn. <i>Panicum maximum</i>  | Guinea Grass               |        |          |      |                        |                      |                  |
| <i>Microlaena stipoides</i> var. <i>stipoides</i>         | Weeping Meadow Grass       |        |          |      |                        |                      |                  |
| <i>Oplismenus imbecillis</i>                              | Basket Grass               |        |          |      |                        |                      |                  |
| <i>Panicum effusum</i>                                    | Hairy Panic                |        |          |      |                        |                      |                  |
| <i>Paspalidium distans</i>                                | Spreading Panic Grass      |        |          |      |                        |                      |                  |
| * <i>Paspalum dilatatum</i>                               | Paspalum                   |        |          |      |                        |                      |                  |
| <i>Paspalum distichum</i>                                 | Water Couch                |        |          |      |                        |                      |                  |
| * <i>Paspalum urvillei</i>                                | Vasey Grass                |        |          |      |                        |                      |                  |
| <i>Phragmites australis</i>                               | Native Reed                |        |          |      |                        |                      |                  |
| * <i>Poa annua</i>                                        | Winter Grass               |        |          |      |                        |                      | Aug, Sept        |
| <i>Rytidosperma fulvum</i>                                | Wallaby Grass              |        |          |      |                        |                      |                  |
| * <i>Setaria parviflora</i> syn. <i>Setaria gracillis</i> | Slender Pigeon Grass       |        |          |      |                        |                      |                  |
| * <i>Setaria sphacelata</i>                               | South African Pigeon Grass |        |          |      |                        |                      |                  |
| * <i>Sporobolus africanus</i>                             | Parramatta Grass           |        |          |      |                        |                      |                  |
| * <i>Sorghum halepense</i>                                | Johnson Grass              |        |          |      |                        |                      |                  |
| * <i>Sporobolus africanus</i>                             | (African) Parramatta Grass |        |          |      |                        |                      |                  |
| <i>Sporobolus creber</i>                                  | Slender Rats Tail          |        |          |      |                        |                      |                  |
| <i>Themeda triandra</i> syn. <i>Themeda australis</i>     | Kangaroo Grass             |        |          |      |                        |                      | Oct, Nov         |
| <b>Typhaceae</b>                                          |                            |        |          |      |                        |                      |                  |
| <i>Typha orientalis</i>                                   | Cumbungi                   |        |          |      |                        |                      |                  |
| <b>MAGNOLIIDAE (Dicotyledons)</b>                         |                            |        |          |      |                        |                      |                  |
| <b>Acanthaceae</b>                                        |                            |        |          |      |                        |                      |                  |
| <i>Brunoniella australis</i>                              | Blue Trumpet               |        |          |      |                        |                      |                  |
| <i>Rostellularia adscendens</i>                           |                            |        |          |      |                        |                      |                  |
| <b>Aceraceae</b>                                          |                            |        |          |      |                        |                      |                  |
| * <i>Acer pseudoplatanus</i>                              | Sycamore Maple             |        |          |      |                        |                      |                  |

| SCIENTIFIC NAME                                         | COMMON NAME                | BC ACT | EPBC ACT | SAIL | REGIONALLY SIGNIFICANT | BIOSECURITY ACT 2015 | FLOWERING PERIOD |
|---------------------------------------------------------|----------------------------|--------|----------|------|------------------------|----------------------|------------------|
| <b>Aizoaceae</b>                                        |                            |        |          |      |                        |                      |                  |
| * <i>Galenia pubescens</i>                              | Galenia                    |        |          |      |                        |                      |                  |
| <b>Amaranthaceae</b>                                    |                            |        |          |      |                        |                      |                  |
| <i>Alternanthera denticulata</i>                        | Lesser Joyweed             |        |          |      |                        |                      |                  |
| * <i>Gomphrena celosioides</i>                          | Gomphrena Weed             |        |          |      |                        |                      |                  |
| <b>Anacardiaceae</b>                                    |                            |        |          |      |                        |                      |                  |
| * <i>Schinus areira</i> syn. <i>Schinus molle</i>       | Pepper Tree                |        |          |      |                        |                      |                  |
| <b>Apiaceae</b>                                         |                            |        |          |      |                        |                      |                  |
| <i>Centella asiatica</i>                                | Indian Pennywort           |        |          |      |                        |                      |                  |
| * <i>Cyclospermum leptophyllum</i>                      | Slender Celery             |        |          |      |                        |                      |                  |
| <i>Daucus glochidiatus</i>                              | Native Carrot              |        |          |      |                        |                      |                  |
| * <i>Foeniculum vulgare</i>                             | Fennel                     |        |          |      |                        |                      |                  |
| <b>Apocynaceae</b>                                      |                            |        |          |      |                        |                      |                  |
| * <i>Araujia sericifera</i> (syn. <i>A. hortorum</i> ). | Moth Plant                 |        |          |      |                        |                      |                  |
| * <i>Gomphocarpus fruticosus</i>                        | Narrow-leaved Cottonbush   |        |          |      |                        |                      |                  |
| <i>Parsonsia straminea</i> var. <i>straminea</i>        | Common Silkpod             |        |          |      | W?                     |                      |                  |
| <b>Araliaceae</b>                                       |                            |        |          |      |                        |                      |                  |
| * <i>Hedera helix</i>                                   | English Ivy                |        |          |      |                        |                      |                  |
| * <i>Hydrocotyle bonariensis</i>                        | Kurnell Curse              |        |          |      |                        |                      |                  |
| <i>Polyscias sambucifolia</i>                           | Elderberry Panax           |        |          |      |                        |                      |                  |
| <b>Asteraceae</b>                                       |                            |        |          |      |                        |                      |                  |
| * <i>Ageratina adenophora</i>                           | Crofton Weed               |        |          |      |                        |                      | Sept             |
| * <i>Aster subulatus</i> syn. <i>Aster squamatus</i>    | Bushy Starwort             |        |          |      |                        |                      |                  |
| * <i>Bidens pilosa</i>                                  | Cobblers Pegs              |        |          |      |                        |                      |                  |
| * <i>Bidens subalternans</i>                            | Greater Beggar's Ticks     |        |          |      |                        |                      |                  |
| <i>Brachyscome multifida</i>                            | Cut-leaved Daisy           |        |          |      |                        |                      | Sept             |
| <i>Calotis cuneifolia</i>                               | Blue Burr-daisy            |        |          |      |                        |                      |                  |
| <i>Calotis lappulacea</i>                               | Yellow Burr-daisy          |        |          |      |                        |                      | Sept             |
| * <i>Carthamus lanatus</i>                              | Saffron Thistle            |        |          |      |                        |                      |                  |
| <i>Cassinia sifton</i> syn <i>Cassinia arcuata</i>      | Sifton Bush, Chinese Scrub |        |          |      |                        |                      |                  |
| <i>Chrysocephalum apiculatum</i>                        | Yellow Buttons             |        |          |      |                        |                      |                  |



| SCIENTIFIC NAME                                                     | COMMON NAME           | BC ACT | EPBC ACT | SAII | REGIONALLY SIGNIFICANT | BIOSECURITY ACT 2015 | FLOWERING PERIOD |
|---------------------------------------------------------------------|-----------------------|--------|----------|------|------------------------|----------------------|------------------|
| <i>Chrysocephalum semipapposum</i>                                  | Clustered Everlasting |        |          |      |                        |                      |                  |
| * <i>Cirsium vulgare</i>                                            | Spear Thistle         |        |          |      |                        |                      | Sept             |
| * <i>Conyza bonariensis</i>                                         | Flax-leaved Fleabane  |        |          |      |                        |                      |                  |
| * <i>Coreopsis lanceolata</i>                                       | Coreopsis             |        |          |      |                        |                      |                  |
| <i>Cotula australis</i>                                             | Carrot Weed           |        |          |      |                        |                      |                  |
| <i>Cymbonotus lawsonianus</i>                                       | Bear's Ear            |        |          |      |                        |                      |                  |
| <i>Euchiton sphaericus</i>                                          | Common Cudweed        |        |          |      |                        |                      |                  |
| * <i>Facelis retusa</i>                                             | Facelis               |        |          |      |                        |                      |                  |
| * <i>Gamochaeta coarctata</i> syn. <i>Gamochaeta spicata</i>        | Spiked Cudweed        |        |          |      |                        |                      |                  |
| <i>Glossocardia bidens</i> syn. <i>Glossogyne tannensis</i>         | Cobbler's Tack        |        |          |      |                        |                      |                  |
| * <i>Hypochaeris radicata</i>                                       | Catsear, Flatweed     |        |          |      |                        |                      |                  |
| * <i>Lactuca serriola</i>                                           | Prickly Lettuce       |        |          |      |                        |                      |                  |
| <i>Lagenophora stipitata</i> (syn. <i>Lagenifera stipitata</i> )    | Blue Bottle-daisy     |        |          |      |                        |                      |                  |
| <i>Olearia elliptica</i>                                            | Sticky Daisy Bush     |        |          |      |                        |                      | 20200310         |
| <i>Ozothamnus diosmifolium</i>                                      | White Dogwood         |        |          |      |                        |                      | Sept             |
| * <i>Senecio madagascariensis</i>                                   | Fireweed              |        |          |      |                        |                      | Sept, Oct        |
| * <i>Sonchus asper</i>                                              | Prickly Sowthistle    |        |          |      |                        |                      |                  |
| * <i>Sonchus oleraceus</i>                                          | Common Sow Thistle    |        |          |      |                        |                      |                  |
| * <i>Tagetes minuta</i>                                             | Stinking Roger        |        |          |      |                        |                      |                  |
| * <i>Taraxacum officinale</i>                                       | Dandelion             |        |          |      |                        |                      |                  |
| * <i>Tragopogon porrifolius</i>                                     | Salsify               |        |          |      |                        |                      |                  |
| <i>Vittadinia cuneata</i> var. <i>cuneata</i>                       | Fuzzweed              |        |          |      |                        |                      | Sept, Oct        |
| * <i>Xanthium occidentale</i>                                       | Noogoora Burr         |        |          |      |                        |                      |                  |
| * <i>Xanthium spinosum</i>                                          | Bathurst Burr         |        |          |      |                        |                      |                  |
| <b>Basellaceae</b>                                                  |                       |        |          |      |                        |                      |                  |
| * <i>Anredera cordifolia</i>                                        | Madeira Vine          |        |          |      |                        |                      |                  |
| <b>Bignoniaceae</b>                                                 |                       |        |          |      |                        |                      |                  |
| * <i>Dolichandra unguis-cati</i> syn. <i>Macfadyena unguis-cati</i> | Cats Claw Creeper     |        |          |      |                        |                      |                  |
| <i>Pandorea pandorana</i>                                           | Wonga-wonga Vine      |        |          |      |                        |                      | Aug, Sept        |
| <b>Boraginaceae</b>                                                 |                       |        |          |      |                        |                      |                  |
| * <i>Echium plantagineum</i>                                        | Paterson's Curse      |        |          |      |                        |                      |                  |
| * <i>Heliotropium amplexicaule</i>                                  | Blue Heliotrope       |        |          |      |                        |                      |                  |

| SCIENTIFIC NAME                                            | COMMON NAME                | BC ACT | EPBC ACT | SAIL | REGIONALLY SIGNIFICANT | BIOSECURITY ACT 2015 | FLOWERING PERIOD |
|------------------------------------------------------------|----------------------------|--------|----------|------|------------------------|----------------------|------------------|
| <b>Brassicaceae</b>                                        |                            |        |          |      |                        |                      |                  |
| * <i>Capsella bursa-pastoris</i>                           | Shepherd's Purse           |        |          |      |                        |                      |                  |
| * <i>Lepidium africanum</i>                                | Peppercress                |        |          |      |                        |                      |                  |
| * <i>Rapistrum rugosum</i>                                 | Turnip Weed, Giant Mustard |        |          |      |                        |                      |                  |
|                                                            |                            |        |          |      |                        |                      |                  |
| <b>Cactaceae</b>                                           |                            |        |          |      |                        |                      |                  |
| * <i>Opuntia aurantiaca</i>                                | Tiger Pear                 |        |          |      |                        |                      |                  |
| * <i>Opuntia stricta</i>                                   | Prickly Pear               |        |          |      |                        |                      |                  |
|                                                            |                            |        |          |      |                        |                      |                  |
| <b>Campanulaceae</b>                                       |                            |        |          |      |                        |                      |                  |
| <i>Lobelia concolor</i>                                    | Poison Pratia              |        |          |      |                        |                      |                  |
| <i>Lobelia purpurascens</i>                                | White Root                 |        |          |      |                        |                      |                  |
| <i>Wahlenbergia communis</i>                               | Native Bluebell            |        |          |      |                        |                      |                  |
| <i>Wahlenbergia gracilis</i>                               | Sprawling Bluebell         |        |          |      |                        |                      |                  |
|                                                            |                            |        |          |      |                        |                      |                  |
| <b>Caryophyllaceae</b>                                     |                            |        |          |      |                        |                      |                  |
| * <i>Cerastium glomeratum</i>                              | Mouse Ear Chickweed        |        |          |      |                        |                      |                  |
| * <i>Paronychia brasiliensis</i>                           | Brazilian Whitlow          |        |          |      |                        |                      |                  |
| * <i>Petrorhagia nanteuillii</i>                           | Proliferous Pink           |        |          |      |                        |                      | Sept, Oct        |
| * <i>Stellaria media</i>                                   | Common Chickweed           |        |          |      |                        |                      | Aug, Sept        |
|                                                            |                            |        |          |      |                        |                      |                  |
| <b>Cassythaceae</b>                                        |                            |        |          |      |                        |                      |                  |
| <i>Cassytha pubescens</i>                                  | Common Devils Twine        |        |          |      |                        |                      |                  |
|                                                            |                            |        |          |      |                        |                      |                  |
| <b>Casuarinaceae</b>                                       |                            |        |          |      |                        |                      |                  |
| <i>Allocasuarina littoralis</i>                            | Black She-oak              |        |          |      |                        |                      |                  |
| <i>Allocasuarina luehmannii</i>                            | Bulloak                    |        |          |      |                        |                      |                  |
| <i>Allocasuarina torulosa</i>                              | Forest Oak                 |        |          |      |                        |                      |                  |
| <i>Casuarina cunninghamiana</i> ssp. <i>cunninghamiana</i> | River She-oak              |        |          |      |                        |                      |                  |
| <i>Casuarina glauca</i>                                    | Swamp She-oak              |        |          |      |                        |                      |                  |
|                                                            |                            |        |          |      |                        |                      |                  |
| <b>Celastraceae</b>                                        |                            |        |          |      |                        |                      |                  |
| <i>Denhamia silvestris</i> syn. <i>Maytenus silvestris</i> | Narrow leaved Orangebark   |        |          |      | U                      |                      | Oct, April       |
|                                                            |                            |        |          |      |                        |                      |                  |
|                                                            |                            |        |          |      |                        |                      |                  |
| <b>Chenopodiaceae</b>                                      |                            |        |          |      |                        |                      |                  |
| * <i>Chenopodium alba</i>                                  | Fat Hen                    |        |          |      |                        |                      |                  |

| SCIENTIFIC NAME                              | COMMON NAME            | BC ACT | EPBC ACT | SAIL | REGIONALLY SIGNIFICANT | BIOSECURITY ACT 2015 | FLOWERING PERIOD |
|----------------------------------------------|------------------------|--------|----------|------|------------------------|----------------------|------------------|
| <i>Einadia hastata</i>                       | Berry Saltbush         |        |          |      |                        |                      |                  |
| <i>Einadia nutans</i>                        | Nodding Saltbush       |        |          |      |                        |                      |                  |
| <i>Enchylaena tomentosa</i>                  | Ruby Saltbush          |        |          |      |                        |                      |                  |
| <i>Maireana microphylla</i>                  | Small-leaf Bluebush    |        |          |      | E?                     |                      |                  |
| <i>Sclerolaena muricata</i>                  | Black Roly Poly        |        |          |      |                        |                      |                  |
|                                              |                        |        |          |      |                        |                      |                  |
| <b>Convolvulaceae</b>                        |                        |        |          |      |                        |                      |                  |
| <i>Convolvulus erubescens</i>                | Australian Bindweed    |        |          |      |                        |                      |                  |
| <i>Dichondra repens</i>                      | Kidney Weed            |        |          |      |                        |                      |                  |
| <i>Polymeria calycina</i>                    | Swamp Bindweed         |        |          |      |                        |                      |                  |
|                                              |                        |        |          |      |                        |                      |                  |
| <b>Crassulaceae</b>                          |                        |        |          |      |                        |                      |                  |
| <i>*Bryophyllum delagoense</i>               | Mother of Millions     |        |          |      |                        |                      | Noxious weed     |
| <i>Crassula sieberana</i>                    | Australian Stonecrop   |        |          |      |                        |                      |                  |
|                                              |                        |        |          |      |                        |                      |                  |
| <b>Dilleniaceae</b>                          |                        |        |          |      |                        |                      |                  |
| <i>Hibbertia aspera</i>                      | Rough Guinea Flower    |        |          |      |                        |                      | Nov              |
| <i>Hibbertia diffusa</i>                     | Guinea Flower          |        |          |      |                        |                      |                  |
| <i>Hibbertia linearis</i>                    | Guinea Flower          |        |          |      |                        |                      |                  |
| <i>Hibbertia obtusifolia</i>                 | Hoary Guinea Flower    |        |          |      |                        |                      |                  |
| <i>Hibbertia pedunculata</i>                 |                        |        |          |      |                        |                      |                  |
| <i>Hibbertia scandens</i>                    | Climbing Guinea Flower |        |          |      |                        |                      | Sept, Oct        |
|                                              |                        |        |          |      |                        |                      |                  |
| <b>Droseraceae</b>                           |                        |        |          |      |                        |                      |                  |
| <i>Drosera peltata</i> subsp. <i>peltata</i> | Sundew                 |        |          |      |                        |                      |                  |
|                                              |                        |        |          |      |                        |                      |                  |
| <b>Ericaceae formerly Epacridaceae</b>       |                        |        |          |      |                        |                      |                  |
| <i>Acrotriche divaricata</i>                 | Ground Berry           |        |          |      | N                      |                      |                  |
| <i>Epacris pulchella</i>                     | NSW Coral Heath        |        |          |      |                        |                      | April, May       |
| <i>Leucopogon juniperinus</i>                | Prickly Bearded Heath  |        |          |      |                        |                      | July, Aug        |
| <i>Lissanthe strigosa</i>                    | Native Cranberry       |        |          |      |                        |                      |                  |
| <i>Melichrus urceolatus</i>                  | Urn Heath              |        |          |      |                        |                      |                  |
|                                              |                        |        |          |      |                        |                      |                  |
| <b>Euphorbiaceae</b>                         |                        |        |          |      |                        |                      |                  |
| <i>Euphorbia drummondii</i>                  | Caustic Weed           |        |          |      |                        |                      |                  |
| <i>*Euphorbia peplus</i>                     | Petty Spurge           |        |          |      |                        |                      |                  |
| <i>*Ricinus communis</i>                     | Castor Oil Plant       |        |          |      |                        |                      |                  |



| SCIENTIFIC NAME                                             | COMMON NAME           | BC ACT  | EPBC ACT | SAII | REGIONALLY SIGNIFICANT | BIOSECURITY ACT 2015 | FLOWERING PERIOD |
|-------------------------------------------------------------|-----------------------|---------|----------|------|------------------------|----------------------|------------------|
| <b>Fabaceae Subfamily (Caesalpinioideae)</b>                |                       |         |          |      |                        |                      |                  |
| <i>*Senna pendula</i> var. <i>glabrata</i>                  |                       |         |          |      |                        |                      |                  |
| <b>Fabaceae Subfamily (Faboideae)</b>                       |                       |         |          |      |                        |                      |                  |
| <i>Chorizema parviflorum</i>                                | Eastern Flame Pea     |         |          |      |                        |                      |                  |
| <i>Daviesia genistifolia</i>                                | Broom Bitter Pea      |         |          |      |                        |                      |                  |
| <i>Daviesia ulicifolia</i>                                  | Gorse Bitter Pea      |         |          |      |                        |                      | Aug              |
| <i>Desmodium varians</i>                                    | Slender Tick-trefoil  |         |          |      |                        |                      |                  |
| <i>Dillwynia retorta</i> subsp. <i>retorta</i>              | Heathy Parrot Pea     |         |          |      |                        |                      | July, Aug, Sept  |
| <i>Glycine clandestina</i> subsp. complex                   | Love Creeper          |         |          |      |                        |                      | Sept             |
| <i>Glycine tabacina</i> sp. complex                         | Love Creeper          |         |          |      |                        |                      |                  |
| <i>Gompholobium pinnatum</i>                                |                       |         |          |      |                        |                      |                  |
| <i>Hardenbergia violacea</i>                                | False Sarsaparilla    |         |          |      |                        |                      | Aug, Sept        |
| <i>Indigofera australis</i>                                 | Austral Indigo        |         |          |      |                        |                      | July, Aug        |
| <i>Jacksonia scoparia</i>                                   | Dogwood               |         |          |      |                        |                      | Sept, Oct        |
| <i>Kennedia rubicunda</i>                                   | Dusky Coral Pea       |         |          |      |                        |                      | Sept, Oct        |
| <i>*Medicago polymorpha</i>                                 | Burr Medic            |         |          |      |                        |                      |                  |
| <i>*Medicago sativa</i>                                     | Lucerne, Alfalfa      |         |          |      |                        |                      |                  |
| <i>*Melilotus indicus</i>                                   | Hexham Scent          |         |          |      |                        |                      |                  |
| <i>Pultenaea microphylla</i>                                |                       |         |          |      |                        |                      |                  |
| <i>Pultenaea spinosa</i> syn. <i>Pultenaea cunninghamii</i> | Spiny Bush-pea        |         |          |      |                        |                      | Sept             |
| <i>Templetonia stenophylla</i>                              | Leafy Templetonia     |         |          |      |                        |                      | Fruit Sept       |
| <i>*Trifolium arvense</i>                                   | Haresfoot Clover      |         |          |      |                        |                      |                  |
| <i>*Trifolium campestre</i>                                 | Hop Clover            |         |          |      |                        |                      | Sept, Oct        |
| <i>*Trifolium repens</i>                                    | White Clover          |         |          |      |                        |                      | Sept, Oct        |
| <i>*Vicia sativa</i>                                        | Common Vetch          |         |          |      |                        |                      |                  |
| <i>Zornia dyctiocarpa</i> var. <i>dyctiocarpa</i>           | Zornia                |         |          |      |                        |                      |                  |
| <b>Fabaceae (Subfamily Mimosoideae)</b>                     |                       |         |          |      |                        |                      |                  |
| <i>Acacia decora</i>                                        | Western Silver Wattle |         |          |      | U*                     |                      |                  |
| <i>Acacia elata</i>                                         | Cedar Wattle          |         |          |      |                        |                      |                  |
| <i>Acacia elongata</i>                                      | Swamp Wattle          |         |          |      |                        |                      | Aug              |
| <i>Acacia falcata</i>                                       | Falcata Wattle        |         |          |      | W                      |                      |                  |
| <i>Acacia implexa</i>                                       | Hickory               |         |          |      |                        |                      |                  |
| <i>Acacia longifolia</i> subsp. <i>longifolia</i>           | Sydney Golden Wattle  |         |          |      |                        |                      | July, Aug        |
| <i>Acacia parvipinnula</i>                                  | Silver-stemmed Wattle |         |          |      |                        |                      |                  |
| <i>Acacia pendula</i>                                       | Weeping Myall         | E - pop |          |      | E R T*                 |                      |                  |

| SCIENTIFIC NAME                 | COMMON NAME              | BC ACT | EPBC ACT | SAII | REGIONALLY SIGNIFICANT | BIOSECURITY ACT 2015 | FLOWERING PERIOD |
|---------------------------------|--------------------------|--------|----------|------|------------------------|----------------------|------------------|
| <i>Acacia podalyriifolia</i>    | Queensland Silver Wattle |        |          |      |                        |                      |                  |
| <i>Acacia salicina</i>          | Cooba                    |        |          |      | E                      |                      | Mar,             |
| <i>Acacia saligna</i>           | Golden Wreath Wattle     |        |          |      |                        |                      | Aug, Sept        |
| <i>Acacia suaveolens</i>        | Sweet-scented Wattle     |        |          |      |                        |                      |                  |
| <i>Acacia ulicifolia</i>        | Prickly Moses            |        |          |      |                        |                      | Mar, Apr         |
| <i>Acacia uncinata</i>          | Gold Dust Wattle         |        |          |      |                        |                      |                  |
| <b>Gentianaceae</b>             |                          |        |          |      |                        |                      |                  |
| * <i>Cenaurium erythraea</i>    | Common Centaury          |        |          |      |                        |                      |                  |
| <b>Geraniaceae</b>              |                          |        |          |      |                        |                      |                  |
| <i>Erodium crinitum</i>         | Blue Heronsbill          |        |          |      |                        |                      | Sept, Oct        |
| <i>Geranium homeanum</i>        | Cranesbill               |        |          |      |                        |                      |                  |
| <i>Geranium solanderi</i>       | Native Geranium          |        |          |      |                        |                      |                  |
| <b>Goodeniaceae</b>             |                          |        |          |      |                        |                      |                  |
| <i>Goodenia heterophylla</i>    | Variable-leaved Goodenia |        |          |      |                        |                      |                  |
| <i>Goodenia ovata</i>           | Hop Goodenia             |        |          |      |                        |                      | Sept             |
| <i>Goodenia pinnatifida</i>     |                          |        |          |      |                        |                      |                  |
| <i>Goodenia rotundifolia</i>    |                          |        |          |      |                        |                      |                  |
| <b>Haloragaceae</b>             |                          |        |          |      |                        |                      |                  |
| <i>Gonocarpus tetragynus</i>    | Poverty Raspwort         |        |          |      |                        |                      |                  |
| <b>Hypericaceae</b>             |                          |        |          |      |                        |                      |                  |
| <i>Hypericum gramineum</i>      | Native St John's Wort    |        |          |      |                        |                      |                  |
| <i>Hypericum perforatum</i>     | St John's Wort           |        |          |      |                        |                      |                  |
| <b>Lamiaceae</b>                |                          |        |          |      |                        |                      |                  |
| <i>Mentha satureioides</i>      | Native Pennyroyal        |        |          |      |                        |                      |                  |
| <i>Plectranthus parviflorus</i> | Cockspur Flower          |        |          |      |                        |                      |                  |
| * <i>Stachys arvensis</i>       | Stagger Weed             |        |          |      |                        |                      |                  |
| <b>Lauraceae</b>                |                          |        |          |      |                        |                      |                  |
| * <i>Cinnamomum camphora</i>    | Camphor Laurel           |        |          |      |                        |                      |                  |
| <b>Linaceae</b>                 |                          |        |          |      |                        |                      |                  |
| <i>Linum marginale</i>          | Native Flax              |        |          |      |                        |                      | Sept             |

| SCIENTIFIC NAME                                       | COMMON NAME               | BC ACT | EPBC ACT | SAIL | REGIONALLY SIGNIFICANT | BIOSECURITY ACT 2015 | FLOWERING PERIOD |
|-------------------------------------------------------|---------------------------|--------|----------|------|------------------------|----------------------|------------------|
| <i>*Linum trigynum</i>                                | French Flax               |        |          |      |                        |                      |                  |
| <b>Loranthaceae</b>                                   |                           |        |          |      |                        |                      |                  |
| <i>Amyema miquelii</i>                                | Box Mistletoe             |        |          |      |                        |                      |                  |
| <i>Amyema pendula</i>                                 | Drooping Mistletoe        |        |          |      |                        |                      |                  |
| <i>Dendrophthoe vitellina</i>                         | Apostle Mistletoe         |        |          |      |                        |                      | Sept, Oct, Nov   |
| <b>Lythraceae</b>                                     |                           |        |          |      |                        |                      |                  |
| <i>Lythrum hyssopifolia</i>                           | Hyssop Loosestrife        |        |          |      |                        |                      |                  |
| <b>Malvaceae</b>                                      |                           |        |          |      |                        |                      |                  |
| <i>Brachychiton populneus</i> subsp. <i>populneus</i> | Kurrajong                 |        |          |      |                        |                      |                  |
| <i>*Malva parviflora</i>                              | Small-flowered Mallow     |        |          |      |                        |                      | Sept             |
| <i>*Modiola carliniana</i>                            | Red-flowered Mallow       |        |          |      |                        |                      |                  |
| <i>*Pavonia hastata</i>                               | Pink Parvonia             |        |          |      |                        |                      |                  |
| <i>Sida corrugata</i>                                 | Corrugated Sida           |        |          |      |                        |                      |                  |
| <i>Sida hackettiana</i> syn <i>Sida subspicata</i>    | Golden Rod, Spiked Sida   |        |          |      |                        |                      |                  |
| <i>*Sida rhombifolia</i>                              | Paddys Lucerne            |        |          |      |                        |                      |                  |
| <b>Meliaceae</b>                                      |                           |        |          |      |                        |                      |                  |
| <i>Melia azedarach</i> var. <i>australasica</i>       | White Cedar               |        |          |      |                        |                      |                  |
| <b>Myrtaceae</b>                                      |                           |        |          |      |                        |                      |                  |
| <i>Angophora bakeri</i>                               | Narrow-leaved Apple       |        |          |      |                        |                      | Nov, Dec         |
| <i>Angophora floribunda</i>                           | Rough-barked Apple        |        |          |      |                        |                      |                  |
| <i>Callistemon linearifolius</i>                      | Netted Bottlebrush?       | V      |          |      |                        |                      |                  |
| <i>Callistemon linearis</i>                           | Narrow-leaved Bottlebrush |        |          |      |                        |                      |                  |
| <i>Callistemon rigidus</i>                            | Stiff Bottlebrush         |        |          |      |                        |                      |                  |
| <i>Callistemon salignus</i>                           | Willow Bottlebrush        |        |          |      |                        |                      |                  |
| <i>Callistemon viminalis</i>                          | Weeping Bottlebrush       |        |          |      |                        |                      |                  |
| <i>Calytrix tetragona</i>                             | Common Fringe-myrtle      |        |          |      |                        |                      |                  |
| <i>Corymbia maculata</i>                              | Spotted Gum               |        |          |      |                        |                      | Mar, Apr         |
| <i>Eucalyptus amplifolia</i> subsp. <i>amplifolia</i> | Cabbage Gum               |        |          |      |                        |                      |                  |
| <i>Eucalyptus blakelyi</i>                            | Blakely's Red Gum         |        |          |      |                        |                      |                  |
| <i>Eucalyptus crebra</i>                              | Narrow-leaved Ironbark    |        |          |      |                        |                      | Sept, Oct        |
| <i>Eucalyptus fibrosa</i> subsp. <i>fibrosa</i>       | Broad-leaved Ironbark     |        |          |      |                        |                      |                  |
| <i>Eucalyptus glaucina</i>                            | Slaty Red Gum             | V      | V        |      |                        |                      |                  |
| <i>Eucalyptus globoidea</i>                           | White Stringybark         |        |          |      |                        |                      |                  |



| SCIENTIFIC NAME                                                 | COMMON NAME                 | BC ACT | EPBC ACT | SAII | REGIONALLY SIGNIFICANT | BIOSECURITY ACT 2015 | FLOWERING PERIOD    |
|-----------------------------------------------------------------|-----------------------------|--------|----------|------|------------------------|----------------------|---------------------|
| <i>Eucalyptus moluccana</i>                                     | Grey Box                    |        |          |      |                        |                      | Mar                 |
| <i>Eucalyptus parramattensis</i> subsp. <i>decadens</i>         | Drooping Red Gum            | V      | V        |      |                        |                      |                     |
| <i>Eucalyptus punctata</i>                                      | Grey Gum                    |        |          |      |                        |                      |                     |
| <i>Eucalyptus siderophloia</i>                                  | Grey Ironbark               |        |          |      |                        |                      |                     |
| <i>Eucalyptus sideroxylon</i>                                   | Mugga                       |        |          |      |                        |                      |                     |
| <i>Eucalyptus tereticornis</i>                                  | Forest Red Gum              |        |          |      |                        |                      | May, Aug, Sept, Oct |
| <i>Eucalyptus umbra</i>                                         | Thick-leaved White Mahogany |        |          |      |                        |                      |                     |
| <i>Euryomyrtus ramosissima</i> subsp. <i>ramosissima</i>        | Rosy Baeckea                |        |          |      |                        |                      | Sept, Oct           |
| <i>Gaudium multicaule</i> syn <i>Leptospermum multicaule</i>    | Silver Tea-tree             |        |          |      |                        |                      |                     |
| <i>Kunzea ambigua</i>                                           | Tick Bush                   |        |          |      |                        |                      | Oct, Nov            |
| <i>Leptospermum polygalifolium</i> subsp. <i>cismontanum</i>    |                             |        |          |      |                        |                      |                     |
| <i>Leptospermum polygalifolium</i> subsp. <i>polygalifolium</i> | Teatree                     |        |          |      |                        |                      | Sept                |
| <i>Lophostemon confertus</i>                                    |                             |        |          |      |                        |                      |                     |
| <i>Melaleuca decora</i>                                         | White Feather Honeymyrtle   |        |          |      |                        |                      | Nov, Dec            |
| <i>Melaleuca linearifolia</i>                                   | Snow in Summer              |        |          |      |                        |                      | Oct, Nov            |
| <i>Melaleuca nodosa</i>                                         | Ball Honeymyrtle            |        |          |      |                        |                      | Late Sept, Oct      |
| <i>Melaleuca quinquenervia</i>                                  | Broad-leaved Paperbark      |        |          |      |                        |                      | Feb, Mar            |
| <i>Melaleuca styphelioides</i>                                  | Prickly-leaved Paperbark    |        |          |      |                        |                      |                     |
| <b>Oleaceae</b>                                                 |                             |        |          |      |                        |                      |                     |
| * <i>Ligustrum sinense</i>                                      | Small-leaved Privet         |        |          |      |                        |                      | Sept                |
| <i>Notelaea longifolia</i>                                      | Mock Olive                  |        |          |      |                        |                      |                     |
| <i>Notelaea microcarpa</i>                                      | Native Olive                |        |          |      |                        |                      |                     |
| * <i>Olea europaea</i> subsp. <i>cuspidata</i>                  | African Olive               |        |          |      |                        |                      |                     |
| <b>Onagraceae</b>                                               |                             |        |          |      |                        |                      |                     |
| <i>Ludwigia peploides</i> subsp. <i>montevidensis</i>           | Water Primrose              |        |          |      |                        |                      |                     |
| <b>Oxalidaceae</b>                                              |                             |        |          |      |                        |                      |                     |
| <i>Oxalis corniculata</i>                                       | Creeping Oxalis             |        |          |      |                        |                      |                     |
| <i>Oxalis perennans</i>                                         | -                           |        |          |      |                        |                      |                     |
| <b>Phyllanthaceae</b>                                           |                             |        |          |      |                        |                      |                     |

| SCIENTIFIC NAME                                             | COMMON NAME              | BC ACT | EPBC ACT | SAII | REGIONALLY SIGNIFICANT | BIOSECURITY ACT 2015 | FLOWERING PERIOD |
|-------------------------------------------------------------|--------------------------|--------|----------|------|------------------------|----------------------|------------------|
| <i>Breynia oblongifolia</i>                                 | Coffee Bush              |        |          |      |                        |                      |                  |
| <i>Phyllanthus hirtellus</i>                                | Thyme Spurge             |        |          |      |                        |                      |                  |
| <i>Poranthera microphylla</i>                               |                          |        |          |      |                        |                      |                  |
|                                                             |                          |        |          |      |                        |                      |                  |
| <b>Pittosporaceae</b>                                       |                          |        |          |      |                        |                      |                  |
| <i>Billardiera scandens</i>                                 | Apple Dumplings          |        |          |      |                        |                      | Sept             |
| <i>Bursaria spinosa</i> subsp. <i>spinosa</i>               | Blackthorn               |        |          |      |                        |                      | Jan              |
| <i>Pittosporum undulatum</i>                                | Sweet Pittosporum        |        |          |      |                        |                      | Aug, Sept        |
|                                                             |                          |        |          |      |                        |                      |                  |
| <b>Plantaginaceae</b>                                       |                          |        |          |      |                        |                      |                  |
| <i>Plantago debilis</i>                                     |                          |        |          |      |                        |                      |                  |
| * <i>Plantago lanceolata</i>                                | Plantain                 |        |          |      |                        |                      |                  |
|                                                             |                          |        |          |      |                        |                      |                  |
| <b>Polygonaceae</b>                                         |                          |        |          |      |                        |                      |                  |
| <i>Persicaria decipens</i>                                  | Slender Knotweed         |        |          |      |                        |                      |                  |
| * <i>Polygonum aviculare</i>                                | Wireweed                 |        |          |      |                        |                      |                  |
| <i>Rumex brownii</i>                                        | Swamp Dock               |        |          |      |                        |                      |                  |
| * <i>Rumex crispus</i>                                      | Curled Dock              |        |          |      |                        |                      |                  |
| * <i>Rumex sagittatus</i> syn. <i>Acetosa sagittata</i>     | Turkey Rhubarb           |        |          |      |                        |                      |                  |
|                                                             |                          |        |          |      |                        |                      |                  |
| <b>Portulacaceae</b>                                        |                          |        |          |      |                        |                      |                  |
| <i>Portulaca oleracea</i>                                   | Purslane, Pigweed        |        |          |      |                        |                      |                  |
|                                                             |                          |        |          |      |                        |                      |                  |
| <b>Primulaceae</b>                                          |                          |        |          |      |                        |                      |                  |
| * <i>Lysimachia arvensis</i> syn. <i>Anagallis arvensis</i> | Scarlet Pimpernel        |        |          |      |                        |                      |                  |
|                                                             |                          |        |          |      |                        |                      |                  |
| <b>Proteaceae</b>                                           |                          |        |          |      |                        |                      |                  |
| <i>Banksia collina</i>                                      |                          |        |          |      |                        |                      | Mar              |
| <i>Grevillea montana</i>                                    | Mountain Grevillea       |        |          |      |                        |                      |                  |
| <i>Grevillea parviflora</i> subsp. <i>parviflora</i>        | Small-flowered Grevillea | V      | V        |      |                        |                      | Sept             |
| <i>Grevillea robusta</i>                                    | Silky Oak                |        |          |      |                        |                      |                  |
| <i>Hakea sericea</i>                                        | Needlebush               |        |          |      |                        |                      |                  |
| <i>Isopogon anemonifolius</i>                               | Drumsticks               |        |          |      |                        |                      |                  |
| <i>Lambertia formosa</i>                                    | Mountain Devil           |        |          |      |                        |                      |                  |
| <i>Persoonia linearis</i>                                   | Narrow-leaved Geebung    |        |          |      |                        |                      |                  |
|                                                             |                          |        |          |      |                        |                      |                  |
| <b>Ranunculaceae</b>                                        |                          |        |          |      |                        |                      |                  |
| <i>Clematis aristida</i>                                    | Old Man's Beard          |        |          |      |                        |                      |                  |

| SCIENTIFIC NAME                                     | COMMON NAME         | BC ACT | EPBC ACT | SAIL | REGIONALLY SIGNIFICANT | BIOSECURITY ACT 2015 | FLOWERING PERIOD |
|-----------------------------------------------------|---------------------|--------|----------|------|------------------------|----------------------|------------------|
| <i>Clematis glycinoides</i> var. <i>glycinoides</i> | Headache Vine       |        |          |      |                        |                      | Aug, Sept        |
| <i>Ranunculus inundatus</i>                         | River Buttercup     |        |          |      |                        |                      |                  |
| <i>Ranunculus lappaceus</i>                         | Common Buttercup    |        |          |      |                        |                      |                  |
| <b>Rhamnaceae</b>                                   |                     |        |          |      |                        |                      |                  |
| <i>Pomaderris lanigera</i>                          | Woolly Pomaderris   |        |          |      |                        |                      |                  |
| <b>Rosaceae</b>                                     |                     |        |          |      |                        |                      |                  |
| * <i>Rubus fruticosus</i> ssp. <i>aggregate</i>     | Blackberry          |        |          |      |                        |                      |                  |
| <b>Rubiaceae</b>                                    |                     |        |          |      |                        |                      |                  |
| <i>Asperula conferta</i>                            | Common Woodruff     |        |          |      |                        |                      |                  |
| * <i>Galium aparine</i>                             | Clevers             |        |          |      |                        |                      |                  |
| <i>Opercularia diphylla</i>                         | Stinkweed           |        |          |      |                        |                      |                  |
| <i>Pomax umbellata</i>                              | Pomax               |        |          |      |                        |                      |                  |
| * <i>Richardia humistrata</i>                       |                     |        |          |      |                        |                      |                  |
| <b>Rutaceae</b>                                     |                     |        |          |      |                        |                      |                  |
| <i>Boronia polygalifolia</i>                        | Dwarf Boronia       |        |          |      |                        |                      |                  |
| <i>Geijera parviflora</i>                           | Wilga               |        |          |      |                        |                      |                  |
| <i>Zieria smithii</i>                               | Sandfly Zieria      |        |          |      |                        |                      | Sept, Oct        |
| <b>Santalaceae</b>                                  |                     |        |          |      |                        |                      |                  |
| <i>Choretrum candollei</i>                          | White Sour Bush     |        |          |      |                        |                      |                  |
| <i>Exocarpus cupressiformis</i>                     | Cherry Ballart      |        |          |      |                        |                      |                  |
| <b>Sapindaceae</b>                                  |                     |        |          |      |                        |                      |                  |
| * <i>Cardiospermum grandiflorum</i>                 | Balloon Vine        |        |          |      |                        |                      |                  |
| <i>Dodonaea viscosa</i> subsp. <i>cuneata</i>       | Wedge-leaf Hop Bush |        |          |      |                        |                      |                  |
| <i>Dodonaea triquetra</i>                           | Hop Bush            |        |          |      |                        |                      |                  |
| <b>Scrophulariaceae</b>                             |                     |        |          |      |                        |                      |                  |
| <i>Eremophila debilis</i>                           | Amulla              |        |          |      |                        |                      |                  |
| <i>Myoporum montanum</i>                            | Western Boobialla   |        |          |      |                        |                      |                  |
| * <i>Verbascum virgatum</i>                         | Twiggy Mullein      |        |          |      |                        |                      |                  |
| <i>Veronica plebeia</i>                             | Speedwell           |        |          |      |                        |                      |                  |
| <b>Solanaceae</b>                                   |                     |        |          |      |                        |                      |                  |



| SCIENTIFIC NAME                           | COMMON NAME           | BC ACT | EPBC ACT | SAII | REGIONALLY SIGNIFICANT | BIOSECURITY ACT 2015 | FLOWERING PERIOD |
|-------------------------------------------|-----------------------|--------|----------|------|------------------------|----------------------|------------------|
| <i>*Cestrum parqui</i>                    | Green Cestrum         |        |          |      |                        |                      | Sept, Oct        |
| <i>*Datura stramonium</i>                 | Common Thornapple     |        |          |      |                        |                      |                  |
| <i>*Lycium ferocissimum</i>               | African Boxthorn      |        |          |      |                        |                      |                  |
| <i>Solanum cinereum</i>                   | Narrawa Burr          |        |          |      |                        |                      |                  |
| <i>*Solanum mauritianum</i>               | Wild Tobacco          |        |          |      |                        |                      |                  |
| <i>*Solanum nigrum</i>                    | Blackberry Nightshade |        |          |      |                        |                      |                  |
| <i>*Solanum pseudocapsicum</i>            | Jerusalem Cherry      |        |          |      |                        |                      |                  |
| <i>Solanum prinophyllum</i>               | Forest Nightshade     |        |          |      |                        |                      |                  |
| <i>*Solanum seaforthianum</i>             | Climbing Nightshade   |        |          |      |                        |                      | March, April     |
| <i>Solanum stelligerum</i>                | Devils Needle's       |        |          |      |                        |                      |                  |
| <b>Stackhousiaceae</b>                    |                       |        |          |      |                        |                      |                  |
| <i>Stackhousia viminea</i>                | Slender Stackhousia   |        |          |      |                        |                      |                  |
| <b>Stylidiaceae</b>                       |                       |        |          |      |                        |                      |                  |
| <i>Stylidium graminifolium</i>            | Trigger Plant         |        |          |      |                        |                      | Sept             |
| <b>Thymelaeaceae</b>                      |                       |        |          |      |                        |                      |                  |
| <i>Pimelea linifolia</i>                  | Rice Flower           |        |          |      |                        |                      | Sept             |
| <b>Urticaceae</b>                         |                       |        |          |      |                        |                      |                  |
| <i>Urtica incisa</i>                      | Stinging Nettle       |        |          |      |                        |                      |                  |
| <i>*Urtica urens</i>                      | Small Nettle          |        |          |      |                        |                      |                  |
| <b>Verbenaceae</b>                        |                       |        |          |      |                        |                      |                  |
| <i>*Lantana camara</i>                    | Lantana               |        |          |      |                        |                      | Noxious          |
| <i>*Phyla canescens</i>                   | Lippia                |        |          |      |                        |                      | Noxious          |
| <i>*Verbena bonariensis</i>               | Purple Top            |        |          |      |                        |                      |                  |
| <i>Verbena gaudichaudii</i>               |                       |        |          |      |                        |                      |                  |
| <i>*Verbena rigida</i> var. <i>rigida</i> | Veined Verbena        |        |          |      |                        |                      |                  |
| <b>Vitaceae</b>                           |                       |        |          |      |                        |                      |                  |
| <i>Cayratia clematidea</i>                | Slender Grape         |        |          |      |                        |                      |                  |
| <b>Zygophyllaceae</b>                     |                       |        |          |      |                        |                      |                  |
| <i>Tribulus</i> sp.                       | Caltrop               |        |          |      |                        |                      |                  |

## **APPENDIX E**

### **SURVEYED SIGNIFICANT TREE DATA**

**Significant Tree Data Key for Table E1.**

- **\*DBH** – Diameter at Breast Height. Tree trunk diameter measured at breast height (1.4 metres above ground level).
- **\*Tree Height** –(m)
- **Coordinates - GDA – 2020, MGA 56**
- **Habitat/Hollows** –
  - Class 1** – very large sized hollow openings (i.e., >20cm) suitable for species such as Owls
  - Class 2** – large sized hollow openings (i.e., 15-20cm) suitable for species such as Owls and Possums
  - Class 3** – medium sized hollow-openings (i.e., 5-15cm) suitable for species such as Gliders and Possums
  - Class 4** – small sized hollow openings (i.e., <5cm) suitable for species such as microchiropteran bats.
- Spout** - Hollow opening towards sky offering little protection from the weather
- Arboreal Termite Nest** – provides potential nesting opportunities for hollow-dependent birds, such as kingfishers and kookaburras



**Table E1: Details of significant trees likely impacted by the REZ.**

| No. | Species of Tree                 | Easting | Northing | Height (m) | DBH (m)    | Habitat Present |         |         |         | Comments                                                           | Removal Required?        |
|-----|---------------------------------|---------|----------|------------|------------|-----------------|---------|---------|---------|--------------------------------------------------------------------|--------------------------|
|     |                                 |         |          |            |            | Class 1         | Class 2 | Class 3 | Class 4 |                                                                    |                          |
| T1  | Stringy                         | 357791  | 6369986  | 20         | 1.13       |                 |         |         | 1       |                                                                    | Likely retain            |
| T2  | <i>A. floribunda</i>            | 357766  | 6370028  |            | 1.02       |                 |         | 1?      |         | Large scar in trunk                                                | Likely retain            |
| T3  | <i>A. floribunda</i>            | 357754  | 6370028  |            | 1.11       |                 |         |         |         | Branch leaning into easement                                       | Likely retain            |
| T4  |                                 | 354258  | 6373718  | 17         | 0.42       |                 |         |         | 1       | Dead branches. Scar in trunk. Hollow suitable for bats and lizards | Likely removal           |
| T5  | <i>E. tereticornis</i>          | 349586  | 6376906  | 14         | 0.45       |                 |         |         |         |                                                                    | Likely removal           |
| T6  | <i>E. tereticornis</i>          | 349586  | 6376900  | 15         | 0.42       |                 |         |         |         |                                                                    | Likely removal           |
| T7  | <i>E. tereticornis</i>          | 349576  | 6376927  | 15         | 0.77       |                 |         |         |         |                                                                    | Likely removal           |
| T8  | <i>C. maculata</i>              | 349136  | 6377953  | 15         | 0.45       |                 |         |         |         |                                                                    | Possible branch trimming |
| T9  | <i>E. fibrosa</i>               | 347684  | 6382740  | 16         | 0.63, 0.70 |                 |         |         |         |                                                                    | Possible branch trimming |
| T10 | <i>E. tereticornis</i>          | 346839  | 6383916  | 18         | 0.83       |                 |         | 1       |         |                                                                    | Possible branch trimming |
| T11 | <i>C. maculata</i>              | 344329  | 6385055  | 22         | 1.03       |                 |         | 1?      |         | Scar at base                                                       | Possible branch trimming |
| T12 | <i>E. tereticornis</i>          | 342397  | 6385186  |            | 1.08       |                 | 3       | 4       | 3       | Large scratches on trunk                                           | Possible branch trimming |
| T13 | <i>E. moluccana</i>             | 342002  | 6385113  | 18         | 1.13       |                 |         |         | 1       |                                                                    | Possible branch trimming |
| T14 | <i>E. moluccana</i>             | 341946  | 6385101  | 17         | 0.89       |                 | 1       | 2       | 2       |                                                                    | Possible branch trimming |
| T15 | <i>E. moluccana</i>             | 340546  | 6385062  | 13         | 0.73       |                 |         |         |         | Hollow 4m up trunk                                                 | Near original route      |
| T18 | <i>C. maculata</i>              | 336972  | 6385823  |            | 1.0        | 1               |         |         |         |                                                                    | Likely removal           |
| T19 | <i>E. crebra</i>                | 319014  | 6399108  |            | 0.9        |                 |         | 2       | 2       | Cracks in trunk                                                    | Likely removal           |
| T20 | <i>E. moluccana</i>             | 319078  | 6399191  | 18         | 0.86       |                 |         |         | 1       |                                                                    | Likely removal           |
| T21 | <i>Allocasuarina luehmannii</i> | 303476  | 6430401  | 10         | 0.47       |                 |         |         |         |                                                                    | Likely retain            |
| T22 | <i>E. blakelyi</i>              | 304229  | 6430944  | 8          | 0.2        |                 |         |         |         |                                                                    | Likely removal           |
| T23 | <i>E. blakelyi</i>              | 305917  | 6432402  | 20         | 1.06       |                 |         | 2       | 1       |                                                                    | Likely removal           |
| T24 | <i>E. parramattensis</i> subsp. | 359373  | 6369253  | 7          | 0.52       |                 | 1       | 1       | 1       |                                                                    | Likely removal           |

| No. | Species of Tree        | Easting | Northing | Height (m) | DBH (m)    | Habitat Present |         |         |         | Comments                                                                      | Removal Required?   |
|-----|------------------------|---------|----------|------------|------------|-----------------|---------|---------|---------|-------------------------------------------------------------------------------|---------------------|
|     |                        |         |          |            |            | Class 1         | Class 2 | Class 3 | Class 4 |                                                                               |                     |
|     | <i>decadens</i>        |         |          |            |            |                 |         |         |         |                                                                               |                     |
| T25 | <i>E. blakelyi</i>     | 305909  | 6432385  | 20         | 1.01       | 1               | 3       |         | 3       |                                                                               | Likely removal      |
| T26 | <i>A. floribunda</i>   | 326167  | 6388110  | 16         | 0.81       |                 |         |         |         |                                                                               | Likely removal      |
| T27 | <i>E. crebra</i>       | 326052  | 6388147  | 18         | 1.27       |                 |         |         |         | Cracks in trunk                                                               | Likely removal      |
| T28 | <i>E. crebra</i>       | 325938  | 6388179  | 15         | 0.69       |                 |         |         |         |                                                                               | Likely removal      |
| T29 | <i>E. crebra</i>       | 325892  | 6388206  |            | 0.78       |                 |         |         |         | Dead, hollow trunk leaning against tree                                       | Likely removal      |
| T30 | <i>E. moluccana</i>    | 325555  | 6388366  | 11         | 0.6        | 1               | 1       |         | 1       |                                                                               | Likely removal      |
| T31 | <i>E. crebra</i>       | 325562  | 6388369  | 16         | 0.83       |                 |         |         |         | Cracks in base of tree. Broken off branch                                     | Likely removal      |
| T32 | <i>E. moluccana</i>    | 325537  | 6388381  |            | 0.94       |                 |         | 2       | 2       |                                                                               | Likely removal      |
| T33 | <i>E. moluccana</i>    | 324520  | 6388660  |            | 0.68       |                 |         |         |         |                                                                               | Likely removal      |
| T34 | <i>A. floribunda</i>   | 324187  | 6388779  | 17         | 0.97       |                 |         |         |         |                                                                               | Likely removal      |
| T35 | <i>E. moluccana</i>    | 321847  | 6389673  | 18         | 0.83       |                 |         |         | 1       |                                                                               | Likely removal      |
| T36 | Dead tree              | 321864  | 6389664  |            | 0.9        |                 |         |         | 2?      | Dead <i>E. moluccana</i> . Cracks in trunk                                    | Likely removal      |
| T37 | <i>E. moluccana</i>    | 321843  | 6389844  | 18         | 0.75       |                 |         |         |         |                                                                               | Likely removal      |
| T44 | <i>E. moluccana</i>    | 340168  | 6385115  |            | 0.99       | 1               | 1       | 2       | 3       | Base of tree is within private property and tree is leaning into road reserve | Near original route |
| T45 | <i>E. moluccana</i>    | 340262  | 6385100  | 18         | 0.35, 0.78 |                 |         |         |         | Scar at base of tree                                                          | Near original route |
| T46 | <i>E. fibrosa</i>      | 340348  | 6385082  | 20         | 0.77       |                 |         |         |         |                                                                               | Near original route |
| T47 | <i>C. maculata</i>     | 340373  | 6385073  | 17         | 0.65       |                 |         |         |         |                                                                               | Near original route |
| T48 | Dead tree              | 340744  | 6385424  | 12         | 0.55, 0.55 |                 | 1       | 2       |         |                                                                               | Likely retain       |
| T65 | <i>E. tereticornis</i> | 340738  | 6385427  | 15         | 0.61, 0.63 |                 |         |         |         |                                                                               | Likely retain       |
| T66 | <i>E. tereticornis</i> | 340768  | 6385433  | 7          | 0.17       |                 |         |         |         |                                                                               | Likely removal      |
| T67 | <i>E. fibrosa</i>      | 340928  | 6385238  | 18         | 1.11       |                 |         |         |         |                                                                               | Likely retain       |
| T68 | <i>E. tereticornis</i> | 340853  | 6385322  | 10         | 0.31       |                 |         |         |         |                                                                               | Likely removal      |

| No. | Species of Tree        | Easting | Northing | Height (m) | DBH (m) | Habitat Present |         |         |         | Comments             | Removal Required?        |
|-----|------------------------|---------|----------|------------|---------|-----------------|---------|---------|---------|----------------------|--------------------------|
|     |                        |         |          |            |         | Class 1         | Class 2 | Class 3 | Class 4 |                      |                          |
| T69 | <i>E. moluccana</i>    | 340913  | 6385300  |            | 1.1     |                 | 1       | 1       | 2       | Scar at base of tree | Possible branch trimming |
| T70 | <i>E. moluccana</i>    | 340882  | 6385353  |            | 1.08    |                 | 2       |         |         |                      | Likely retain            |
| T71 | <i>E. fibrosa</i>      | 340896  | 6385276  |            | 0.78    |                 |         |         | 1       |                      | Likely retain            |
| T72 | <i>E. fibrosa</i>      | 340942  | 6385206  | 18         | 0.96    |                 |         |         |         |                      | Likely removal           |
| T73 | <i>E. fibrosa</i>      | 34044   | 6385086  |            | 0.61    |                 |         |         | 1       |                      | Likely removal           |
| T74 | Dead tree              | 340905  | 6385212  | 17         | 0.81    |                 |         |         |         |                      | Likely retain            |
| T77 | <i>E. tereticornis</i> | 338975  | 6385638  | 16         | 0.8     | 1               | 1       | 1       |         |                      | Likely removal           |
| T78 | <i>E. tereticornis</i> | 338945  | 6385643  | 18         | 1.03    | 1               | 1       | 2       | 3       |                      | Likely removal           |
| T79 | <i>E. tereticornis</i> | 338997  | 6385620  | 18         | 0.88    |                 |         | 1       | 1       |                      | Likely retain            |
| T80 | <i>E. tereticornis</i> | 339059  | 6385616  | 17         | 1       |                 |         |         | 1?      |                      | Likely removal           |
| T81 | <i>E. tereticornis</i> | 339085  | 6385603  | 17         | 0.85    |                 |         |         | 1       |                      | Possible branch trimming |
| T82 | <i>E. tereticornis</i> | 339001  | 6385624  | 16         | 0.55    |                 |         |         | 1       |                      | Likely retain            |
| T83 | <i>C. maculata</i>     | 339163  | 6385575  | 22         | 0.81    |                 |         |         | 1?      |                      | Likely retain            |
| T84 | <i>C. maculata</i>     | 339222  | 6385566  |            | 0.56    |                 |         |         | 1       |                      | Likely retain            |
| T85 | <i>E. crebra</i>       | 339219  | 6385562  |            | 0.69    |                 |         |         | 1?      |                      | Likely retain            |
| T86 | <i>E. tereticornis</i> | 338962  | 6385634  |            | 0.85    |                 |         |         | 1?      |                      | Likely removal           |
| T87 | Dead Ironbark          | 338996  | 6385630  | 18         | 0.81    |                 |         |         |         |                      | Likely removal           |



## **APPENDIX F**

### **SIGNIFICANT FLORA SPECIES INFORMATION**

**Table F.1** *Eucalyptus parramattensis* subsp. *decadens* (Drooping Red Gum) recorded within REZ.

| Tree No. | Species                                                 | COORDINATES<br>GDA – 2020<br>Easting | COORDINATES<br>GDA – 2020<br>Northing | *DBH (M) | *Height (M) | Comments<br>Habitat Present                                    | Removal Required?                                    |
|----------|---------------------------------------------------------|--------------------------------------|---------------------------------------|----------|-------------|----------------------------------------------------------------|------------------------------------------------------|
| P1       | <i>Eucalyptus parramattensis</i> subsp. <i>decadens</i> | 359430                               | 6369215                               | 0.14     | 5           |                                                                | Possible Removal                                     |
| P2       | <i>E. parramattensis</i> subsp. <i>decadens</i>         | 359392                               | 6369239                               | 0.36     | 9           |                                                                | Possible Removal                                     |
| P3       | <i>E. parramattensis</i> subsp. <i>decadens</i>         | 359382                               | 6369248                               | 0.19     | 5           |                                                                | Possible Removal                                     |
| P4       | <i>E. parramattensis</i> subsp. <i>decadens</i>         | 359373                               | 6369253                               | 0.52     | 7           | 1 x class 2 hollow<br>1 x class 3 hollow<br>1 x class 4 hollow | Possible Removal. Tree 24 in Significant Tree Table. |
| P5       | <i>E. parramattensis</i> subsp. <i>decadens</i>         | 359373                               | 6369253                               | 0.13     | 4           |                                                                | Possible Removal                                     |
| P6       | <i>E. parramattensis</i> subsp. <i>decadens</i>         | 359369                               | 6369253                               | 0.12     | 4           |                                                                | Possible Removal                                     |
| P7       | <i>E. parramattensis</i> subsp. <i>decadens</i>         | 359374                               | 6369270                               | 0.2      | 5           |                                                                | Possible Removal                                     |
| P8       | <i>E. parramattensis</i> subsp. <i>decadens</i>         | 359374                               | 6369273                               | 0.28     | 8           |                                                                | Possible Removal                                     |
| P9       | <i>E. parramattensis</i> subsp. <i>decadens</i>         | 359361                               | 6369279                               | 0.36     | 9           |                                                                | Possible Removal                                     |
| P10      | <i>E. parramattensis</i> subsp. <i>decadens</i>         | 359172                               | 6369283                               | 0.12     | 4           |                                                                | Possible Removal                                     |
| P11      | <i>E. parramattensis</i> subsp. <i>decadens</i>         | 359170                               | 6369280                               | 0.15     | 5           |                                                                | Possible Removal                                     |
| P12      | <i>E. parramattensis</i> subsp. <i>decadens</i>         | 359145                               | 6369297                               | 0.12     | 4           |                                                                | Possible Removal                                     |
| P13      | <i>E. parramattensis</i> subsp. <i>decadens</i>         | 359141                               | 6369305                               | 0.07     | 2           |                                                                | Possible Removal                                     |
| P14      | <i>E. parramattensis</i> subsp. <i>decadens</i>         | 359132                               | 6369302                               | 0.14     | 3.5         |                                                                | Possible Removal                                     |
| P15      | <i>E. parramattensis</i> subsp. <i>decadens</i>         | 359137                               | 6369303                               | 0.07     | 4           |                                                                | Possible Removal                                     |
| P16      | <i>E. parramattensis</i> subsp. <i>decadens</i>         | 359134                               | 6369305                               | 0.13     | 3           |                                                                | Possible Removal                                     |
| P17      | <i>E. parramattensis</i> subsp. <i>decadens</i>         | 359115                               | 6369318                               | 0.13     | 4           |                                                                | Possible Removal                                     |
| P18      | <i>E. parramattensis</i> subsp. <i>decadens</i>         | 359112                               | 6369314                               | 0.12     | 4           |                                                                | Possible Removal                                     |

| Tree No. | Species                                         | COORDINATES<br>GDA – 2020<br>Easting | COORDINATES<br>GDA – 2020<br>Northing | *DBH (M)  | *Height (M) | Comments<br>Habitat Present | Removal Required? |
|----------|-------------------------------------------------|--------------------------------------|---------------------------------------|-----------|-------------|-----------------------------|-------------------|
| P19      | <i>E. parramattensis</i> subsp. <i>decadens</i> | 359109                               | 6369315                               | 0.13      | 4           |                             | Possible Removal  |
| P20      | <i>E. parramattensis</i> subsp. <i>decadens</i> | 359107                               | 6369316                               | 0.2       | 4           |                             | Possible Removal  |
| P21      | <i>E. parramattensis</i> subsp. <i>decadens</i> | 359106                               | 6369319                               | 0.22 0.16 | 4.5         |                             | Possible Removal  |
| P22      | <i>E. parramattensis</i> subsp. <i>decadens</i> | 359079                               | 6369331                               | 0.25      | 7           |                             | Possible Removal  |
| P23      | <i>E. parramattensis</i> subsp. <i>decadens</i> | 359074                               | 6369337                               | 0.09 0.09 | 4           |                             | Possible Removal  |
| P24      | <i>E. parramattensis</i> subsp. <i>decadens</i> | 359050                               | 6369356                               | 0.1       | 4           |                             | Possible Removal  |
| P25      | <i>E. parramattensis</i> subsp. <i>decadens</i> | 358995                               | 6369387                               | 0.12      | 4.5         |                             | Possible Removal  |
| P26      | <i>E. parramattensis</i> subsp. <i>decadens</i> | 358972                               | 6369410                               | 0.16      | 4.5         |                             | Possible Removal  |
| P27      | <i>E. parramattensis</i> subsp. <i>decadens</i> | 358967                               | 6369403                               | 0.2       | 5           |                             | Possible Removal  |
| P28      | <i>E. parramattensis</i> subsp. <i>decadens</i> | 358964                               | 6369403                               | 0.12      | 4           |                             | Possible Removal  |
| P29      | <i>E. parramattensis</i> subsp. <i>decadens</i> | 358929                               | 6369422                               | 0.3       | 6           |                             | Possible Removal  |
| P30      | <i>E. parramattensis</i> subsp. <i>decadens</i> | 359015                               | 6369379                               | 0.1       | 2           |                             | Possible Removal  |
| P31      | <i>E. parramattensis</i> subsp. <i>decadens</i> | 359161                               | 6369290                               | 0.19      | 5           |                             | Possible Removal  |
| P32      | <i>E. parramattensis</i> subsp. <i>decadens</i> | 359172                               | 6369279                               | 0.2       | 4.5         |                             | Possible Removal  |

**Table F.2** *Diuris tricolor* (Pine Donkey Orchid) recorded within REZ.

| Tree No. | Species                                     | COORDINATES<br>GDA – 2020<br>Easting | COORDINATES<br>GDA – 2020<br>Northing | Comments                                                                                           | Removal Required?                                           |
|----------|---------------------------------------------|--------------------------------------|---------------------------------------|----------------------------------------------------------------------------------------------------|-------------------------------------------------------------|
| D1       | <i>Diuris tricolor</i> (Pine Donkey Orchid) | 305420                               | 6432084                               | Two plants recorded just outside the existing eastern edge of the easement NE of Muswellbrook STS. | Avoid will remain <i>in-situ</i> .<br>Remain <i>in-situ</i> |



**Table F.3** *Eucalyptus glaucina* (Slaty Red Gum) recorded within REZ or within proximity.

| Tree No. | Species                                    | COORDINAT<br>ES<br>GDA – 2020<br>Easting | COORDINATES<br>GDA – 2020<br>Northing | *DBH (M)            | *Height (M) | Comments<br>Habitat Present                                                 | Removal Required?   |
|----------|--------------------------------------------|------------------------------------------|---------------------------------------|---------------------|-------------|-----------------------------------------------------------------------------|---------------------|
| S1       | <i>Eucalyptus glaucina</i> (Slaty Red Gum) | 340654                                   | 6385034                               | 0.19, 0.26          | 10          | Possible specimen                                                           | Near original route |
| S2       | <i>Eucalyptus glaucina</i>                 | 339385                                   | 6385521                               | thin                | 2           | Likely outside easement. Possible specimen                                  |                     |
| S3       | <i>Eucalyptus glaucina</i>                 | 339379                                   | 6385516                               | 0.04                | 3           | Likely outside easement. Previously cut and growing back. Possible specimen |                     |
| S4       | <i>Eucalyptus glaucina</i>                 | 338903                                   | 6385631                               | 0.07                | 4           | Likely outside easement. Possible specimen                                  |                     |
| S5       | <i>Eucalyptus glaucina</i>                 | 338903                                   | 6385631                               | 0.12                | 6           |                                                                             |                     |
| S6       | <i>Eucalyptus glaucina</i>                 | 338782                                   | 6385680                               |                     | 1.5         |                                                                             |                     |
| S7       | <i>Eucalyptus glaucina</i>                 | 338045                                   | 6385766                               |                     |             |                                                                             |                     |
| S8       | <i>Eucalyptus glaucina</i>                 | 333185                                   | 6387116                               | 0.43                | 10          | Possible specimen                                                           |                     |
| S9       | <i>Eucalyptus glaucina</i>                 | 328017                                   | 6387897                               | 0.21, 0.29,<br>0.34 | 8           |                                                                             |                     |
| S10      | <i>Eucalyptus glaucina</i>                 | 327964                                   | 6387876                               | 0.45                | 11          | Possible specimen                                                           |                     |
| S11      | <i>Eucalyptus glaucina</i>                 | 327320                                   | 6387820                               | 0.2                 | 7           | Potential removal. Possible specimen                                        |                     |
| S12      | <i>Eucalyptus glaucina</i>                 | 327321                                   | 6387817                               | 0.18                | 8           | Potential removal. Possible specimen                                        |                     |
| S13      | <i>Eucalyptus glaucina</i>                 | 327322                                   | 6387818                               | 0.22                | 9           | Possible specimen                                                           |                     |

**Table F.4** *Grevillea parviflora* subsp. *parviflora* (Pine Donkey Orchid) recorded within REZ.

| Tree No. | Species                                              | COORDINATES<br>GDA – 2020<br>Easting | COORDINATES<br>GDA – 2020<br>Northing | Comments | Removal Required?                  |
|----------|------------------------------------------------------|--------------------------------------|---------------------------------------|----------|------------------------------------|
| G1       | <i>Grevillea parviflora</i> subsp. <i>parviflora</i> | 357426                               | 6370406                               |          | Avoid will remain <i>in-situ</i> . |
| G2       | <i>Grevillea parviflora</i> subsp. <i>parviflora</i> | 357419                               | 6370423                               |          | Avoid will remain <i>in-situ</i> . |
| G3       | <i>Grevillea parviflora</i> subsp. <i>parviflora</i> | 357308                               | 6370507                               |          | Avoid will remain <i>in-situ</i> . |
| G4       | <i>Grevillea parviflora</i> subsp. <i>parviflora</i> | 357296                               | 6370514                               |          | Avoid will remain <i>in-situ</i> . |
| G5       | <i>Grevillea parviflora</i> subsp. <i>parviflora</i> | 357251                               | 6370546                               |          | Avoid will remain <i>in-situ</i> . |
| G6       | <i>Grevillea parviflora</i> subsp. <i>parviflora</i> | 357283                               | 6370535                               |          | Avoid will remain <i>in-situ</i> . |
| G7       | <i>Grevillea parviflora</i> subsp. <i>parviflora</i> | 357428                               | 6370425                               |          | Avoid will remain <i>in-situ</i> . |

**Table F.5** *Acacia pendula* (Weeping Myall) recorded within REZ or in proximity.

| Tree No. | Species               | COORDINATE S<br>GDA – 2020<br>Easting | COORDINATES<br>GDA – 2020<br>Northing | *DBH (M)   | *Height (M) | Comments<br>Habitat Present                                   | Removal<br>Required? |
|----------|-----------------------|---------------------------------------|---------------------------------------|------------|-------------|---------------------------------------------------------------|----------------------|
| 1        | <i>Acacia pendula</i> | 339978                                | 6385341                               |            |             | Approximately 5 specimens. Outside of easement, Lower Belford | No                   |
| 2        | <i>Acacia pendula</i> | 311984                                | 6418382                               |            |             | Liddell STS Hub at Antiene, within planted treed belt         | No                   |
| 3        | <i>Acacia pendula</i> | 312028                                | 6418360                               |            |             | Liddell STS Hub at Antiene, within planted treed belt         | No                   |
| 4        | <i>Acacia pendula</i> | 312018                                | 6418364                               |            |             | Liddell STS Hub at Antiene, within planted treed belt         | No                   |
| 5        | <i>Acacia pendula</i> | 312004                                | 6418373                               |            |             | Liddell STS Hub at Antiene, within planted treed belt         | No                   |
| 6        | <i>Acacia pendula</i> | 311999                                | 6418376                               |            |             | Liddell STS Hub at Antiene, within planted treed belt         | No                   |
| 7        | <i>Acacia pendula</i> | 311990                                | 6418380                               |            |             | Liddell STS Hub at Antiene, within planted treed belt         | No                   |
| 8        | <i>Acacia pendula</i> | 312149                                | 6418620                               |            |             | Liddell STS Hub at Antiene, within planted treed belt         | Yes                  |
| 9        | <i>Acacia pendula</i> | 312149                                | 6418613                               |            |             | Liddell STS Hub at Antiene, within planted treed belt         | Yes                  |
| 10       | <i>Acacia pendula</i> | 312167                                | 6418602                               |            |             | Liddell STS Hub at Antiene, within planted treed belt         | No                   |
| 11       | <i>Acacia pendula</i> | 312149                                | 6418620                               | 0.15       |             | Liddell STS Hub at Antiene, within planted treed belt         | No                   |
| 12       | <i>Acacia pendula</i> | 312149                                | 6418613                               | 0.18, 0.12 |             | Liddell STS Hub at Antiene, within planted treed belt         | No                   |
| 13       | <i>Acacia pendula</i> | 312167                                | 6418602                               | 0.13       |             | Liddell STS Hub at Antiene, within planted treed belt         | No                   |
| 14       | <i>Acacia pendula</i> | 320735                                | 6392532                               | 0.15       |             | Golden Highway, Warkworth                                     | Possible             |
| 15       | <i>Acacia pendula</i> | 320736                                | 6392531                               |            |             | Golden Highway, Warkworth                                     | Possible             |
| 16       | <i>Acacia pendula</i> | 320722                                | 6392538                               | 0.13       |             | Golden Highway, Warkworth                                     | Possible             |
| 17       | <i>Acacia pendula</i> | 320727                                | 6392538                               | 0.2        |             | Golden Highway, Warkworth                                     | Possible             |
| 18       | <i>Acacia pendula</i> | 320727                                | 6392538                               |            |             | Golden Highway, Warkworth                                     | Possible             |
| 19       | <i>Acacia pendula</i> | 320727                                | 6392553                               |            |             | Golden Highway, Warkworth                                     | Possible             |

## **APPENDIX G**

### **FAUNA SPECIES LIST**



## VERTEBRATE FAUNA LIST

Family sequencing and taxonomy follow for each fauna class:

### Fish

Allen, G.R., Midgley, S.H. & Allen, M. (2002). *Field Guide to the Freshwater Fishes of Australia*. Western Australian Museum, Perth.

### Herpetofauna

Cogger, H.G. (2014). *Reptiles and Amphibians of Australia* (7th edn.). CSIRO Publishing.

### Birds

Pizzey and Knight (2012) (9th edn).

### Mammals

Van Dyck, S. and Strahan, R. (Ed) (2008). *The Mammals of Australia* (3rd edn). New Holland Publishers, Australia –

Churchill, S. (2008). *Australian Bats*. (2nd edn.). Allen & Unwin Australia.

(?) - Indicates a species identified without certainty or to a Genus level only.

\* - Indicates an introduced species.

Threatened species addressed within this assessment appear in **bold** font.

Introduced species are indicated by an asterisk (“\*”).

**The following standard abbreviations are used to indicate subspecific taxa:**

**subsp.** -subspecies

**var.-** variety

**x -** hybrid between the two indicated species

### **Biodiversity Conservation Act 2016 (BC Act)**

**V** Vulnerable

**E1** Endangered

**E2** Endangered Population

**E4A** Critically Endangered Population

### **Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act)**

**V** **Vulnerable**

**E** **Endangered**

**CE** **Critically Endangered Population**

**M** **Migratory**

### **Regionally Significant Fauna Species.**

**+** Region includes Gosford, Wyong, Cessnock, Maitland, Lake Macquarie, Newcastle and Port Stephens LGA's. Produced from Stage 1 of the LHCCREMS – Regional Biodiversity Conservation Strategy.

### **Observation Type**

**O** - Observed (sighted)

**W** - Heard call

**OW** – Observed and heard call

**X** - In scat

**P** – Scat

**T** - Trapped or netted

**H** – Hair, feathers or skin

**A** - Stranded/Beached

**G** – Crushed cones

**R** – Road Kill

**D** – Dog Kill

**Q** – Camera

**C** – Cat Kill

**V** – Fox Kill

**K** – Dead

**S** – Shot

**I** – Fossil/subfossil

**FB** – Burrow

**F** – Tracks, scratching

**Z** – In raptor/owl Pellet

**U** – Ultrasonic recording

**M** - Miscellaneous

**E** – Nest/roost

**B** - Burnt

**Y** – Bones, teeth or shell

**N** – Not located

**AR** – Acoustic Recording

Table F1 Total Fauna List

| SCIENTIFIC NAME                                 | COMMON NAME            | BC ACT | EPBC ACT | REGIONALLY SIGNIFICANT | OBSERVATION TYPE |
|-------------------------------------------------|------------------------|--------|----------|------------------------|------------------|
| <b>Phylum - Chordata</b>                        |                        |        |          |                        |                  |
| <b>Subphylum - Vertebrata</b>                   |                        |        |          |                        |                  |
|                                                 |                        |        |          |                        |                  |
| <b>Class - Actinopterygii</b>                   |                        |        |          |                        |                  |
|                                                 |                        |        |          |                        |                  |
| <b>Order Cyprinodontiformes</b>                 |                        |        |          |                        |                  |
|                                                 |                        |        |          |                        |                  |
| <b>Family Poeciliidae</b>                       |                        |        |          |                        |                  |
| <i>*Gambusia holbrooki</i>                      | Mosquitofish           |        |          |                        |                  |
|                                                 |                        |        |          |                        |                  |
| <b>Class Amphibia - Amphibians</b>              |                        |        |          |                        |                  |
|                                                 |                        |        |          |                        |                  |
| <b>Order Salientia - Frogs</b>                  |                        |        |          |                        |                  |
|                                                 |                        |        |          |                        |                  |
| <b>Family Myobatrachidae - 'Southern Frogs'</b> |                        |        |          |                        |                  |
| <i>Crinia signifera</i>                         | Common Eastern Froglet |        |          |                        |                  |
| <i>Limnodynastes peronii</i>                    | Striped Marsh Frog     |        |          |                        |                  |
| <i>Limnodynastes tasmaniensis</i>               | Spotted Grass Frog     |        |          | +                      |                  |
|                                                 |                        |        |          |                        |                  |
| <b>Family Hylidae - Tree Frogs</b>              |                        |        |          |                        |                  |
| <i>Litoria peronii</i>                          | Peron's Tree Frog      |        |          |                        |                  |
|                                                 |                        |        |          |                        |                  |
| <b>Class Reptilia - Reptiles</b>                |                        |        |          |                        |                  |
|                                                 |                        |        |          |                        |                  |
| <b>Order Squamata – Lizards and Snakes</b>      |                        |        |          |                        |                  |
| <b>Suborder Sauria - Lizards</b>                |                        |        |          |                        |                  |
| <b>Family Agamidae - Dragons</b>                |                        |        |          |                        |                  |
| <i>Pogona barbata</i>                           | Eastern Bearded Dragon |        |          | +                      |                  |

| SCIENTIFIC NAME                                 | COMMON NAME             | BC ACT | EPBC ACT | REGIONALLY SIGNIFICANT | OBSERVATION TYPE |
|-------------------------------------------------|-------------------------|--------|----------|------------------------|------------------|
| <b>Family Varanidae - Monitors</b>              |                         |        |          |                        |                  |
| <i>Varanus varius</i>                           | Lace Monitor            |        |          |                        |                  |
| <b>Family Scinidae - Skinks</b>                 |                         |        |          |                        |                  |
| <i>Anomalopus leuckartii</i>                    | Two-clawed worm-skink   |        |          |                        |                  |
| <i>Carlia tetradactyla</i>                      | Rainbow Skink           |        |          | +                      |                  |
| <i>Cryptoblepharus pulcher</i>                  | Fence Skink, Wall Skink |        |          |                        |                  |
| <i>Ctenotus robustus</i>                        | Striped Skink           |        |          |                        |                  |
| <i>Egernia striolata</i>                        | Tree Skink              |        |          | +                      |                  |
| <i>Eulamprus tenuis</i>                         | Bar-sided Skink         |        |          | +                      |                  |
| <i>Lampropholis delicata</i>                    | Grass Skink             |        |          |                        |                  |
| <b>Suborder Serpentes - Snakes</b>              |                         |        |          |                        |                  |
| <b>Family Elapidae - Venomous Snakes</b>        |                         |        |          |                        |                  |
| <i>Pseudechis porphyriacus</i>                  | Red-bellied Black Snake |        |          |                        |                  |
| <i>Pseudonaja textilis</i>                      | Eastern Brown Snake     |        |          | +                      |                  |
| <b>Class Aves - Birds</b>                       |                         |        |          |                        |                  |
| <b>Family Phasianidae</b>                       |                         |        |          |                        |                  |
| <i>Coturnix pectoralis</i>                      | Stubble Quail           |        |          |                        |                  |
| <b>Family Anatidae - Ducks, Swans and Geese</b> |                         |        |          |                        |                  |
| <i>Anas superciliosa</i>                        | Pacific Black Duck      |        |          |                        |                  |
| <i>Chenonetta jubata</i>                        | Australian Wood Duck    |        |          |                        |                  |
| <b>Family Podicipedidae - Grebes</b>            |                         |        |          |                        |                  |
| <i>Tachybaptus novaehollandiae</i>              | Australian Grebe        |        |          |                        |                  |
| <b>Family Columbidae - Pigeons, Doves</b>       |                         |        |          |                        |                  |
| <i>Geopelia humeralis</i>                       | Bar-shouldered Dove     |        |          |                        |                  |
| <i>Leucosarcia melanoleuca</i>                  | Wonga Pigeon            |        |          |                        |                  |



| SCIENTIFIC NAME                                                 | COMMON NAME                    | BC ACT | EPBC ACT | REGIONALLY SIGNIFICANT | OBSERVATION TYPE |
|-----------------------------------------------------------------|--------------------------------|--------|----------|------------------------|------------------|
| <i>Ocyphaps lophotes</i>                                        | Crested Pigeon                 |        |          |                        |                  |
| <i>Phaps chalcoptera</i>                                        | Common Bronzewing              |        |          |                        |                  |
|                                                                 |                                |        |          |                        |                  |
| <b>Family Phalacrocoridae - Cormorants</b>                      |                                |        |          |                        |                  |
| <i>Phalacrocorax fuscescens</i>                                 | Pied Cormorant                 |        |          |                        |                  |
| <i>Phalacrocorax sulcirostris</i>                               | Little Black Cormorant         |        |          |                        |                  |
| <i>Phalacrocorax varius</i>                                     | Little Pied Cormorant          |        |          |                        |                  |
|                                                                 |                                |        |          |                        |                  |
| <b>Family Ardeidae - Herons, Egrets and Bitterns</b>            |                                |        |          |                        |                  |
| <i>Ardea alba</i>                                               | Great Egret                    |        |          |                        |                  |
| <i>Ardea ibis</i>                                               | Cattle Egret                   |        | M        |                        |                  |
| <i>Ardea pacifica</i>                                           | White-necked Heron             |        |          |                        |                  |
| <i>Egretta novaehollandiae</i>                                  | White-faced Heron              |        |          |                        |                  |
|                                                                 |                                |        |          |                        |                  |
| <b>Family Threskiornithidae - Ibises and Spoonbills</b>         |                                |        |          |                        |                  |
| <i>Threskiornis molucca</i>                                     | Australian White Ibis          |        |          |                        |                  |
| <i>Threskiornis spinicollis</i>                                 | Straw-necked Ibis              |        |          |                        |                  |
|                                                                 |                                |        |          |                        |                  |
| <b>Family Accipitridae - Osprey, Hawks, Eagles and Harriers</b> |                                |        |          |                        |                  |
| <i>Accipiter fasciatus</i>                                      | Brown Goshawk                  |        |          |                        |                  |
| <i>Accipiter cirrhocephalus</i>                                 | Collared Sparrowhawk           |        |          |                        |                  |
| <i>Aquila audax</i>                                             | Wedge-tailed Eagle             |        |          |                        |                  |
| <i>Elanus axillaris</i>                                         | Black-shouldered Kite          |        |          |                        |                  |
| <b><i>Haliaeetus leucogaster</i></b>                            | <b>White-bellied Sea-Eagle</b> | V      | M        | +                      |                  |
| <i>Haliastur sphenurus</i>                                      | Whistling Kite                 |        |          |                        |                  |
|                                                                 |                                |        |          |                        |                  |
| <b>Family Falconidae - Falcons</b>                              |                                |        |          |                        |                  |
| <i>Falco cenchroides</i>                                        | Nankeen Kestrel                |        |          |                        |                  |
| <i>Falco longipennis</i>                                        | Australian Hobby               |        |          |                        |                  |
| <i>Falco peregrinus</i>                                         | Peregrine Falcon               |        |          | +                      |                  |
|                                                                 |                                |        |          |                        |                  |

| SCIENTIFIC NAME                                             | COMMON NAME                  | BC ACT | EPBC ACT | REGIONALLY SIGNIFICANT | OBSERVATION TYPE |
|-------------------------------------------------------------|------------------------------|--------|----------|------------------------|------------------|
| <b>Family Rallidae</b>                                      |                              |        |          |                        |                  |
| <i>Fulica atra</i>                                          | Eurasian Coot                |        |          |                        |                  |
| <i>Gallinula tenebrosa</i>                                  | Dusky Moorhen                |        |          |                        |                  |
| <i>Porphyrio porphyrio</i>                                  | Purple Swamp Hen             |        |          |                        |                  |
|                                                             |                              |        |          |                        |                  |
| <b>Family Charadriidae Plover, Dotterels, Lapwings</b>      |                              |        |          |                        |                  |
| <i>Vanellus miles</i>                                       | Masked Lapwing               |        |          |                        |                  |
|                                                             |                              |        |          |                        |                  |
| <b>Family Cacatuidae - Cockatoos and Corellas</b>           |                              |        |          |                        |                  |
| <i>Cacatua galerita</i>                                     | Sulphur-crested Cockatoo     |        |          |                        |                  |
| <i>Cacatua roseicapilla</i>                                 | Galah                        |        |          |                        |                  |
| <i>Cacatua sanguinea</i>                                    | Little Corella               |        |          |                        |                  |
| <i>Calyptorhynchus funereus</i>                             | Yellow-tailed Black-Cockatoo |        |          | +                      |                  |
|                                                             |                              |        |          |                        |                  |
| <b>Family Psittacidae - Parrots, Rosellas and Lorikeets</b> |                              |        |          |                        |                  |
| <i>Alisterus scapularis</i>                                 | King Parrot                  |        |          |                        |                  |
| <i>Glossopsitta concinna</i>                                | Musk Lorikeet                |        |          |                        |                  |
| <i>Neophema pulchella</i>                                   | Turquoise Parrot             | V      |          |                        |                  |
| <i>Platycercus elegans</i>                                  | Crimson Rosella              |        |          |                        |                  |
| <i>Platycercus eximius</i>                                  | Eastern Rosella              |        |          |                        |                  |
| <i>Psephotus haematonotus</i>                               | Red-rumped Parrot            |        |          |                        |                  |
| <i>Trichoglossus chlorolepidotus</i>                        | Scaly-breasted Lorikeet      |        |          |                        |                  |
| <i>Trichoglossus haematodus</i>                             | Rainbow Lorikeet             |        |          |                        |                  |
|                                                             |                              |        |          |                        |                  |
| <b>Family Cuculidae - Cuckoos</b>                           |                              |        |          |                        |                  |
| <i>Cacomantis flabelliformis</i>                            | Fan-tailed Cuckoo            |        |          |                        |                  |
| <i>Cacomantis variolosus</i>                                | Brush Cuckoo                 |        |          |                        |                  |
| <i>Eudynamis orientalis</i>                                 | Common Koel                  |        |          |                        |                  |
| <i>Scythrops novaehollandiae</i>                            | Channel-billed Cuckoo        |        |          |                        |                  |
|                                                             |                              |        |          |                        |                  |
| <b>Family Halcyonidae - Tree Kingfishers</b>                |                              |        |          |                        |                  |
| <i>Dacelo novaeguineae</i>                                  | Laughing Kookaburra          |        |          |                        |                  |

| SCIENTIFIC NAME                                                                           | COMMON NAME                | BC ACT   | EPBC ACT | REGIONALLY SIGNIFICANT | OBSERVATION TYPE |
|-------------------------------------------------------------------------------------------|----------------------------|----------|----------|------------------------|------------------|
| <i>Todiramphus sanctus</i>                                                                | Sacred Kingfisher          |          |          |                        |                  |
| <b>Family Climacteridae - Treecreepers</b>                                                |                            |          |          |                        |                  |
| <i>Climacteris erythrops</i>                                                              | Red-browed Treecreeper     |          |          | +                      |                  |
| <i>Climacteris picumnus victoriae</i>                                                     | <b>Brown Treecreeper</b>   | <b>V</b> |          | +                      |                  |
| <i>Cormobates leucophaea</i>                                                              | White-throated Treecreeper |          |          |                        |                  |
| <b>Family Maluridae</b>                                                                   |                            |          |          |                        |                  |
| <i>Malurus cyaneus</i>                                                                    | Superb Fairy-wren          |          |          |                        |                  |
| <i>Malurus lamberti</i>                                                                   | Variegated Fairy-wren      |          |          |                        |                  |
| <b>Family Pardalotidae - Pardalotes, Gerygones, Scrubwrens, Heathwrens and Thornbills</b> |                            |          |          |                        |                  |
| <i>Acanthiza chrysorrhoa</i>                                                              | Yellow-rumped Thornbill    |          |          |                        |                  |
| <i>Acanthiza nana</i>                                                                     | Yellow Thornbill           |          |          |                        |                  |
| <i>Acanthiza pusilla</i>                                                                  | Brown Thornbill            |          |          |                        |                  |
| <i>Gerygone mouki</i>                                                                     | Brown Gerygone             |          |          |                        |                  |
| <i>Gerygone olivacea</i>                                                                  | White-throated Gerygone    |          |          |                        |                  |
| <i>Pardalotus punctatus</i>                                                               | Spotted Pardalote          |          |          |                        |                  |
| <i>Pardalotus striatus</i>                                                                | Striated Pardalote         |          |          |                        |                  |
| <b>Family Meliphagidae - Honeyeaters</b>                                                  |                            |          |          |                        |                  |
| <i>Acanthorhynchus tenuirostris</i>                                                       | Eastern Spinebill          |          |          |                        |                  |
| <i>Anthochaera carunculata</i>                                                            | Red Wattlebird             |          |          |                        |                  |
| <i>Caligavis chrysops</i>                                                                 | Yellow-faced Honeyeater    |          |          |                        |                  |
| <i>Entomyzon cyanotis</i>                                                                 | Blue-faced Honeyeater      |          |          |                        |                  |
| <i>Lichenostomus fuscus</i>                                                               | Fuscous Honeyeater         |          |          |                        |                  |
| <i>Lichenostomus melanops</i>                                                             | Yellow-tuft Honeyeater     |          |          |                        |                  |
| <i>Manorina melanocephala</i>                                                             | Noisy Miner                |          |          |                        |                  |
| <i>Philemon corniculatus</i>                                                              | Noisy Friarbird            |          |          |                        |                  |



| SCIENTIFIC NAME                                                           | COMMON NAME               | BC ACT | EPBC ACT | REGIONALLY SIGNIFICANT | OBSERVATION TYPE |
|---------------------------------------------------------------------------|---------------------------|--------|----------|------------------------|------------------|
| <b>Family Pomatostomidae Australian Babblers</b>                          |                           |        |          |                        |                  |
| <i>Pomatostomus temporalis temporalis</i>                                 | Grey-crowned Babbler      | V      |          | +                      |                  |
| <b>Family Petroicidae - Robins and Jacky Winter</b>                       |                           |        |          |                        |                  |
| <i>Eopsaltria australis</i>                                               | Eastern Yellow Robin      |        |          |                        |                  |
| <i>Microeca fascians</i>                                                  | Jacky Winter              |        |          |                        |                  |
| <b>Family Pachycephalidae - Whistlers, Shrike-tit and Shrike-thrushes</b> |                           |        |          |                        |                  |
| <i>Colluricincla harmonica</i>                                            | Grey Shrike-thrush        |        |          |                        |                  |
| <i>Pachycephala pectoralis</i>                                            | Golden Whistler           |        |          |                        |                  |
| <i>Pachycephala rufiventris</i>                                           | Rufous Whistler           |        |          |                        |                  |
| <b>Family Monarchidae - Monarchs, Flycatchers and Magpie-Lark</b>         |                           |        |          |                        |                  |
| <i>Grallina cyanoleuca</i>                                                | Magpie-lark               |        |          |                        |                  |
| <b>Family Rhipiduridae - Fantails</b>                                     |                           |        |          |                        |                  |
| <i>Rhipidura albiscapa</i> syn. <i>Rhipidura fuliginosa</i>               | Grey Fantail              |        |          |                        |                  |
| <i>Rhipidura leucophrys</i>                                               | Willie Wagtail            |        |          |                        |                  |
| <b>Family Campephagidae - Cuckoo-shrikes and Trillers</b>                 |                           |        |          |                        |                  |
| <i>Coracina novaehollandiae</i>                                           | Black-faced Cuckoo-shrike |        |          |                        |                  |
| <b>Family Hirundinidae - Swallows and Martins</b>                         |                           |        |          |                        |                  |
| <i>Cecropis nigricans</i>                                                 | Tree Martin               |        |          |                        |                  |
| <i>Hirundo neoxena</i>                                                    | Welcome Swallow           |        |          |                        |                  |
| <b>Family Sylviidae - Old World Warblers</b>                              |                           |        |          |                        |                  |
| <i>Acrocephalus stentoreus</i>                                            | Clamorous Reed Warbler    |        |          |                        |                  |

| SCIENTIFIC NAME                                                              | COMMON NAME             | BC ACT | EPBC ACT | REGIONALLY SIGNIFICANT | OBSERVATION TYPE |
|------------------------------------------------------------------------------|-------------------------|--------|----------|------------------------|------------------|
| <i>Cisticola exilis</i>                                                      | Golden-headed Cisticola |        |          |                        |                  |
| <b>Family Zosteropidae - White-eyes</b>                                      |                         |        |          |                        |                  |
| <i>Zosterops lateralis</i>                                                   | Silvereye               |        |          |                        |                  |
| <b>Family Artamidae - Wood-swallows, Butcherbirds, Magpie and Currawongs</b> |                         |        |          |                        |                  |
| <i>Cracticus nigrogularis</i>                                                | Pied Butcherbird        |        |          |                        |                  |
| <i>Cracticus tibicen</i> syn. <i>Gymnorhina tibicen</i>                      | Australian Magpie       |        |          |                        |                  |
| <i>Strepera graculina</i>                                                    | Pied Currawong          |        |          |                        |                  |
| <b>Family Corvidae - Crows, Raven</b>                                        |                         |        |          |                        |                  |
| <i>Corvus coronoides</i>                                                     | Australian Raven        |        |          |                        |                  |
| <b>Corcoracidae - Chough and Apostlebird</b>                                 |                         |        |          |                        |                  |
| <i>Corcorax melanorhamphos</i>                                               | White-winged Chough     |        |          |                        |                  |
| <b>Family Estrildidae - Grassfinches</b>                                     |                         |        |          |                        |                  |
| <i>Neochima temporalis</i>                                                   | Red-browed Finch        |        |          |                        |                  |
| <i>Taeniopygia bichenovii</i>                                                | Double-barred Finch     |        |          |                        |                  |
| <b>Family Dicaeidae</b>                                                      |                         |        |          |                        |                  |
| <i>Dicaeum hirundinaceum</i>                                                 | Mistletoe Bird          |        |          |                        |                  |
| <b>Family Turdidae - Thrushes</b>                                            |                         |        |          |                        |                  |
| * <i>Turdus merula</i>                                                       | Common Blackbird        |        |          |                        |                  |
| <b>Family Sturnidae - Starlings and Mynas</b>                                |                         |        |          |                        |                  |
| * <i>Sturnus tristis</i> syn <i>Acridotheres tristis</i>                     | Indian Myna             |        |          |                        |                  |
| * <i>Sturnus vulgaris</i>                                                    | Common Starling         |        |          |                        |                  |
| <b>Class Mammalia - Mammals</b>                                              |                         |        |          |                        |                  |
| <b>Subclass Prototheria - Monotremes</b>                                     |                         |        |          |                        |                  |

| SCIENTIFIC NAME                                                  | COMMON NAME           | BC ACT | EPBC ACT | REGIONALLY SIGNIFICANT | OBSERVATION TYPE |
|------------------------------------------------------------------|-----------------------|--------|----------|------------------------|------------------|
| <b>Order Monotremata</b>                                         |                       |        |          |                        |                  |
| <b>Family Tachyglossidae - Echidna</b>                           |                       |        |          |                        |                  |
| <i>Tachyglossus aculeatus</i>                                    | Echidna               |        |          | +                      |                  |
| <b>Subclass Marsupialia - Marsupials</b>                         |                       |        |          |                        |                  |
| <b>Order Dasyuromorphia – Carnivorous Marsupials</b>             |                       |        |          |                        |                  |
| <b>Order Diprotodontia</b>                                       |                       |        |          |                        |                  |
| <b>Suborder Vombatiformes</b>                                    |                       |        |          |                        |                  |
| <b>Family Vombatidae - Wombats</b>                               |                       |        |          |                        |                  |
| <i>Vombatus ursinus</i>                                          | Common Wombat         |        |          |                        |                  |
| <b>Superfamily - Macropodoidae</b>                               |                       |        |          |                        |                  |
| <b>Family Macropodidae - Kangaroos, Wallabies</b>                |                       |        |          |                        |                  |
| <i>Macropus giganteus</i>                                        | Eastern Grey Kangaroo |        |          | +                      |                  |
| <i>Notamacropus rufogriseus</i> syn. <i>Macropus rufogriseus</i> | Red-necked Wallaby    |        |          | +                      |                  |
| <b>Subclass Eutheria - Eutherian Mammals</b>                     |                       |        |          |                        |                  |
| <b>Order Rodentia</b>                                            |                       |        |          |                        |                  |
| <b>Family Muridae - Rodents</b>                                  |                       |        |          |                        |                  |
| * <i>Mus musculus</i>                                            | House Mouse           |        |          |                        |                  |
| <b>Order Lagomorpha</b>                                          |                       |        |          |                        |                  |



| SCIENTIFIC NAME                | COMMON NAME     | BC ACT | EPBC ACT | REGIONALLY SIGNIFICANT | OBSERVATION TYPE |
|--------------------------------|-----------------|--------|----------|------------------------|------------------|
| <b>Family Leporidae</b>        |                 |        |          |                        |                  |
| * <i>Lepus capensis</i>        | European Hare   |        |          |                        |                  |
| * <i>Oryctolagus cuniculus</i> | European Rabbit |        |          | O                      |                  |
|                                |                 |        |          |                        |                  |
| <b>Order Carnivora</b>         |                 |        |          |                        |                  |
|                                |                 |        |          |                        |                  |
| <b>Family Canidae</b>          |                 |        |          |                        |                  |
| * <i>Vulpes vulpes</i>         | Red Fox         |        |          |                        |                  |
|                                |                 |        |          |                        |                  |
| <b>Order Perissodactyla</b>    |                 |        |          |                        |                  |
| <b>Family Equidae</b>          |                 |        |          |                        |                  |
| * <i>Equus caballus</i>        | Horse           |        |          |                        |                  |
|                                |                 |        |          |                        |                  |
| <b>Order Artiodactyla</b>      |                 |        |          |                        |                  |
| <b>Family Suidae</b>           |                 |        |          |                        |                  |
| * <i>Sus scrofa</i>            | Pig             |        |          |                        |                  |
|                                |                 |        |          |                        |                  |
| <b>Family Bovidae</b>          |                 |        |          |                        |                  |
| * <i>Bos taurus</i>            | Cattle          |        |          |                        |                  |
|                                |                 |        |          |                        |                  |
| <b>Family Cervidae</b>         |                 |        |          |                        |                  |
| * <i>Cervus elaphus</i>        | Red Deer        |        |          |                        |                  |
|                                |                 |        |          |                        |                  |